

Homework 2

1. Give Kripke counter-models to:

(a) $\neg(p \wedge \neg q) \rightarrow (p \rightarrow q)$ [2pts]

(b) $\neg\neg(p \vee q) \rightarrow \neg\neg p \vee \neg\neg q$ [2pts]

(c) $\neg\neg p \vee (\neg\neg p \rightarrow p)$ [2pts]

Proofs are not needed.

2. Exercise 4 of the syllabus, on p 16:

Prove that persistency transfers to formulas (i.e., if $w \models \phi$ and wRv then $v \models \phi$, for all propositional formulas ϕ). [4pts]

3. Show directly, **without** using Theorem 30 ($\vdash_{\text{CPC}} \phi$ iff $\vdash_{\text{IPC}} \phi^n$), that $((\varphi \rightarrow \chi) \wedge (\psi \rightarrow \chi) \rightarrow (\varphi \vee \psi \rightarrow \chi))^n$ is provable in **IPC**.

You **are** allowed to use the following fact: $\vdash_{\text{IPC}} \varphi^n \leftrightarrow \neg\neg\varphi^n$. [4pts]

4.* **Definition:**

- φ is **negative** iff there is some ψ such that $\vdash_{\text{IPC}} \varphi \leftrightarrow \neg\psi$
- φ has the **down property** iff for each w which is not an end-point, if for all x with wRx and $w \neq x$ we have $x \models \varphi$, then $w \models \varphi$.

Show that φ is negative iff it has the down property. [4pts]