

Dick de Jongh
Institute for Logic, Language and Computation
University of Amsterdam
dickdij@science.uva.nl

Intuitionistic and Modal Logic

- Universal models for modal logic, application to unifiability
- Universal models for IPC, Jančov formulas, intermediate logics
- Heyting algebras
- Formulas of one variable
- Modal Logic, Translations
- Kripke models, Metatheorems
- Proofs and proof systems (IPC)
- Introduction to intuitionism

Overview

- euclidean geometry,
- Brouwer, Foundations of analysis (Cauchy, Weierstrass), non-
- Brouwer (1881-1963)
- 1997.
- A. Chagrov and M. Zakharyaschev, Modal Logic, Oxford University Press,
- Series DS-2006-02
- See also Nick Bezhanishvili, Lattices of Intermediate and Cylindric Modal Logics, Dissertation, Universiteit van Amsterdam, ILLC Dissertation Series DS-2006-02
- Prepublications PP-2006-25, www.illc.uva.nl
- First three lectures partly based on Intuitionistic Logic by Nick Bezhanishvili and Dick de Jongh, Lecture Notes for ESSLI 2005, ILLC-

Introduction, Brouwer

- **Dissertation** 1907: *Over de Grondslagen van de Wijskunde* (About the Foundations of Mathematics).

- Period that Foundations of Mathematics was hot issue.
- Frege 1879 BegriFFschrift, 1884 Grundlagen der Mathematik, 1903 Grundgesetze der Arithmetik,
- 1901: Russells paradox, Russell 1903, Principles of Mathematics,
- Hilbert 1889 Grundlagen der Geometrie, 1900 Mathematische Probleme, 1904 Über die Grundlagen der Logik und der Arithmetik,
- Cantor . . . , Peano, Schröder, Huntington, Veblen, . . .

Foundations of mathematics

- Predecessors of Brouwer: Kronecker: „God made the natural numbers, the rest is human work”, French semi-intuitionists (e.g. E. Borel).
- Unhappy feeling from these mathematicians about abstractness of mathematics, proving the existence of objects by reasoning by contradiction, so that no object really arises from the proof: $\neg \forall x \neg Ax \leftarrow \exists x Ax$.

Preursors of Intuitionism

- Intuitionism is as one of the three basic points of view opposed to Platonism and formalism. View that mathematics and mathematical truths are creations of the human mind: true = provable. N.B! provable in the informal, not formal sense.
- Intuitionism, Platonism, Formalism
- Intuitionism. Most famous modern representatives: Frege, Gödel. View that mathematical objects have independent existence outside of space-time, that mathematical truths are independent of us. At the time mixed with Logicism, Frege's idea that mathematics is no more than logic, since mathematics can be reduced to it, a view supported by Russell (not a Platonist) at the time.
- Formalism. Most famous modern representative: Hilbert. View that there are no mathematical objects, no mathematical truths, just formal systems and derivations in them.

- Principle of the excluded third (law of the exclude middle) $A \vee \neg A$.
- Criticism of 'classical' logical laws,
- Mathematics is a mental activity, the "exact part of human thought", writing mathematics down is only an aide,
- Logic follows mathematics, is not its basis, logical rules extracted from mathematics,
- Foundations unnecessary, in fact impossible,

Brouwer's ideas

- Brouwer's programme: rebuilding of mathematics according to intuitionistic principles.
- Only partially successful. Not accepted by mathematicians in practice.
- But study of intuitionistic proofs and formal systems very alive. Only by fully accepting intuitionistic methods does one get proofs that guarantees to exhibit objects that are proved to exist. One gets this way the constructive part of mathematics.
- Less popular but fascinating are Brouwer's choice sequences which have classically inconsistent properties.
- Our course will mostly concentrate on propositional calculus.

Brouwer's programme

- Theorem There exist irrational numbers r and s such that r_s is rational.
- Proof Well-known since Euclid, $\sqrt{2}$ is irrational.
- Now either $\sqrt{2}\sqrt{2}$ is rational or it is not.
- In the first case take $r = \sqrt{2}$, $s = \sqrt{2}$. Then $r_s = (\sqrt{2}\sqrt{2})\sqrt{2} = \sqrt{2}^2 = 2$, i.e. rational.
- In the second case take $r = \sqrt{2}$, $s = \sqrt{2}$. Then $r_s = 2$, i.e. rational.
- So, we have found r and s as required, only we cannot tell r is, it is either $\sqrt{2}\sqrt{2}$ or $\sqrt{2}$ (in reality of course the latter) and $= \sqrt{2}$.

Example of nonconstructive proof

- Heyting, 1928-1930:
 - Earlier incomplete version in Kolmogorov 1925,
 - Hilbert type system. We first give natural deduction variant of which first version was given by Gentzen.
 - $\neg\phi$ is defined as $\phi \rightarrow \perp$ where \perp stands for a contradiction, an obviously false statement like $\bot = 0$.

Heyting

Natural Deduction

$\frac{d}{D(x) \in x \in}$	$(x) \in x \in \frac{}{E(t)}$	E
\vdots		
$(c) \in$		
$\frac{(t)\phi}{(x)\phi x A}$	$\frac{(x)\phi x A}{(x)\phi}$	A
χ $\frac{\chi \chi}{\chi \chi} \quad \phi \wedge \phi$ $\vdots \quad \vdots$ $\phi \quad \phi$	$\frac{\phi \wedge \phi}{\phi} \quad \frac{\phi \wedge \phi}{\phi}$	\wedge
$\frac{\phi \quad \phi}{\phi \vee \phi \quad \phi \vee \phi}$	$\frac{\phi \vee \phi}{\phi} \quad \frac{\phi \vee \phi}{\phi}$	\vee
$\frac{\phi}{\phi \leftarrow \phi \quad \phi}$	$\frac{\phi \leftarrow \phi}{\phi}$ \vdots $[\phi]$	\leftarrow
$\frac{\phi}{T}$	none	T
introduction	elimination	

$\phi \vdash$ $\overline{\top}$

⋮

 $\phi \leftarrow \top$

To get classical logic one adds the rule that if \perp is derived from ϕ , then one can conclude to ϕ dropping the assumption $\perp\phi$.

Classical Logic

- Brouwer-Heyting-Kolmogorov interpretation of connectives and quantifiers.
- Natural deduction closely related to BHK.
- Interpretation by means of proofs (nonformal, nonsyntactical objects, mind constructions),
- A proof of $\phi \vee \psi$ consists of proof of ϕ or of proof of ψ (plus conclusion),
- A proof of $\phi \wedge \psi$ consists of proof of ϕ plus proof of ψ (plus conclusion),
- A proof of $\phi \rightarrow \psi$ consists of method that applied to any conceivable proof of ϕ will deliver proof of ψ ,

BHK-interpretation

- A proof of $\forall x \phi(x)$ consists of method that applied to any element d of domain will deliver proof of $\phi(d)$,
- A proof of $\exists x \phi(x)$ consists of object d from domain plus proof of $\phi(d)$ (plus conclusion),
- Proof of $\neg\phi$ is method that given any proof of ϕ gives proof of \perp ,
- Nothing is a proof of \perp ,

BHK-interpretation, continued

- $\vdash (\phi \wedge \psi) \leftarrow (\phi \leftarrow \phi) \wedge (\psi \leftarrow \psi)$ •
- $\vdash ((\chi \leftarrow \phi) \wedge (\phi \leftarrow \phi)) \leftarrow (\chi \wedge \phi \leftarrow \phi)$ •
- $\vdash (\phi \vdash \psi) \wedge (\psi \vdash \phi) \leftarrow (\phi \vee \psi) \vdash$ •
- (excluded middle)
 - other examples of such invalid formulas are $\phi \vee \neg \phi$, (the law of the excluded middle)
- $\vdash (\phi \vee \psi) \vdash \leftarrow (\phi \vdash \phi) \wedge (\psi \vdash \psi)$ •
- $\vdash (\phi \wedge \psi) \vdash \leftarrow (\phi \vdash \phi) \wedge (\psi \vdash \psi)$ •
- $\vdash (\phi \vee \psi) \vdash \leftarrow \neg \phi \vee \neg \psi$ •
- Morgan laws only $\vdash (\phi \wedge \psi) \leftarrow \neg \phi \vee \neg \psi$ is not valid,
- A disjunction is hard to prove: e.g. of the four directions of the de Morgan laws only $\vdash (\phi \wedge \psi) \leftarrow \neg \phi \vee \neg \psi$ is not valid,

Valid and invalid reasoning

- $\cdot ((\phi \wedge \psi) \vee (\phi \wedge \psi)) \text{ (needs } ex \text{ falso!)}.$
- $\cdot \chi \leftarrow (\phi \vee \psi) \leftrightarrow ((\chi \leftarrow \phi) \leftarrow \psi)$
- $\cdot ((\chi \leftarrow \phi) \wedge \psi) \leftrightarrow (\chi \leftarrow \phi) \vee (\chi \leftarrow \psi)$
- $\cdot (\chi \leftarrow \phi) \vee (\phi \leftarrow \psi) \leftrightarrow (\chi \vee \phi \leftarrow \psi)$
- On the other hand, many basic laws naturally remain valid, commutativity and associativity of conjunction and disjunction, both distributivity laws,
- statements directly based on the two-valuedness of truth values are not valid, e.g. $\neg \neg \phi \leftarrow \phi$ or $((\phi \leftarrow \psi) \leftarrow \psi) \leftarrow \phi$ (Peirce's law),
- of the four directions of the classically valid interactions between negations and quantifiers only $\neg \forall x \phi \leftarrow \exists x \neg \phi$ is not valid,
- An existential statement is hard to prove:

Valid and invalid reasoning, continued

$$\begin{array}{c}
 , \phi \leftarrow \top \bullet \\
 , ((\chi \leftarrow \phi \wedge \phi) \leftarrow (\chi \leftarrow \phi)) \leftarrow (\chi \leftarrow \phi) \bullet \\
 , \phi \wedge \phi \leftarrow \phi \quad \phi \wedge \phi \leftarrow \phi \bullet \\
 , (\phi \vee \phi \leftarrow \phi) \leftarrow \phi \bullet \\
 , \phi \leftarrow \phi \vee \phi \quad \phi \leftarrow \phi \vee \phi \bullet
 \end{array}$$

- The first two axioms plus modus ponens are sufficient for proving the deduction theorem. (corresponding to implication introduction).

The only rule is **modus ponens** from ϕ and $\phi \leftarrow \psi$ conclude ψ .

$$((\chi \leftarrow \phi) \leftarrow (\phi \leftarrow \phi)) \leftarrow ((\chi \leftarrow \phi) \leftarrow \phi) \bullet$$

$$(\phi \leftarrow \phi) \leftarrow \phi \bullet$$

Hilbert type system

- To get CPC add $((\phi \rightarrow \psi) \rightarrow \psi) \leftarrow \psi$ (Peirce's Law) or $\top \vdash \psi \leftarrow \psi$.

Classical propositional calculus

- $w \models \phi \leftarrow \psi \iff \forall w' (w R w' \text{ and } w' \models \psi \leftarrow \phi)$,
- $w \models \phi \vee \psi \iff w \models \phi \text{ or } w \models \psi$,
- $w \models \phi \vee \psi \iff w \models \phi \text{ and } w \models \psi$,
- $w R w' \& w \in V(d) \iff w' \in V(d)$.
- For models \mathfrak{M} a persistent valuation V is added. Persistence means:
 - An accessibility relation R , which is a \leq -partial order,
 - A set of worlds W , also nodes, points
 - Frames, (usually \mathcal{G}):

Kripke frames and models

- \Leftrightarrow for finite models $\rightarrow \Box_w (\Diamond w \wedge \Box w \text{ end point} \Leftarrow \Diamond \Box \Diamond)$.
- Note that $w \models \neg \varphi \Leftrightarrow \Box_{w'} (\Diamond w' \wedge \Box w' \wedge \Box \varphi)$
- $\Diamond w \wedge \Box \varphi \Leftarrow \Box \Diamond \varphi$.
- Persistence for formulas follows:
 $w \models \neg \varphi \Leftrightarrow \Box_{w'} (\Diamond w' \wedge \neg \varphi)$ (follows from definition of $\neg \varphi$ as $\varphi \leftarrow \perp$),
 $w \models \neg \varphi \Leftrightarrow \Box_{w'} (\Diamond w' \wedge \neg \varphi) \Leftarrow \neg \varphi \models \perp$,
 $w \not\models \perp$.
- Frames will usually have a **root** w_0 : $w_0 R w$ for all w .


Kripke frames and models, continued

- Persistence transfers to formulas here as well.
- $w \models \forall x \phi(x) \iff$, for each w' , with $w R w'$, and all $d \in D^{w'}$, $w' \models \phi(d)$,
- $w \models \exists x \phi(x) \iff$, for some $d \in D^w$, $w \models \phi(d)$,
- with names for the elements of the domains:
- $w R w' \iff D^w \subseteq D^{w'}$.
- Increasing domains D^w :

Kripke frames and models, predicate logic

- $d \sqcap \neg d \vdash (d \leftarrow d \sqcap) (p)$
- $(c) (d \leftarrow d) \wedge (b \leftarrow d \sqcap) \vdash (b \leftarrow d \sqcap) (r)$
- $(q) (r \leftarrow d) \wedge (b \leftarrow d) \vdash (r \wedge b \leftarrow d)$
- $(a) d \leftarrow d \sqcap \vdash d \sqcap \wedge d$
- These figures give counterexamples to respectively:


Figure 1: Counter-models for the propositional formulas



Counter-models to propositional formulas

- (a) $\Box \Diamond (Ax \vee \Box Ax)$, if domain constant \mathbb{N} (and also against $\Box x \Box Ax \leftarrow \Box \Diamond \Box Ax$),
- (b) $\Diamond \Box (A \vee Bx) \leftarrow A \vee \Diamond x Bx$.
- These figures give counterexamples to:

Figure 2: Counter-models for the predicate formulas



Counter-models to predicate formulas

- φ is valid in a model \mathfrak{M} , $\mathfrak{M} \models \varphi$, if φ is satisfied in all worlds in the model. φ is valid in a frame $\mathbf{mathfrak{F}}$, $\mathbf{mathfrak{F}} \models \varphi$ if φ is valid in all models on the frame.
- Soundness (\Leftarrow) just means checking all axioms in Hilbert type system (plus the fact that modus ponens leaves validity intact).
- Completeness Theorem:
- $\vdash_{IPC} \varphi$ iff φ is valid in all (finite) frames.
- Soundness (\Leftarrow) just means checking all axioms in Hilbert type system (plus the fact that modus ponens leaves validity intact).

Soundness and Completeness

- Before the completeness proof an application of completeness.
- Gilivenko's Theorem, Theorem 5:
- $\vdash \text{CPC } \varphi \text{ iff } \vdash \text{IPC } \varphi$ (CPC is classical propositional calculus).
- \Rightarrow is of course trivial.
- \Leftarrow Exercise.
- e.g. $\vdash \text{IPC} \neg(\varphi \vee \neg\varphi)$.
- Gilivenko's Theorem does not extend to predicate logic, exercise.

Gilivenko's theorem

- Lindenbaum type lemma needed.
or $\psi \in T$.
- A set T of formulas has the disjunction property if $\phi \vee \psi \in T$ implies $\phi \in T$
- A set T of formulas is a theory if T is closed under IPC-induction.
- Theories with the disjunction property,
- Basic entities in Henkin type completeness proof are:

Proof of Completeness

- $\Delta^n \not\models \text{IPC } X, \Delta \not\models \text{IPC } X,$
- Δ is the union of all Δ^n .
- $\Delta^{n+1} = \Delta^n$ otherwise.
- $\Delta^{n+1} = \Delta^n \cup \{\varphi_n\}$ if this does not prove X ,
- $\Delta^0 = T \cup \{\psi\},$
- Enumerate all formulas: $\varphi_0, \varphi_1, \dots$ and define:
 - Proof.
- Lemma 10 If $T \cup \{\psi\} \not\models \text{IPC } X$, then a theory with the disjunction property Δ exists such that $T \subseteq \Delta$, $\psi \in \Delta$ and $X \notin \Delta$.

Lemma

- **Claim:** Δ has the disjunction property:
- Δ is a theory.
- Assume $\phi \vee \psi \in \Delta$, $\phi \notin \Delta$, $\psi \notin \Delta$.
- Let $\phi = \phi_m$ and $\psi = \psi_n$ and w.l.o.g. let $n < m$.
- $\Delta^n \cup \{\phi\} \vdash_{IPC} \chi$ and $\Delta^n \cup \{\psi\} \vdash_{IPC} \chi$, and thus $\Delta^n \cup \{\phi \vee \psi\} \vdash_{IPC} \chi$.
- But $\Delta^n \cup \{\phi \vee \psi\} \not\subseteq \Delta$ and $\Delta \not\vdash_{IPC} \chi$, Contradiction.

Proof lemma, continuation

get the n -canonical model (or n -Henkin model).

- The construction can be restricted to formulas in n variables. We then
 - Evaluation of V_C of canonical model: $\Gamma \models V_C(p) \Leftrightarrow p \in \Gamma$.
 - Frame of canonical model is $\mathcal{G}_C = (W_C, R_C)$.
 - $R_C = \overline{C}$,
 - W_C is the set of all consistent theories with the disjunction property,
 - \mathfrak{M}_C

Canonical model

- Theorem 12. $\Gamma \vdash_{IPC} \varphi$ iff φ is valid in all Kripke models of Γ for IPC.
- For the Completeness side (\Rightarrow) we show: if $\Gamma \not\vdash_{IPC} \varphi$, then $\varphi \notin \Delta$ for some Δ containing Γ in the canonical model.
- First show by induction on φ that $\Theta \models \varphi \Leftrightarrow \varphi \in \Theta$.
- Most cases easy: it is for example necessary to show that $\varphi \wedge X \in \Theta \Leftrightarrow \varphi \in \Theta \wedge X \in \Theta$. This follows immediately from the fact that Θ is a theory (closed under IPC-induction). The corresponding fact for \vee is the disjunction property.

Completeness of IPC

- The hardest is showing that, if $\phi \rightarrow \chi \notin \Theta$, then a theory Δ with the disjunction property such that $\Theta \subseteq \Delta$ exists with $\phi \in \Delta$ and $\chi \notin \Delta$.
- But this is the content of Lemma 10.
- Now assume $\Gamma \not\vdash_{IPC} \phi$. Then $\Gamma \not\vdash_{IPC} \top \rightarrow \phi$. Lemma 10 supplies the required Δ .

Completeness of IPC, continued

- **Theorem** For finite T , $T \vdash_{IPC} \varphi$ iff φ is valid in all finite Kripke models of T for IPC.
- **Proof.** The proof can be done by **filtration**. We will not do that here. Or by reducing the whole discussion to the set of subformulas of $T \cup \{\varphi\}$ (a so-called **adequate** set, both in the definition of the (reduced) canonical model as well as in the proof.)
- Same for a language with only **finite** many propositional variables. (Model will not be finite!)

Finite Model Property

- Let C_0, C_1, C_2, \dots be a sequence of disjoint countably infinite sets of new constants. It suffices to consider theories in the languages \mathcal{L}_n obtained by adding $C_0 \cup C_1 \dots \cup C_n$ to the original language \mathcal{L} . We consider theories containing $\exists x\phi(x) \rightarrow \phi(c_\phi)$ as in the classical Henkin proof. That will immediately guarantee that the theories besides the disjunction property, also have the analogous existence property. The proof then proceeds as in the propositional case. The role of the additional constants becomes clear in the induction step for the universal quantifier.
- If Θ is a theory in \mathcal{L}_n . To show is:
 - $\forall x\phi(x) \in \Theta$ iff, for each d and Θ' , in \mathcal{L}_m ($m \leq n$) with $\Theta \subseteq \Theta'$, $\phi(d) \in \Theta'$.
 - \Leftarrow is of course obvious because Θ' is a theory.

Completeness of Predicate Logic

For \Rightarrow assume that $\forall x \phi(x) \notin \Theta$. Then, for some new constant d in C^{n+1} , $\Theta \not\vdash \phi(d)$. And hence Θ can be extended to a Henkin theory Θ' , with the disjunction property in C^{n+1} that does not prove $\phi(d)$ either.

Completeness of Predicate Logic, continued

- Definition 7. $R(u) = \{u' \in W \mid uRu'\}$,
- The generated subframe \mathfrak{F}^u of \mathfrak{F} is $(R(u), R')$, where R' the restriction of R to $R(u)$.
- The generated submodel \mathcal{A}^u of \mathcal{A} is \mathfrak{F}^u with V restricted to it.
- If $\mathfrak{F}_1 = (W_1, R_1)$ and $\mathfrak{F}_2 = (W_2, R_2)$, then their disjoint union $\mathfrak{F}_1 \sqcup \mathfrak{F}_2$ has as its set of worlds the disjoint union of W_1 and W_2 , and R is $R_1 \cup R_2$. To get the disjoint union of two models the union of the two valuations is added.

Generated subframes and submodels, disjoint unions

- $w \in W, w \in V(d) \text{ iff } f(w) \in V'(d)$.
- p-morphism from \mathcal{A} to \mathcal{A}' , iff f is a p-morphism of the frames and, for all
- If $\mathcal{A} = (W, R, V)$ and $\mathcal{A}' = (W', R', V')$ are models, then $f: W \rightarrow W'$ is a
 - If $f: W \rightarrow W'$ and $R' = \{(w', w) \mid (f(w), f(w')) \in R\}$, then f is a p-morphism (also bounded morphism) from \mathcal{A} to \mathcal{A}' , iff
 - for each $w, w' \in W, w' \in W', wRw', \text{ if } f(w)R'f(w'), \text{ then } f(w)Rf(w')$,
 - for each $w, w' \in W, wRw', \text{ if } f(w)R'f(w'), \text{ then } f(w)Rf(w')$,
 - for each $w, w' \in W, wRw', \text{ if } f(w)R'f(w'), \text{ then } f(w)Rf(w')$.

p-morphisms

- If $\mathfrak{X} \models \phi$, then $\mathfrak{X}_w \models \phi$.
- This implies that if ϕ is falsified in a model, we may w.l.o.g. assume that it is falsified in the root.
- If w' in the generated submodel \mathfrak{M}_w , then, $w' \models \phi$ in \mathfrak{M} iff $w' \models \phi$ in \mathfrak{M}_w .
- Lemma

Properties of Generated Subframes


- If $w \in W^1$, then $w \models \phi$ in $\mathcal{M}^1 \oplus \mathcal{M}^2$ iff $w \models \phi$ in \mathcal{M}^1 , etc.
- If f is a p-morphism from \mathfrak{X} onto \mathfrak{Y} , then $\mathfrak{X} \models \phi$ implies $\mathfrak{Y} \models \phi$.
- If $\mathfrak{X} \models \phi$, then $\mathfrak{X}^w \models \phi$.
- If f is a p-morphism from \mathcal{M} to \mathcal{M}' and $w \in W$, then $w \models \phi$ iff $f(w) \models \phi$.

Properties of p-morphic images, disjoint unions

- Theorem 16. $\vdash \text{IPC} \psi \vee \phi \text{ iff } \vdash \text{IPC} \psi \text{ or } \vdash \text{IPC} \phi$.
- This extends to the predicate calculus and arithmetic.
- Proof. \Rightarrow : Trivial
- \Leftarrow : Assume $\nvdash \text{IPC} \psi$ and $\nvdash \text{IPC} \phi$.
- Let $\mathcal{A} \not\models \psi$ and $\mathcal{C} \not\models \phi$.
- Add a new root w_0 below both \mathcal{A} and \mathcal{C} . In w_0 , $\psi \vee \phi$ is falsified (because of persistence!).

Disjunction property

Figure 3: Providing the disjunction property



- An often used theorem is $\Box\phi \vee \Box\neg\phi \leftrightarrow \Box(\phi \vee \neg\phi)$.

and the rule of **necessitation** $\phi / \Box\phi$

$$\Box(\phi \rightarrow \psi) \leftarrow (\Box\phi \rightarrow \Box\psi)$$

classical propositional calculus **CPC** the axiom scheme

- The basic modal logic **K** has as in addition to the axiom schemes of the
- with an additional 1-place operator \Box (pronounced: **necessary**),
- The language of modal logic is the language of the propositional calculus

Modal Logic

- and $\Box(\Box\phi \leftarrow \phi) \leftarrow \Box\phi$ for GL.
- In addition to this Grzegorczyk's axiom $\Box(\Box\phi \leftarrow \Box\phi) \leftarrow \phi$ for Grz,
- The axioms $\Box\phi \leftarrow \phi$, $\Box\phi \leftarrow \Box\Box\phi$ for S4
- The axiom $\Box(\phi \leftarrow \phi) \leftarrow (\Box\phi \leftarrow \Box\phi)$ of K,
- The modal-logical systems S4, Grz and GL are obtained by adding to

S4, Grz and GL

- $w \models \phi \iff \forall u (wRu \iff u \models \phi)$,
- $w \models \phi \vee \psi \iff w \models \phi \text{ and } w \models \psi$, etc.
- $w \models \forall u \in V(d) \iff w \in V(d)$.
- For models a valuation V is added.
- An accessibility relation R ,
- A set of worlds V , also nodes, points
- Frames:

Kripke frames and models for K

Proof If $T \vdash \phi$, then $\{\Box\phi \mid \phi \in T\} \vdash K\Box\phi$

- **Lemma** If $\{\Box\phi \mid \phi \in T\} \nvdash K\Box\phi$, then $T \nvdash \phi$,
- **Lemma** needed.

- **Maximal consistent sets** (these are of course also theories with the disjunction property),
- Basic entities in Henkin type completeness proof for K are:

Completeness of K

- Value of V_K of canonical model: $\Gamma \in V_K(d) \Leftrightarrow d \models \Gamma$.
- Frame of canonical model is $\mathfrak{F}_K = (W_K, R_K)$.
- $TR_K \Delta \leftrightarrow (\Delta \Box \phi \in \Gamma \Leftrightarrow \phi \in \Delta)$,
- W_K is the set of all maximal consistent sets,
- $\mathcal{M}_K = (W_K, W_K, V_K) = (\mathfrak{F}_K, V_K)$
- The canonical model \mathcal{M}_K is defined as follows:

Canonical model of K

- A modal logic L is said to define or characterize the class of frames \mathfrak{F} such that $\mathfrak{F} \models L$.

$$\mathfrak{F} \models \phi \Leftrightarrow \forall \mathfrak{m} \in \mathfrak{F} (\mathfrak{m} \models \phi)$$

$$\mathfrak{M} \models \phi \Leftrightarrow \forall w \in W (w \models \phi)$$

- Definition

Validity on models, frames, characterization

- GL is complete w.r.t. the finite \leq -partial orders.
- GL characterizes the transitive, conversely well-founded (i.e. irreflexive, asymmetric) frames.
- Grz is complete w.r.t. the finite \leq -partial orders,
- Grz characterizes the reflexive, transitive, conversely well-founded frames,
- S4 is complete w.r.t. \leq -partial orders (reflexive, transitive, anti-symmetric)
- S4 is complete w.r.t. the (finite) reflexive, transitive frames,
- S4 characterizes the reflexive transitive frames,
- GL is complete w.r.t. the finite \leq -partial orders <-partial orders.

Kripke frames, models for S4, Grz and GL

- $\top_u = \top$
- $\phi_u \leftarrow \phi =_u (\phi \leftarrow \phi)$
- $(\phi_u \wedge \psi_u) \leftarrow =_u (\phi \wedge \psi)$
- $\phi_u \vee \psi_u =_u (\phi \vee \psi)$
- $d_u = \perp \perp d$
- Definition 28
- extends to the predicate calculus and arithmetic, has many variations,
- Gödel's negative translation

Translations

- **Theorem 29.** $\vdash_{\text{CPC}} \varphi \text{ iff } \vdash_{\text{IPC}} \varphi_n$.
- **Proof.**
- $\Rightarrow : \vdash_{\text{IPC}} \varphi_n \Leftarrow \vdash_{\text{CPC}} \varphi_n \Leftarrow \vdash_{\text{CPC}} \varphi$.
- $\Leftarrow :$ First prove $\vdash_{\text{IPC}} \varphi_n \leftrightarrow \neg \neg \varphi_n$ (φ_n is negative) (using $\neg \neg$ -introduction).
- Then simply follow the proof of φ in CPC to mimic it with a proof of φ_n .
- in IPC. Exercise.

Properties of Gödel's negative translation

- Gödel noticed the closeness of S4 and IPC when one interprets \Box as *intuitive provability*.
- Definition 32.
- $d_\Box = \Box d$,
- $\phi \vee \Box \psi = \Box (\phi \vee \psi)$,
- $\phi \wedge \Box \psi = \Box (\phi \wedge \psi)$,
- $(\phi \leftarrow \psi) \Box = \Box (\psi \leftarrow \phi)$,
- Theorem 33 $\vdash_{IPC} \phi \text{ iff } \vdash_{S4} \phi \Box \text{ iff } \vdash_{Gz} \phi \Box$.

Gödel's translation of IPC into S4

- **Proof \Leftarrow** : Trivial from S4 to Grz. From IPC to S4 it is simply a matter of using one of the proof systems of IPC and to find the needed proofs in S4, or showing their validity in the S4-frames and using completeness.
- \Rightarrow : It is sufficient to note that it is easily provable by induction on the length of the formula φ that for any world w in a Kripke model with a persistent valuation $w \models \varphi$ iff $w \models \varphi$. This means that if $\vdash_{IPC} \varphi$ one can interpret the finite IPC-countermodel to φ provided by the completeness theorem immediately as a finite Grz-countermodel to φ .



Proof for Gödel's translation of IPC into S4

- e.g. the **Kreisel-Putnam logic** ($\neg\phi \leftarrow \phi \vee (\phi \leftarrow \chi) \vdash (\chi \leftarrow \phi)$),
most do not have disjunction property, some do:
- **Dummett's logic**: $(\phi \leftarrow \phi) \vee (\phi \leftarrow \psi)$,
- e.g. **Weak excluded middle**: $\neg\phi \vee \neg\neg\phi$,
classical logic),
- Logics extending intuitionistic logic by axiom schemes (and sublogics of
intuitionistic logic) (Superintuitionistic logics),

Intermediate Logics

- $f^{n+2}(\phi) = def g^n(\phi \wedge g^{n+1}(\phi))$
- $g^{n+4}(\phi) = def g^{n+3}(\phi \wedge g^n(\phi \leftarrow (\phi) \wedge f^{n+2}(\phi)))$
- $g^3(\phi) = def \phi \leftarrow \phi \leftarrow f^{n+2}(\phi)$
- $g^2(\phi) = def \perp \sqsubset \phi$
- $g_1(\phi) = f_1(\phi) = def \perp \sqsubset \phi$
- $g_0(\phi) = f_0(\phi) = def \phi$
- Definition 36. Rieger-Nishimura Lattice.

The Rieger-Nishimura Lattice and Ladder



subframes generated by w_k will be called RN_k .

- The frame of the Rieger-Nishimura ladder will be called RN . Its $\psi(p)$ can be reached from $\varphi(p)$ by a downward going line.
- In the Rieger-Nishimura lattice a formula $\varphi(p)$ implies $\psi(p)$ in IPC iff fact, in the Rieger-Nishimura Ladder w_i validates $g_n(p)$ for $i \leq n$ only.
- All formulas $f_n(p)$ ($n \leq 2$) and $g_n(p)$ ($n \leq 0$) are non-equivalent in IPC. In to a formula $f_n(p)$ ($n \leq 2$) or $g_n(p)$ ($n \leq 0$), or to \top or \perp .
- Each formula $\varphi(p)$ with only the propositional variable p is IPC-equivalent to Theorem 37.

The Rieger-Nishimura Lattice and Ladder II

- Algebraic completeness of IPC
- Heyting algebras and Kripke frames
- Lattices, distributive lattices and Heyting algebras

Overview

Heyting algebras


Lattices

A partially ordered set (A, \leq) is called a **lattice** if every two element subset of A has a least upper and greatest lower bound.

Let (A, \leq) be a lattice. For $a, b \in A$ let

$$a \wedge b := \inf\{a, b\} \text{ and } a \vee b = \sup\{a, b\}.$$





We assume that every lattice is bounded, i.e., it has a least and a greatest element denoted by \bot and \top respectively.

Lattices, top and bottom

1. $a \vee a = a$, $a \vee a = a$; (*idempotency laws*)
2. $a \vee b = b \vee a$, $a \vee b = b \vee a$; (*commutative laws*)
3. $a \vee (b \vee c) = (a \vee b) \vee c$, $a \vee (b \vee c) = (a \vee b) \vee c$; (*associative laws*)
4. $a \vee \top = a$, $a \vee \top = a$; (*existence of \top and \top*)
5. $a \vee (q \vee a) = a$, $a \vee (q \vee a) = a$. (*absorption laws*)

Proposition 40. A structure $(A, \vee, \wedge, \perp, \top)$ is a lattice iff for every $a, b, c \in A$ the following holds:

Lattices, axioms

Lattices, axioms, continued

Proof.(Sketch)


\Rightarrow Suppose $(A, \vee, \wedge, \perp, \top)$ satisfies the axioms 1–5.

\Leftarrow Check that every lattice satisfies the axioms 1–5.

Define $a \leq b$ by putting $a \vee b = b$ or equivalently by putting $a \wedge b = a$.

Check that (A, \leq) is a lattice. \square

We denote lattices by $(A, \vee, \wedge, \perp, \top)$.



The lattices M_5 and N_5 are not distributive.

- $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$
- $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$

the **distributive laws**:

Definition 41. A lattice $(A, \vee, \wedge, \perp, \top)$ is called **distributive** if it satisfies

Distributive lattices

Theorem 43. A lattice L is distributive iff M_5 and N_5 are not sublattices of L .

Distributive lattices, characterization

$$a \leftarrow b = \bigwedge \{c \in A : a \vee c \leq b\}.$$

In every Heyting algebra \mathcal{A} we have that

$$c \leq a \leftarrow b \text{ iff } a \vee c \leq b.$$

Definition 44. A distributive lattice $(A, \wedge, \vee, \perp, \top)$ is said to be a **Heyting algebra** if for every $a, b \in A$ there exists an element $a \leftarrow b$ such that for every $c \in A$ we have:

Heyting algebras

$$1. \quad a \leftarrow a = \top$$

$$2. \quad a \vee (a \leftarrow q) = a \vee q$$

$$3. \quad q = (q \leftarrow a) \vee a$$

$$4. \quad a \leftarrow (a \vee c) = (a \leftarrow a) \vee (a \leftarrow c)$$

Theorem 47. A (distributive) lattice $\mathfrak{A} = (A, \wedge, \vee, \perp, \top)$ is a Heyting algebra iff there is a binary operation \rightarrow on A such that for every $a, b, c \in A$:

Heyting algebras, axioms

$$a \vee \bigwedge_{i \in I} b^i = \bigwedge_{i \in I} a \vee b^i.$$

Proposition 45. A complete distributive lattice $(A, \wedge, \vee, \perp, \top)$ is a Heyting algebra iff it satisfies the infinite distributivity law

We say that a lattice (A, \wedge, \vee) is **complete** if for every subset $X \subseteq A$ there exist $\inf(X) =: \bigwedge X$ and $\sup(X) =: \bigvee X$.

Complete distributive lattices

- Every Boolean algebra is a Heyting algebra.

$$a \leftarrow b = \begin{cases} q & \text{if } a < b, \\ \perp & \text{if } a \leq b, \end{cases}$$

For every $a, b \in C$ we have

- Every chain C with a least and greatest element is a Heyting algebra.
- Every finite distributive lattice is a Heyting algebra.

More examples

- Proposition 49. Let $\mathcal{A} = (A, \wedge, \vee, \neg, \top)$ be a Heyting algebra. Then the following three conditions are equivalent:
1. \mathcal{A} is a Boolean algebra;
 2. $a \vee \neg a = \top$ for every $a \in A$;
 3. $\neg \neg a = a$, for every $a \in A$.
- For every element a of a Heyting algebra let $\neg a := a \rightarrow \perp$.

Boolean algebras

- $R_1(U) = \bigcup_{u \in U} R_1(u)$
 - $R(U) = \bigcup_{u \in U} R(u)$
 - $R_1(u) = \{v \in W : uRv\}$
 - $R(u) = \{v \in W : uRv\}$
- For every $u \in W$ and $U \subseteq W$ let
- Let $\mathfrak{F} = (W, R)$ be an intuitionistic Kripke frame.

frames

The connection between Heyting algebras and Kripke

Proposition. $(Up(\mathcal{X}), \cup, \cap, \leftarrow, \emptyset)$ is a Heyting algebra.

$$\cdot(A \setminus \cap) = R^{-1}(A \setminus M)$$

$$U \leftarrow A = \{u \in M : \text{for every } v \in M \text{ with } uRv \text{ if } v \in U \text{ then } u \in V\}$$

For $U, V \in Up(\mathcal{X})$, let

Let $Up(\mathcal{X})$ be the set of all upsets of \mathcal{X} .

A subset $U \subseteq W$ is called an **upset** if $w \in U$ and wRv implies $v \in U$.

Heyting algebras and Kripke frames, continued

A triple $\mathfrak{F} = (W, R, A)$ is called a **general frame**.
 A is a Heyting algebra.
Let A be a set of upsets of \mathfrak{F} closed under \cup , \cap , \neg and containing \emptyset .

General Frames

- The duality does not generalize easily to general frames in general. We use the **descriptive frames**. They are general frames with two additional properties:
 - **finite intersection property**:
 \mathcal{F} is **compact** if $\forall A \subseteq \mathcal{A}, \exists U \subseteq \{W \mid U \in A\} (A \cap U \neq \emptyset)$,
 - **refined** if $\forall w, u \in W, \neg(wRu) \iff \exists U \in A (w \in U \wedge u \notin U)$,
 - \mathcal{F} is **isomorphic** to $(A, \sqcup, \sqcap, \rightarrow, \emptyset)$.

Descriptive Frames

For $Y \subseteq X$, the **interior** of Y is the set $\mathbf{I}(Y) = \bigcup\{U \in \mathcal{O} : U \subseteq Y\}$.

- If $U_i \in \mathcal{O}$, for every $i \in I$, then $\bigcup_{i \in I} U_i \in \mathcal{O}$
- If $U, V \in \mathcal{O}$, then $U \cup V \in \mathcal{O}$
- $X, \emptyset \in \mathcal{O}$

Definition 51. A pair $\mathcal{X} = (X, \mathcal{O})$ is called a **topological space** if $X \neq \emptyset$ and \mathcal{O} is a set of subsets of X such that

The connection of Heyting algebras and topology

For every $U, V \in O$ let

$$(A \cap (U \setminus X))I = A \leftarrow U$$

Proposition. $(O, \cup, U, \leftarrow, \emptyset)$ is a Heyting algebra.

Heyting algebras and topology, continued

- $a \vee b \in F$ implies $a \in F$ or $b \in F$
 - A filter F is called prime if
 - $a \in F$ and $a \leq b$ imply $b \in F$
 - $a, b \in F$ implies $a \vee b \in F$
- $F \subseteq A$ is called a filter if
- Let $\mathfrak{A} = (A, \wedge, \vee, \rightarrow, \perp)$ be a Heyting algebra.
- How to obtain a Kripke frame from a Heyting algebra?

Kripke frames from Heyting algebras

Kripke frames from Heyting algebras, continued

If \mathcal{A} is a Boolean algebra, then every prime filter of \mathcal{A} is maximal.

This is not the case for Heyting algebras.

Let $W := \{F : F \text{ is a prime filter of } \mathcal{A}\}.$

For $F, F' \in W$ we say that $F \leq F'$ if $F \subseteq F'$.

(W, R) is an intuitionistic Kripke frame.

An algebra \mathcal{A}' is called a **homomorphic image** of \mathcal{A} if there exists a homomorphism from \mathcal{A} onto \mathcal{A}' .

$$h(\top) = \top \bullet$$

$$h(q \leftarrow a) = (q \leftarrow a) \bullet$$

$$h(a \wedge q) = (a \wedge q) \bullet$$

$$h(a \vee q) = (a \vee q) \bullet$$

A map $h : A \rightarrow A'$ is called a **Heyting homomorphism** if

algebras.

Let $\mathcal{A} = (A, \wedge, \vee, \leftarrow, \top)$ and $\mathcal{A}' = (A', \wedge', \vee', \leftarrow', \top')$ be Heyting

Basic algebraic operations, homomorphisms

$$(, \top, \top) =: \top \bullet$$

$$(, q, \leftarrow, a, q \leftarrow a) =: (, q, q) \leftarrow (, a, a) \bullet$$

$$(, a, \wedge, b, a, \wedge, b) =: (, q, q) \wedge (, a, a) \bullet$$

$$(, a, \vee, b, a, \vee, b) =: (, q, q) \vee (, a, a) \bullet$$

A **product** $\mathcal{A} \times \mathcal{A}'$ of \mathcal{A} and \mathcal{A}' is the algebra $(A \times A', \wedge, \vee, \leftarrow, \top)$, where

\mathcal{A}' is a **subalgebra** of \mathcal{A} if $A' \subseteq A$ and for every $a, b \in A'$, $a \wedge b, a \vee b, a \leftarrow b, \top \in A'$.

Basic algebraic operations, subalgebras

For a homomorphism $h : \mathcal{A} \rightarrow \mathcal{A}'$, let $\phi(h) : \phi(\mathcal{A}') \rightarrow \phi(\mathcal{A})$ be such that for every element $F \in \phi(\mathcal{A}')$ we have $\phi(h)(F) = h^{-1}(F)$.

$$\mathcal{A} \hookrightarrow \phi(\mathcal{A}) = (W, R).$$

We define $\phi : \text{Heyt} \rightarrow \text{Kripke}$ and $\Psi : \text{Kripke} \rightarrow \text{Heyt}$.

Let **Kripke** denote the category of intuitionistic Kripke frames and d -morphisms.

Let **Heyt** be a category whose objects are Heyting algebras and whose morphisms are Heyting homomorphisms.

Categories

Define a functor $\Psi : \text{Kripke} \rightarrow \text{Heyt}$.

For every Kripke frame \mathfrak{X} let $\Psi(\mathfrak{X}) = (\mathcal{U}^p(\mathfrak{X}), \cup, \cap, \leftarrow, \emptyset)$.

If $f : \mathfrak{X} \rightarrow \mathfrak{Y}$ is a p -morphism, then $\Phi(f) : \Phi(\mathfrak{X}) \rightarrow \Phi(\mathfrak{Y})$ is such that
for every element of $U \in \Phi(\mathfrak{Y})$ we have $\Phi(f)(U) = (f^{-1}(U))_f$.

Categories, continued

- If $\mathcal{A} \times \mathcal{B}$ is a product of \mathcal{A} and \mathcal{B} , then $\phi(\mathcal{A} \times \mathcal{B})$ is isomorphic to the disjoint union $\phi(\mathcal{A}) \sqcup \phi(\mathcal{B})$.
- If $\mathcal{A} \times \mathcal{B}$ is a subalgebra of \mathcal{B} , then $\phi(\mathcal{A})$ is a p -morphic image of $\phi(\mathcal{B})$.
- If \mathcal{A} is a subalgebra of \mathcal{B} , then $\phi(\mathcal{A})$ is isomorphic to a generated subframe of $\phi(\mathcal{B})$.
- If \mathcal{A} is a homomorphic image of \mathcal{B} , then $\phi(\mathcal{A})$ is isomorphic to a frame.

Theorem 57. Let \mathcal{A} and \mathcal{B} be Heyting algebras and \mathcal{G} and \mathcal{S} Kripke frames.

Duality

- the product $\Psi(\mathfrak{X}) \times \Psi(\mathfrak{G})$.
- If $\mathfrak{X} \oplus \mathfrak{G}$ is a disjoint union of \mathfrak{X} and \mathfrak{G} , then $\Psi(\mathfrak{X} \oplus \mathfrak{G})$ is isomorphic to
 - If \mathfrak{X} is a d -morphic image of \mathfrak{G} , then $\Psi(\mathfrak{X})$ is a subalgebra of $\Psi(\mathfrak{G})$.
2. • If \mathfrak{X} is a generated subframe of \mathfrak{G} , then $\Psi(\mathfrak{X})$ is isomorphic to a homomorphic image of $\Psi(\mathfrak{G})$.

Duality, continued

NO!

Is $\Psi(\text{Kripke})$ isomorphic to Heyt ?

Is $\phi(\text{Heyt})$ isomorphic to Kripke ?

Duality, continued 2

\mathfrak{A} such that \mathfrak{A} is isomorphic to $Up(\mathfrak{X})$.
Theorem 63. For every finite Heyting algebra \mathfrak{A} there exists a Kripke frame

finite Kripke frames respectively, are dually equivalent.
 Restrictions of φ and Ψ to the categories of finite Heyting algebras and

Open question 62. Characterization of Kripke frames in $\Psi(\text{Heyt})$.

Not every Heyting algebra is complete.

$\Psi(\mathfrak{X}) = (Up(\mathfrak{X}), \cup, \cap, \neg, \emptyset)$ is a complete lattice.

Duality, continued 3

- \mathfrak{L} is not isomorphic to a generated subframe of $\Psi(\mathfrak{L})$.
- \mathfrak{A} is not isomorphic to a homomorphic image of $\Psi(\mathfrak{A})$.
- \mathfrak{L} is a p -morphic image of $\Psi(\mathfrak{L})$.
- \mathfrak{A} is a subalgebra of $\Psi(\mathfrak{A})$.

Proposition.

For every Heyting algebra \mathfrak{A} the algebra $\Psi(\mathfrak{A})$ is called a canonical filter extension of \mathfrak{A} . For every Kripke frame \mathfrak{L} the frame $\Psi(\mathfrak{L})$ is called a prime extension of \mathfrak{L} . For every Kripke frame \mathfrak{L} the frame $\Psi(\mathfrak{L})$ is called a descriptive frames.

Duality, continued, 4

Let K be a class of algebras of the same signature. We say that K is a **Variety** if K is closed under homomorphic images, subalgebras and products. We say that K is a **Variety** if K is closed under homomorphic images, subalgebras and products. **Theorem.** (Tarski) K is a variety iff $K = \text{HSP}(K)$, where **H**, **S** and **P** are respectively the operations of taking homomorphic images, subalgebras and products. **Theorem 64.** (Birkhoff) A class of algebras forms a variety iff it is equationally defined.

Heyt is a variety.

Algebraic completeness

$$\begin{aligned}
 \top &= (\top)u & \bullet \\
 (\phi)u &\leftarrow (\phi)u = (\phi \leftarrow \phi)u & \bullet \\
 (\phi)u \wedge (\phi)u &= (\phi \wedge \phi)u & \bullet \\
 (\phi)u \vee (\phi)u &= (\phi \vee \phi)u & \bullet
 \end{aligned}$$

We extend the valuation from P to the whole of $Form$ by putting:

Let $\mathfrak{A} = (A, \wedge, \vee, \leftarrow, \perp)$ be a Heyting algebra. A function $u : P \rightarrow A$ is called a **valuation** into the Heyting algebra \mathfrak{A} .

Let $Form$ be the set of all formulas in this language.

Let P be the (finite or infinite) set of propositional variables.

Valuations on Heyting algebras

algebra.

Proposition 66. (Soundness) $\text{IPC} \vdash \phi$ implies that ϕ is valid in every Heyting

ϕ is valid in \mathcal{A} if ϕ is true for every valuation in \mathcal{A} .

A formula ϕ is true in \mathcal{A} under v if $v(\phi) = \top$.

Soundness

$$[\phi \leftarrow \sigma] = [\phi] \leftarrow [\sigma] \bullet$$

$$[\phi \wedge \sigma] = [\phi] \wedge [\sigma] \bullet$$

$$[\phi \vee \sigma] = [\phi] \vee [\sigma] \bullet$$

Define the operations on $Form/\equiv$ by letting:

$$Form/\equiv := \{[\phi] : \phi \in Form\}.$$

Let $[\phi]$ denote the \equiv -equivalence class containing ϕ .

$$\phi \equiv \psi \text{ iff } \vdash_{IPC} \phi \leftrightarrow \psi.$$

Define an equivalence relation \equiv on $Form$ by putting

Completeness

The operations on Form/\equiv are well-defined.
That is, if $\phi' \equiv \phi''$ and $\psi' \equiv \psi''$, then $\phi' \circ \psi' \equiv \phi'' \circ \psi''$, for $\circ \in \{\wedge, \vee, \rightarrow\}$.
Denote by $F(w)$ the algebra $(\text{Form}/\equiv, \wedge, \vee, \rightarrow, \top)$.
We call $F(w)$ the Lindenbaum-Tarski algebra of IPC or the w -generated free Heyting algebra.

Completeness 2

semantics.

Corollary 69. IPC is sound and complete with respect to algebraic

3. $\text{IPC} \vdash \varphi$ iff φ is valid in $F(n)$, for any formula φ in n variables.

2. $\text{IPC} \vdash \varphi$ iff φ is valid in $F(\omega)$.

1. $F(a)$, for $a \leq \omega$ is a Heyting algebra.

Theorem 68.

Completeness 3

Jankov formulas and intermediate logics

Fix a propositional language \mathcal{L}^n consisting of finitely many propositional

letters p_1, \dots, p_n for $n \in \omega$.

Let $\mathfrak{M} = (W, R, V)$ be an intuitionistic Kripke model.

With every point w of \mathfrak{M} , we associate a sequence $i_1 \dots i_n$ such that for $k = 1, \dots, n$:

$$i_k = \begin{cases} 1 & \text{if } w \models p_k, \\ 0 & \text{if } w \not\models p_k, \end{cases}$$

We call the sequence $i_1 \dots i_n$ associated with w the **color** of w and denote it by $\text{col}(w)$.

Colors

Colors are ordered according to the relation \leq such that $i_1 \dots i_n \leq i'_1 \dots i'_n$ if for every $k = 1, \dots, n$ we have that $i_k \leq i'_k$.

The set of colors of length n ordered by \leq forms an n -element Boolean

algebra.

We write $i_1 \dots i_n < i'_1 \dots i'_n$ if $i_1 \dots i_n \leq i'_1 \dots i'_n$ and $i_1 \dots i_n \neq i'_1 \dots i'_n$.

Covers, anti-chains

For a Kripke frame $\mathfrak{F} = (W, R)$ and $w, u \in W$, we say that a point w is an **immediate successor** of a point u if $w \neq u$, wRu , and there is no $v \in W$ such that $w \neq v$, $v \neq u$, vRu and vRw .

We say that a set A **totally covers** a point u and write $u \subset A$ if A is the set of all immediate successors of u .

$A \subseteq W$ is an **anti-chain** if $|A| < 1$ and for every $w, u \in A$, if $w \neq u$ then $\neg(wRu)$ and $\neg(uRw)$

The 2-universal model $U(2) = (U(2), R, V)$ of **IPC** is the smallest Kripke model satisfying the following three conditions:


1. $\max(U(2))$ consists of 2^2 points of distinct colors.

2. If $u \in U(2)$, then for every color $i_1 i_2 < \text{col}(u)$, there exists $v \in U(2)$ such that $v \prec u$ and $\text{col}(v) = i_1 i_2$.


3. For every finite anti-chain $A \subset U(2)$ and every color $i_1 i_2$, such that $i_1 i_2 \leq \text{col}(u)$ for all $u \in A$, there exists $v \in U(2)$ such that $v \prec A$ and $\text{col}(v) = i_1 i_2$.




The construction of the n -universal model



The construction of the n -universal model





The construction of the n -universal model




The construction of the n -universal model

The construction of the n -universal model







The construction of the n -universal model



The construction of the n -universal model

The construction of the n -universal model







The construction of the n -universal model

1 • 0


J-universal model




L-universal model



J-universal model




J-universal model



J-universal model

J-universal model is called the **Rieger-Nishimura ladder**.



J-universal model

Theorem. For every formula φ in the language \mathcal{L}_n , we have that

$$\text{IPC } \varphi \quad \text{iff} \quad U(n) \models \varphi.$$

Call a set $V \subseteq U(n)$ **definable** if there is a formula φ such that

$$V = \{w \in U(n) : w \models \varphi\}.$$

Every upset of the Rieger-Nishimura ladder is definable.

Not every upset of the n -universal model (for $n > 1$) is definable.

Theorem 42. The Heyting algebra of all definable subsets of the n -universal model is isomorphic to the free n -generated Heyting algebra.

$$\varphi_u = \Box\varphi_u.$$

and

$$\varphi_u := \bigvee \{ d_k : u \models d_k \} \vee \{ d_k = u : u \models d_k \}$$

Let u be a maximal point of $U(n)$. Then

φ_u defines $R(u)$ and ψ_u defines $U(n) \setminus R_{-1}(u)$.

For every formula $u \in U(n)$ we construct formulas φ_u and ψ_u such that

All the point generated upsets of $U(n)$ are definable.

Which upsets of $U(n)$ are definable?

Point generated upsets of $U(n)$

φ_w and ψ_w are called de Jongh formulas.

where w_1, \dots, w_n are all the immediate successors of w .

$$\bigwedge_{u_i}^n \varphi_{w_i} \leftarrow \varphi_w =: \psi_w$$

If φ_w is defined, then

Point generated upsets of $U(n)$, 2

$$(\phi_m \wedge \bigwedge_u \psi^u \leftarrow \bigwedge_u \psi^u) \wedge (n \text{drop}(n) \vee (n \text{drop} \vee \bigvee =: (n) \phi)$$

•

- $n \text{drop}(n) =: \{d^k \mid n \not\models d^k \wedge \forall i \leq k \exists u_i \models d^k\}$, the set of atoms which might have been true in u but aren't.
- Let $\text{prop}(u) =: \{d^k \mid u \models d^k\}$, the atoms true in u .

Point generated upsets of $U(n)$, 3

- For every finite Kripke frame \mathfrak{J} , there exists a valuation V , and $n \leq |\mathfrak{J}|$ such that $\mathfrak{M} = (\mathfrak{J}, V)$ is a generated submodel of $\mathcal{U}(n)$.
 - For every Kripke model $\mathfrak{M} = (\mathfrak{J}, V)$, there exists a Kripke model $\mathfrak{M}' = (\mathfrak{J}', V')$ such that \mathfrak{M}' is a generated submodel of $\mathcal{U}(n)$ and \mathfrak{M}' is a p -morphic image of \mathfrak{M} .
- Theorem 82.

Structure of n -universal model

Theorem The n -universal model is isomorphic to the nodes of finite depth in the n -canonical model.

n -Henkin model and n -universal model

- **Proof** By induction on the depth of the nodes it is shown that the submodel generated by a node in the n -Henkin model is isomorphic to the submodel generated by some node in the n -Henkin model and vice versa.
- \Leftarrow : The p -morphism guaranteed by Theorem 82 from the finite Henkin model into the universal model has to be an isomorphism since all nodes of a Henkin model have distinct theories.
- \Rightarrow : For nodes of depth 1 this is trivial. Consider a node of depth $n + 1$. From the induction hypothesis one sees that there is a p -morphism from the nodes of depth $\leq n$. For nodes of depth $n + 1$ one has to use the de Jongh formula to see that in the n -Henkin model can be only one node above the image of depth $n + 1$ which satisfies the formula.

n -Henkin model and n -universal model

Lemma 88. A frame \mathfrak{F} is a p -morphic image of a generated subframe of a frame \mathfrak{G} iff \mathfrak{F} is a generated subframe of a p -morphic image of \mathfrak{G} .

$\mathfrak{G} \not\models X(\mathfrak{F})$ iff \mathfrak{F} is a p -morphic image of a generated subframe of \mathfrak{G} .

Theorem 87. For every finite rooted frame \mathfrak{F} there exists a formula $X(\mathfrak{F})$ such that for every frame \mathfrak{G}

The Jančkova theorem

Theorem 87(Reformulated). For every finite rooted frame \mathfrak{F} there exists a formula $X(\mathfrak{F})$ such that for every frame \mathfrak{G} $\mathfrak{G} \not\models X(\mathfrak{F})$ iff \mathfrak{F} is a generated subframe of a d -morphic image of \mathfrak{G} .

The result follows from the duality of Heyting algebras and Kripke frames.

Heyt has the congruence extension property.

Proof. It is a universal algebraic result that if a variety V has the congruence extension property, then for every algebra $\mathfrak{A} \in V$ we have that $HS(\mathfrak{A}) = SH(\mathfrak{A})$.

Congruence extension property

Proof of the Jankov theorem

Let \mathfrak{F} be a finite rooted frame.
Then exists $n \in \omega$ such that \mathfrak{F} is (isomorphic to) a generated subframe of $\mathcal{U}(n)$.
Let $w \in U(n)$ be the root of \mathfrak{F} . Then \mathfrak{F} is isomorphic to \mathfrak{F}^w .
Let $\chi(\mathfrak{F}) = \phi_w$.

Proof of the Jančák theorem, 2

$\mathfrak{L} \neq \phi^u$ hence if \mathfrak{X} is a generated subframe of a p -morphic image of \mathfrak{Q} then $\mathfrak{Q} \neq \phi^u$.

If $\mathfrak{Q} \neq \phi^u$, then there is a valuation V such that model $\mathfrak{M} \neq \phi^u$, where $\mathfrak{M} = (\mathfrak{Q}, V)$.

We can assume that there is a p -morphic image \mathfrak{M}' of \mathfrak{M} such that \mathfrak{M}' is a generated submodel of $U(n)$.

Then $\mathfrak{M}' \neq \phi^u$. Which implies that \mathfrak{X}^u is a generated subframe of \mathfrak{M}' .


- In the infinite case \leq , in general, is not antisymmetric.
- If we restrict ourselves to only finite Kripke frames, then \leq is a partial order.
- \leq is reflexive and transitive.

$\mathfrak{F} \leq \mathfrak{G}$ if \mathfrak{F} is a p -morphic image of a generated subframe of \mathfrak{G} .

We say that

Let \mathfrak{F} and \mathfrak{G} be Kripke frames.

Applications of Jančov formulas



Consider the sequence Δ of finite Kripke frames shown below

If $\mathfrak{F} \leq \mathfrak{F}'$, then for every frame \mathfrak{G} we have that $\mathfrak{G} \models X(\mathfrak{F})$ implies $\mathfrak{G} \models X(\mathfrak{F}')$.

Let \mathfrak{F} and \mathfrak{F}' be two finite rooted frames.

The Jankovic chain

Properties of the Jankov chain

Lemma 91. Δ forms a \leq -antichain.

For every set Γ of Kripke frames. Let $\text{Log}(\Gamma)$ be the logic of Γ , that is,

$$\text{Log}(\Gamma) = \{\phi : \mathfrak{F} \models \phi \text{ for every } \mathfrak{F} \in \Gamma\}.$$

Theorem 92. For every $\Gamma_1, \Gamma_2 \subseteq \Delta$, if $\Gamma_1 \neq \Gamma_2$, then $\text{Log}(\Gamma_1) \neq \text{Log}(\Gamma_2)$.

Proof. Without loss of generality assume that $\Gamma_1 \not\subseteq \Gamma_2$.

This means that there is $\mathfrak{F} \in \Gamma_1$ such that $\mathfrak{F} \notin \Gamma_2$.

Consider the Jankov formula $\chi(\mathfrak{F})$.

Then $\mathfrak{F} \not\models \chi(\mathfrak{F})$.

Therefore, $\chi(\mathfrak{X}) \notin \text{Log}(\mathbb{T}^1)$ and $\chi(\mathfrak{X}) \in \text{Log}(\mathbb{T}^2)$.
 Suppose $\chi(\mathfrak{X}) \notin \text{Log}(\mathbb{T}^2)$. Now we show that $\chi(\mathfrak{X}) \in \text{Log}(\mathbb{T}^2)$.
 Then there is $\mathfrak{B} \in \mathbb{T}^2$ such that $\mathfrak{B} \neq \chi(\mathfrak{X})$. This means that \mathfrak{X} is a d -morphic image of a generated subframe of \mathfrak{B} . Hence, $\mathfrak{X} \leq \mathfrak{B}$ which contradicts the fact that Δ forms a \leq -antichain.
 Therefore, $\chi(\mathfrak{X}) \notin \text{Log}(\mathbb{T}^1)$ and $\chi(\mathfrak{X}) \in \text{Log}(\mathbb{T}^2)$. Thus, $\text{Log}(\mathbb{T}^1) \neq \text{Log}(\mathbb{T}^2)$.

Properties of the Jančov chain, 2

Corollary 93. There are continuum many intermediate logics.

Continuum many logics

