

Fourier Analysis

Jan Wiergerinck

version March 12, 2010

KORTEWEG – DE VRIES INSTITUUT, UNIVERSITEIT VAN AMSTERDAM, PLANTAGE
MUIDERGRACHT 24, 1018 TV, AMSTERDAM

E-mail address: janwieg@science.uva.nl

Contents

Chapter 1. Classical Fourier series	1
1.1. Introduction and Reminder	1
1.2. Sine versus Cosine Series	3
1.3. Weak Topologies	4
1.4. Lacunary Series	5
1.5. Riesz products	6
1.6. Sidon sets	9
1.7. Exercises	11
1.8. Final remarks, notes, and references	12
Chapter 2. Distributions and their Fourier Series	13
2.1. Introduction	13
2.2. Smooth functions on T	14
2.3. Distributions on T	15
2.4. Operations on Distributions	16
2.5. Fourier Series of Periodic Distributions	18
2.6. Convolution and Multiplication	19
2.7. Exercises	19
Chapter 3. Distributions on \mathbb{R}^n	21
3.1. Smooth Functions	21
3.2. Distributions	24
3.3. Distributions with Compact Support	26
3.4. Convolutions and Product Spaces	28
3.5. Tempered distributions	30
3.6. Exercises	30
3.7. Final remarks	31
Chapter 4. The Fourier Transform	33
4.1. Fourier Transform on \mathcal{S}	33
4.2. Fourier Transform on \mathcal{S}'	34
4.3. The Paley-Wiener-Schwartz Theorem	36
4.4. Fourier Transform and other Function Spaces	38
4.5. Exercises	40
Chapter 5. (Micro) Local Analysis	43
5.1. The Wave Front Set	43
5.2. The windowed Fourier Transform on \mathbb{R}	46
5.3. The Continuous Wavelet Transform	46
5.4. The mathematical zoom	48
5.5. Discrete Wavelets: Frames	50
5.6. Multiresolution Analysis, Orthogonal Bases of Wavelets	52
Bibliography	55
Index	57

Fourier Analysis can indicate the study of Fourier transformations, Fourier series, and their extensions. One studies e.g. the convergence properties of Fourier series of certain classes of functions. It may also indicate the use of the Fourier mechanism in other subjects, e.g. in differential equations or in signal analysis. This course contains something of both these worlds. The course is intended for the master level mathematics at Dutch universities. Thus we assume some knowledge of elementary Fourier Analysis, Functional Analysis and Integration theory. At the UvA the courses Integration theory and Linear analysis of the bachelor program are more than sufficient. For convenience of the reader the essentials of all this are mentioned in the notes, but we do not dwell on the proofs.

The notes are based on a course that I first gave in 1996 at the UvA. They have been modified and slightly been edited, but there are bound to be many typo's and other errors. I would certainly appreciate it if a reader pointed out some to me.

Everything in these notes can be found in the literature, but one may have to look for it a while.

Jan Wiegerinck

Classical Fourier series

1.1. Introduction and Reminder

In this section we recall a few facts from the linear analysis course of the third year. See the notes by Prof. Koornwinder [7] or the book [12]. As usual we identify functions on the unit circle T with on \mathbb{R} ; if f is defined on T , then $g(t) = f(e^{it})$ is the associated 2π -periodic function on \mathbb{R} . We thus denote either by $L^p_{2\pi}$ or by $L^p(T)$, ($1 < p < \infty$) the set of 2π -periodic measurable functions that satisfy

$$(1.1.1) \quad \|f\|_p \stackrel{\text{def}}{=} \left(\int_0^{2\pi} |f(t)|^p \frac{dt}{2\pi} \right)^{1/p} < \infty$$

Notice that we normalized $L^p(T)$ spaces, by taking them with respect to the measure $\frac{dt}{2\pi}$. The pleasant effect is that the norm of the function 1 equals 1. We know that $\|\cdot\|_p$ turns L^p into a Banach space, while $L^2(T)$ is even a Hilbert space.

Other function spaces on T are $L^\infty(T)$ the space of 2π periodic, essentially bounded measurable functions, and $C(T) \subset L^\infty(T)$ the space of continuous 2π -periodic functions. Both spaces are Banach spaces when equipped with the sup-norm.

We will also have use for sequence spaces: $l^p(\mathbb{Z})$ ($1 < p < \infty$) is the space of sequences

$$(1.1.2) \quad \{a = \{a_j\}_{j=-\infty}^\infty : \|a\|_p \stackrel{\text{def}}{=} \left(\sum_{j \in \mathbb{Z}} |a_j|^p \right)^{1/p} < \infty\}$$

Again $l^p(\mathbb{Z})$ is a Banach space with norm $\|\cdot\|_p$, and $l^2(\mathbb{Z})$ is a Hilbert space. Other sequence spaces that we will meet are the space of bounded sequences $c(\mathbb{Z})$, and its subspace $c_0(\mathbb{Z})$, which consists of sequences $\{a_j\}$ tending to 0 if $|j| \rightarrow \infty$.

The *Fourier series* of $f \in L^1(T)$ is

$$(1.1.3) \quad \sum_{-\infty}^{\infty} a_n e^{int},$$

where $a_n = \hat{f}(n) = \int_0^{2\pi} f(s) e^{-ins} \frac{ds}{2\pi}$ are the *Fourier coefficients* of f . We know by the *Riemann-Lebesgue Lemma* that $\hat{f}(n) \rightarrow 0$ if $|n| \rightarrow \infty$, that is, $\hat{f} \in c_0(\mathbb{Z})$. A formal sum of the form (1.1.3) with arbitrary a_n is called a trigonometric series. If (1.1.3) arises from a measure, i.e.

$$a_n = \hat{\mu}(n) = \int_{-\pi}^{\pi} e^{-int} d\mu(t),$$

then the $a_n = \hat{\mu}(n)$ are of course the Fourier coefficients of μ and the series (1.1.3) is called a *Fourier-Stieltjes series*. Of course $|\hat{\mu}(n)| \leq \|\mu\|$, but it is in general not true that $\hat{\mu}(n) \rightarrow 0$, if $|n| \rightarrow \infty$. Taking $\mu = \delta$, point mass at 0, we find $\hat{\delta}(j) = 1/2\pi$ for all j .

We see that taking a Fourier series can be seen as a map $\hat{\cdot}$ from a space of functions, or measures, or more general, to a space of sequences. Natural question are: For what kind of things can one define a Fourier series? Can you say something about the target space if you start in $L^p(T)$? Is $\hat{\cdot}$ surjective to some $l^p(\mathbb{Z})$? Is it maybe even an isometry? Is (1.1.3) convergent in L^p if $f \in L^p$? Is it perhaps convergent in any other sense?

Some of these questions will be answered in the course.

The *Dirichlet kernels* D_N are defined as follows

$$(1.1.4) \quad D_N(t) = \sum_{-N}^N e^{int} = \begin{cases} \frac{\sin((N+1/2)t)}{\sin(t/2)} & \text{if } t \notin 2\pi\mathbb{Z} \\ 2N+1 & \text{if } t \in 2\pi\mathbb{Z}. \end{cases}$$

The N -th *Fejér kernel* K_N is defined by

$$(1.1.5) \quad \begin{aligned} K_N(t) &= \frac{1}{N+1} \sum_{n=0}^N D_n(t) = \sum_{n=-N}^N \frac{N+1-|n|}{N+1} e^{int} \\ &= \begin{cases} \frac{1}{N+1} \left(\frac{\sin((N+1)t/2)}{\sin(t/2)} \right)^2 & \text{if } t \notin 2\pi\mathbb{Z} \\ N+1 & \text{if } t \in 2\pi\mathbb{Z}. \end{cases} \end{aligned}$$

For the N -th partial sum $S_N[f](t) = \sum_{n=-N}^N \hat{f}(n)e^{int}$ of the Fourier series of f we find

$$S_N[f](t) = \int_0^{2\pi} f(s) D_N(t-s) \frac{ds}{2\pi} =: f * D_N(t).$$

Similarly, for the N -th *Césaro sum*, i.e. the average of the partial sums S_0 upto S_N ,

$$\sigma_N[f](t) = \frac{1}{N+1} \sum_{n=0}^N S_n[f](t) = f * K_N(t).$$

For $f, g \in L^2(T)$ Parseval's formula holds:

$$(1.1.6) \quad \int_{-\pi}^{\pi} f \bar{g} \frac{dt}{2\pi} = \sum_{\mathbb{Z}} \hat{f}(n) \bar{\hat{g}}(n),$$

Observe that the lefthand side of 1.1.6 is the inner product in $L^2(T)$, while the righthand side is the inner product in $l^2(\mathbb{Z})$. So 1.1.6 expresses that $\hat{\cdot}$ is an isometry from $L^2(T) \rightarrow l^2(\mathbb{Z})$. In fact it is also surjective. If $\{a_j\}_j \in l^2(\mathbb{Z})$ then the partial sums $\sum_{j=-N}^N a_j e^{ijt}$ form a Cauchy sequence in $L^2(T)$ that converges to some $f \in L^2(T)$ with $\hat{f}(j) = a_j$.

The Fejér kernels K_N are *good kernels*, they have the three characteristic properties of an approximate identity: (a) $K_N \geq 0$. (b) $\int_0^{2\pi} K_N(t) \frac{dt}{2\pi} = 1$. (c) For every $0 < \delta < \pi$, $K_N(t) \rightarrow 0$ as $N \rightarrow \infty$ uniformly on $[\delta, 2\pi - \delta]$. If a family of integral kernels L_N on $[0, 2\pi]$ has these three properties, then for $f \in L^p(T)$ ($1 \leq p < \infty$), $L_N * f$ tends to f in L^p sense as $N \rightarrow \infty$. At a point of continuity a of f one has that $L_N * f(a) \rightarrow f(a)$ and, moreover, for $f \in C(T)$ the convergence of $L_N * f$ to f is uniform.

In particular these things hold for the Fejér kernel, giving the well-known convergence result for the Césaro sums. Every $f \in C(T)$ can be approximated uniformly by goniometric polynomials, e.g. by its Césaro sums.

$$(1.1.7) \quad f(t) = \lim_{N \rightarrow \infty} \sigma_N[f](t).$$

As a consequence, and because $C(T)$ is dense in $L^2(T)$ the exponentials e^{int} , $n \in \mathbb{Z}$, form an orthogonal *basis* of $L^2(T)$. We will need a few additional estimates on K_N :

$$(1.1.8) \quad K_N(t) \leq \min\left\{N+1, \frac{\pi^2}{(N+1)t^2}\right\}$$

and, using Parseval's formula,

$$(1.1.9) \quad \|K_N\|_2^2 = \sum_{n=-N}^N \left(\frac{N+1-|n|}{N+1} \right)^2 \geq N/2.$$

The sum in the middle can of course be computed, but the last estimate follows easily by comparison with $\int_0^{N+1} (1-x/(N+1))^2 dx$.

Convergence of the Fourier series is not nearly as good. The classical result is as follows.

THEOREM 1.1.1. *Suppose that $f \in C(T)$ is Lipschitz continuous, i.e. there exist $\alpha, C > 0$ such that*

$$|f(s) - f(t)| < C|s - t|^\alpha.$$

Then

$$S_N[f](t) \rightarrow f(t), \quad \text{uniformly, as } N \rightarrow \infty.$$

1.2. Sine versus Cosine Series

If we start with an *even function* f , ($f(t) = f(-t)$), in (1.1.3), we will find that $a_n = a_{-n}$ (the sequence is *even*). Then taking together the n -th and $-n$ -th term, we obtain

$$a_0 + \sum_{n=1}^{\infty} a_n (e^{int} + e^{-int}) = a_0 + \sum_{n=1}^{\infty} 2a_n \cos(nt),$$

a *cosine series*. Similarly if f is an *odd function* ($f(t) = -f(-t)$), we find $a_n = -a_{-n}$, (the sequence is *odd*), and we obtain the *sine series*

$$\sum_{n=1}^{\infty} a_n (e^{int} - e^{-int}) = \sum_{n=1}^{\infty} 2ia_n \sin(nt).$$

Notice that every function on $(0, \pi)$ can be extended as an even, but also to an odd 2π -periodic function. In that light it is remarkable that sine and cosine series have different convergence behavior even if the coefficients are the same. (Of course the series belonging to even and odd continuation do not have the same coefficients) In the present section we prove two theorems which enlighten this behavior of sine and cosine series.

THEOREM 1.2.1. *Suppose that $(a_n)_{n=1}^{\infty}$ is an even sequence of positive numbers which tend to 0 if $|n| \rightarrow \infty$. If (a_n) satisfies the convexity condition*

$$a_{n-1} + a_{n+1} - 2a_n \geq 0, \quad (n \geq 1),$$

then there exists $f \in L^1_{2\pi}$ such that $\hat{f}(n) = a_n$.

PROOF. The convexity condition implies that $a_n - a_{n+1}$ is monotonically decreasing to 0. From this we have

$$(1.2.1) \quad \begin{aligned} n(a_n - a_{n+1}) &\leq (a_k - a_{k+1}) + (a_{k+1} - a_{k+2}) \dots (a_n - a_{n+1}) + (k-1)(a_n - a_{n+1}) \\ &= a_k - a_{n+1} + (k-1)(a_n - a_{n+1}) \rightarrow 0, \end{aligned}$$

by choosing k fixed and large, so that a_k is small, and then letting $n \rightarrow \infty$. By cleverly rearranging of the series, so-called *summation by parts*, we also find

$$(1.2.2) \quad \begin{aligned} \sum_{n=1}^N n(a_{n-1} + a_{n+1} - 2a_n) &= \sum_{n=1}^N -n((a_n - a_{n+1}) - (a_{n-1} - a_n)) \\ &= a_0 - a_N - N(a_N - a_{N+1}) \rightarrow a_0, \end{aligned}$$

if $N \rightarrow \infty$. Put

$$f_N(t) = \sum_{n=1}^N n(a_{n-1} + a_{n+1} - 2a_n)K_{n-1}(t).$$

This series has non-negative terms and converges in L^1 sense. For $N > M$

$$(1.2.3) \quad \begin{aligned} \int_0^{2\pi} |f_N - f_M| dt &= \int_0^{2\pi} \sum_{n=M+1}^N n(a_{n-1} + a_{n+1} - 2a_n) \int_0^{2\pi} K_{n-1}(t) dt \\ &= \sum_{n=M+1}^N n(a_{n-1} + a_{n+1} - 2a_n) < \varepsilon, \end{aligned}$$

because of (1.1.5) and (1.2.2), therefore $\lim_N \rightarrow \infty f_N = f \in L^1(T)$. Using again (1.1.5) and a dilated version of (1.2.2), we compute \hat{f} .

$$\hat{f}(p) = \sum_{n>|p|} (a_{n-1} + a_{n+1} - 2a_n)(n - |p|) = \sum_{j=1}^{\infty} j(a_{|p|+j-1} + a_{|p|+j+1} - 2a_{|p|+j}) = a_{|p|}.$$

□

THEOREM 1.2.2. *Suppose that f is in $L^1(T)$ and that $\hat{f}(n) = -\hat{f}(-n) \geq 0$ for $n \geq 0$. Then $\sum_{n \neq 0} \frac{\hat{f}(n)}{n}$ converges.*

PROOF. Let $F(t) = \int_0^t f(s) ds$, $t \in [-\pi, \pi]$. Then F is continuous and $F(-\pi) = F(\pi)$, because $\hat{f}(0) = 0$, i.e. $F \in C(T)$. The Fourier coefficients of F are

$$\hat{F}(n) = \frac{\hat{f}(n)}{in}, \quad (n \neq 0).$$

The Césaro sums of F will converge uniformly to F , therefore, subtracting $\hat{F}(0)$ from F and evaluating at 0,

$$\lim_{N \rightarrow \infty} \sum_{1 \leq |n| \leq N} \frac{N+1-|n|}{N+1} \frac{\hat{f}(n)}{n} = i(F(0) - \hat{F}(0)) = -i\hat{F}(0).$$

All terms in the sum are positive, so this sum converges absolutely. Next

$$\sum_{1 \leq |n| \leq N} \frac{\hat{f}(n)}{n} \leq 2 \sum_{1 \leq |n| \leq N} \left(1 - \frac{|n|}{2N+1}\right) \frac{\hat{f}(n)}{n} < i\hat{F}(0),$$

which proves the theorem. □

COROLLARY 1.2.3. *Let $b_n = \frac{1}{\log(n+2)}$, then $\sum b_n \cos(nt)$ is the Fourier series of an L^1 function, but $\sum b_n \sin(nt)$ is not.*

1.3. Weak Topologies

Occasionally we will use weak-* convergence of measures. In this section we recall this notion for readers who are not familiar with it.

1.3.1. Weak-* convergence. A sequence of Borel measures $(\mu_j)_j$ on a compact Hausdorff space X converges weak-* to μ if for every $f \in C(X)$

$$(1.3.1) \quad \lim_{j \rightarrow \infty} \int f d\mu_j = \int f d\mu.$$

Similarly, in a Hilbert space H with inner product $\langle \cdot, \cdot \rangle$, a sequence f_j converges weakly to f if for every $g \in H$

$$(1.3.2) \quad \lim_{j \rightarrow \infty} \langle f_j, g \rangle = \langle f, g \rangle.$$

This and the Banach-Alaoglu Theorem below is basically all we need; nevertheless we give some background.

1.3.2. The weak topology. Recall that a topology τ_1 on a set X is called *weaker* than τ_2 on X if every τ_1 open set in X is also τ_2 open; then τ_2 is called *stronger* than τ_1 . Also recall that the product topology is defined by requiring that it is the weakest topology on the set theoretical product such that all projections are continuous mappings.

We can do something similar in topological vector spaces. Thus let X be a topological vector space such that its dual X^* separates points of X , i.e. for every $x \in X$ there exists a continuous linear functional $L \in X^*$ with $Lx \neq 0$. This is certainly the case if X is a Banach or a Hilbert space. The *weak topology* on X is the weakest topology that makes all

$L \in X^*$ continuous. Since they are already continuous in the original topology of X , the weak topology is weaker than the original one.

A local subbasis for the weak topology on X consists of the sets

$$V_L^\varepsilon = \{x \in X : |Lx| < \varepsilon\},$$

where $\varepsilon > 0$ and $L \in X^*$. This means that $U \subset X$ is a neighborhood of 0 if there exist $\varepsilon_i > 0$, $L_i \in X^*$ such that $(\cap_{i=1}^p V_{L_i}^{\varepsilon_i}) \subset U$. How does this relate to convergence? Well, $x_j \rightarrow x$ if and only if $x_j - x \rightarrow 0$, that is, every neighborhood U of 0 must, for sufficiently large j , contain $x_j - x$. Therefore, for every choice of finitely many $V_{L_i}^{\varepsilon_i}$, it holds that $x_j - x \in (\cap_{i=1}^p V_{L_i}^{\varepsilon_i})$ if j is sufficiently large. This happens if and only if $Lx_j \rightarrow Lx$ for every L . Compare this to (1.3.2).

1.3.3. The weak-* topology. Recall that X can be seen as a subset of X^{**} via $xL := Lx$, ($x \in X$, $L \in X^*$) and that the subset $X \subset X^{**}$ already separates points on X^* . The *weak-* topology* on X^* is defined as the weakest topology that makes all $x \in X$ continuous functionals on X^* . We do not require continuity of functionals in $X^{**} \setminus X$ (In many important cases, however, this set is empty, compare [13]).

Finally we quote

THEOREM 1.3.1 (Banach-Alaoglu). *If V is a neighborhood of 0 in a topological vector space X and*

$$K_V = \{L \in X^* : |Lx| \leq 1 \text{ for every } x \in V\}.$$

Then K_V is weak- compact.*

A proof can be found in [9].

EXAMPLE 1.3.2. Let $X = C(T)$, $V = \{f \in C(T) : \|f\|_\infty < 1\}$, then $K_V = \{\mu \in M(T) : \|\mu\| \leq 1\}$ is compact. Theorem 1.3.1 tells us that every sequence of Borel measures $(\mu_\alpha)_\alpha$ on T with uniformly bounded mass has a weak-* convergent subsequence. In other words, there exists a subsequence $(\mu_j)_j$ and a measure $\mu \in M(T)$ such that (1.3.1) holds.

1.4. Lacunary Series

DEFINITION 1.4.1. A sequence $\{\lambda_j\}$, $j = 1, 2, \dots$, of positive integers is called (Hadamard-) *lacunary* with constant $q > 1$ if $\lambda_{j+1} > q\lambda_j$ for all $j \geq 1$. A power series is called lacunary if it is of the form $\sum c_j z^{\lambda_j}$, while a trigonometric series is called lacunary if it is of the form $\sum c_j e^{i\lambda_j t} + \sum d_j e^{-i\lambda_j t}$ with $\{\lambda_j\}$ lacunary.

LEMMA 1.4.2. *Suppose that $f \in L_{2\pi}^1$ and $f(t) = O(t)$ as $t \rightarrow 0$. If*

$$(1.4.1) \quad \hat{f}(j) = 0, \quad \text{for all } 1 \leq |n_0 - j| \leq 2N,$$

then

$$|\hat{f}(n_0)| \leq 2\pi^4 (N^{-1} \sup_{|t| \leq N^{-1/4}} |f(t)/t| + N^{-2} \|f\|_1).$$

PROOF. If g_N is any trigonometric polynomial of degree $2N$ with $\hat{g}(0) = 1$, then

$$\hat{f}(n_0) = \int_{-\pi}^{\pi} e^{-in_0 t} f(t) g_N(t) \frac{dt}{2\pi},$$

because (1.4.1) expresses that $S_N[e^{-in_0 t} f(t)] = \hat{f}(n_0)$. We take $g_N = K_N^2 / \|K_N\|_2^2$. Then in view of (1.1.5) and (1.1.8) $\int_{-\pi}^{\pi} g_N \frac{dt}{2\pi} = 1$, $g_N \geq 0$, $g_N(t) \leq \frac{\pi^4 2}{N(N+1)^2 t^4}$. We use this to estimate

$$|\hat{f}(n_0)| \leq \int_{-\pi}^{\pi} |f(t)| g_N(t) \frac{dt}{2\pi} = \int_{|t| \leq N^{-1}} + \int_{N^{-1} \leq |t| \leq N^{-1/4}} + \int_{N^{-1/4} \leq |t| \leq \pi}.$$

Now these three integrals are estimated.

$$\begin{aligned}
\int_{|t| \leq N^{-1}} |f(t)| g_N(t) \frac{dt}{2\pi} &\leq \frac{1}{N} \sup_{|t| \leq N^{-1}} \frac{|f(t)|}{|t|} \int_{-\pi}^{\pi} g_N(t) \frac{dt}{2\pi} = \frac{1}{N} \sup_{|t| \leq N^{-1}} \frac{|f(t)|}{|t|}. \\
\int_{\frac{1}{N} \leq |t| \leq \frac{1}{N^{1/4}}} |f(t)| g_N(t) \frac{dt}{2\pi} &\leq \sup_{|t| \leq \frac{1}{N^{1/4}}} \frac{|f(t)|}{|t|} \int_{\frac{1}{N} \leq |t| \leq \frac{1}{N^{1/4}}} \frac{|t| \pi^4 2}{N(N+1)^2 t^4} \frac{dt}{2\pi} \\
(1.4.2) \qquad \qquad \qquad &\leq \sup_{|t| \leq \frac{1}{N^{1/4}}} \frac{|f(t)|}{|t|} \frac{\pi^4 2N}{2\pi(N+1)^2}. \\
\int_{N^{-1/4} \leq |t| \leq \pi} |f(t)| g_N(t) \frac{dt}{2\pi} &\leq \frac{\pi^4 2}{N(N+1)^2 (N^{-1/4})^4} \int_{-\pi}^{\pi} |f(t)| \frac{dt}{2\pi}.
\end{aligned}$$

These three estimates prove the lemma. \square

COROLLARY 1.4.3. *Suppose that $f = \sum_{n=1}^{\infty} a_n \cos(\lambda_n t) \in L^1_{2\pi}$, with $\lambda_{n+1} \geq q\lambda_n$ and $q > 1$. If f is differentiable at a point p then $a_n = o(\lambda_n^{-1})$.*

PROOF. Considering $f_p(t) = f(t+p)$ which has $\hat{f}_p(n) = e^{inp} \hat{f}(n)$ we may assume that $p = 0$. Replace f by $f - f(0) - f'(0) \sin t$. This has no effect on the tail of the series and now $f(t) = o(|t|)$ at 0. We have $\hat{f}(j) = 0$ for $|j - \lambda_n| < \lambda_n/q$. We apply the Lemma and obtain

$$|\hat{f}(\lambda_n)| \leq \frac{o(1)}{\lambda_n} + \frac{C}{\lambda_n^2} = \frac{o(1)}{\lambda_n}.$$

\square

COROLLARY 1.4.4 (Weierstrass' nowhere differentiable function). .

$$f(t) = \sum_{n=0}^{\infty} \frac{\cos(2^n t)}{2^n}$$

is continuous, but nowhere differentiable.

PROOF. The series is lacunary and uniformly convergent, so f is continuous and the previous corollary gives that f is nowhere differentiable. \square

1.5. Riesz products

Let $\{\lambda_n\}$ be lacunary with $q \geq 3$. A trigonometric polynomial of the form

$$P_N(t) = \prod_{n=1}^N (1 + a_n \cos(\lambda_n t + \varphi_n))$$

is called a (finite) *Riesz product*. Observe that, since $q \geq 3$, an integer M can at most in **one** way be written as

$$M = \sum_1^{\infty} c_n \lambda_n, \quad c_n \in \{-1, 0, 1\}.$$

In fact M will be a finite sum and unless $q = 3$, not all M can be expressed as such a sum. We use this when expanding P_N . A typical factor of P_N is $1 + (a_n e^{i\varphi_n}/2) e^{i\lambda_n t} + (a_n e^{-i\varphi_n}/2) e^{-i\lambda_n t}$. In the expansion of P_N we will thus find exponentials of the form $e^{ikt} = e^{i(\sum c_n \lambda_n)t}$ and by the preceding observation such an exponential can be obtained in at most one way. It follows that

$$(1.5.1) \quad \hat{P}(k) = \begin{cases} \prod \left(\frac{a_n e^{i\varphi_n c_n}}{2} \right) & \text{if } k = \sum c_n \lambda_n, \text{ with } c_n \neq 0, \\ 0 & \text{elsewhere.} \end{cases}$$

Also, from $P_{N+1} = P_N + \frac{a_{N+1} e^{i\varphi_{N+1}}}{2} P_N e^{i\lambda_{N+1} t} + \frac{a_{N+1} e^{-i\varphi_{N+1}}}{2} P_N e^{-i\lambda_{N+1} t}$ we see that the Fourier series of P_{N+1} is obtained from the Fourier series of P_N by adding two copies of P_N

multiplied by a constant, one shifted λ_{N+1} to the right, the other shifted λ_{N+1} to the left. As $q \geq 3$ there is no overlap. In particular, whatever the sequence $\{a_n\}$, if $N \rightarrow \infty$, then \hat{P}_N becomes stationary on every finite subset of \mathbb{Z} . We find that $\lim_{N \rightarrow \infty} \hat{P}_N$ is a well-defined trigonometric series. If $P = \lim P_N$ in some sense then $\hat{P} = \lim \hat{P}_N$.

For us there are two cases of interest:

- (1) Suppose $-1 \leq a_n \leq 1$. Then all P_N are nonnegative and $\int_{-\pi}^{\pi} P_N = 1$. Thus the P_N are (densities of) probability measures on $(-\pi, \pi)$. There exists at least one weak-* limit point. If μ_1, μ_2 are two weak-* limit points of $P_N dt$, we have $\hat{\mu}_1 = \hat{\mu}_2$, so $\mu_1 = \mu_2$ and P_N converges weak-* to a probability measure with Fourier-Stieltjes series $\lim P_N$.
- (2) Suppose that $a_n = ib_n, b_n \in \mathbb{R}$ and $\sum_n |b_n|^2 < \infty$. We have $1 \leq |(1+ib_n \cos(\lambda_n t))| \leq (1+b_n^2)^{1/2}$, therefore, with suitable constant C .

$$(1.5.2) \quad 1 \leq |P_N| \leq e^{(\sum_n \log(1+b_n^2))/2} \leq e^{\frac{1}{2} \sum_n b_n^2} < C.$$

Thus $|P_N|$ is uniformly bounded and P_N converges weak-* to (a measure given by) an $L_{2\pi}^\infty$ function. (If O is open in T with Lebesgue measure $|O|$, and f is continuous $0 \leq f \leq 1$ with support in O , then $|\int f P_N dt| \leq C|O|$; the same goes for the weak-* limit, giving that the weak-* limit is absolutely continuous with respect to Lebesgue measure and the density is in L^∞ .)

LEMMA 1.5.1. *Let $\{\lambda_j\}$ be lacunary with constant q . Put $\lambda_{-j} = -\lambda_j, \lambda_0 = 0$. There exist constants A_q, B_q such that if $f(t) = \sum_{-N}^N c_j e^{i\lambda_j t}$, then*

$$(1.5.3) \quad \begin{aligned} \sum |c_j| &\leq A_q \|f\|_\infty, \\ \|f\|_2 &\leq B_q \|f\|_1. \end{aligned}$$

PROOF. Notice that if we prove the Lemma for real valued f , then it follows for complex valued f with the constants A_q and B_q doubled. We first deal with the case $q \geq 3$ and assume that f is real, which means that $c_j = \bar{c}_{-j}$. To prove the first inequality, we set

$$P_N(t) = \prod_{j=1}^N (1 + \cos(\lambda_j t + \varphi_j)).$$

We choose $\varphi_j = \arg c_j, j \geq 1$. We have

$$\int_{-\pi}^{\pi} P_N \bar{f} \frac{dt}{2\pi} = \sum_{j=-N}^N \hat{P}_N(\lambda_j) \bar{f}(\lambda_j) = \frac{1}{2} \sum_{j=-N}^N e^{i \text{Sign}(j) \varphi_{|j|}} \bar{c}_j = \frac{1}{2} \sum_{-N}^N |c_j|,$$

Also

$$\left| \int_{-\pi}^{\pi} P_N \bar{f} \frac{dt}{2\pi} \right| \leq \|f\|_\infty \int_{-\pi}^{\pi} P_N \frac{dt}{2\pi} = \|f\|_\infty.$$

Thus we have proved the first equality for $q \geq 3$ with $A_q = 4$.

For the second inequality we set

$$P_N(t) = \prod_{j=1}^N \left(1 + i \left(\frac{|c_j|}{\|f\|_2} \right) \cos(\lambda_j t + \varphi_j) \right).$$

We proceed as above and find with the same choice of φ_j

$$(1.5.4) \quad \begin{aligned} \|f\|_2 &= \sum_{j=-N}^N \frac{|c_j|^2}{\|f\|_2} = -2i \sum_{j=-N}^N i \frac{|c_j|}{\|f\|_2} \frac{e^{i \text{Sign}(j) \varphi_{|j|}}}{2} \bar{c}_j \\ &= -2i \sum_{j=-N}^N \hat{P}_N(\lambda_j) \bar{f}(\lambda_j) = -2i \int P_N \bar{f} \frac{dt}{2\pi} \leq 2 \|P_N\|_\infty \|f\|_1. \end{aligned}$$

Since the P_N are uniformly bounded by $e^{1/2}$, compare (1.5.2), we are done. Notice that it is the seemingly artificial factor i that we introduced in P_N , that makes it possible to estimate $\|P_N\|_\infty$.

For $q \geq 3$ we may take $B_q = 4e^{1/2}$.

The general case is done by carefully splitting the lacunary sequence in such a way that the Riesz products associated to the subsequences make sense and will only pick up the terms in the series that we want. Let M be a large integer, depending on q and to be determined in the process. Write for a fixed $0 \leq m < M$ $\lambda_j^m = \lambda_{m+jM}$. We want that $\{\lambda_j^m\}_j$ is lacunary with constant ≥ 3 . Thus we require that M satisfy

$$(1.5.5) \quad q^M \geq 3,$$

hence a Riesz product associated to $\{\lambda_j^m\}_j$ makes sense. Next, we want that each of the frequencies λ_k of f occurs in precisely one Riesz product. Suppose that $n > 0$ is written as $\sum_{j=0}^J c_j \lambda_j^m$, with $c_j \in \{-1, 0, 1\}$ and $c_J = 1$. Then

$$|n - \lambda_J^m| \leq \sum_{j=0}^{J-1} \lambda_j^m \leq \lambda_J^m \sum_{j=0}^{J-1} \frac{\lambda_j^m}{\lambda_J^m} \leq \lambda_J^m \sum_{j=1}^J \frac{1}{q^{jM}} \leq \frac{\lambda_J^m}{q^M - 1}.$$

Thus we want $|\lambda_k - \lambda_J^m| > \frac{\lambda_J^m}{q^M - 1}$ for all $\lambda_k \neq \lambda_J^m$. If $\lambda_k \geq q\lambda_J^m$ this leads to

$$(1.5.6) \quad q - 1 > \frac{1}{1 - q^M} \quad \text{or} \quad q > 1 + \frac{1}{q^M - 1},$$

while, if $\lambda_J^m \geq q\lambda_k$, this leads to

$$(1.5.7) \quad 1 - 1/q > \frac{1}{1 - q^M} \quad \text{or} \quad 1/q < 1 - \frac{1}{q^M - 1}.$$

We take M so large that (1.5.5), (1.5.6), (1.5.7) are satisfied. Now let $P_N^m = \prod_1^N (1 + a_{m+jM} \cos(\lambda_j^m t + \varphi_{m+jM}))$ be one of the Riesz products considered in the first part of the proof. Then

$$\frac{1}{2\pi} \int P_N^m(t) \bar{f}(t) dt = \frac{1}{2} \sum |a_{m+jM}| |c_{m+jM}|.$$

The first part of the proof gives

$$(1.5.8) \quad \begin{aligned} \sum |c_{m+jM}| &\leq 4\|f\|_\infty, \\ \left(\sum |c_{m+jM}|^2\right)^{\frac{1}{2}} &\leq 4e^{1/2}\|f\|_1, \end{aligned}$$

in respectively the first and second case of the lemma. Summing over $m = 1, \dots, M$ gives the result. \square

THEOREM 1.5.2. *Suppose that the Fourier series $\sum_{-\infty}^{\infty} c_j e^{i\lambda_j t}$ of $f \in L^1(T)$ is lacunary, then $f \in L^2(T)$. If f is bounded, then $\sum |c_j| < \infty$.*

PROOF. Let $\sigma_N[f]$ be a Césaro sum. These have L^1 norms, uniformly bounded by $\|f\|_1$. The lemma gives that

$$\sum_{-N}^N \left(1 - \frac{|j|}{N+1}\right)^2 |c_j|^2 \leq B_q M.$$

This implies, by letting $N \rightarrow \infty$, that for fixed J the sum $\sum_{-J}^J |c_j|^2 \leq B_q M$, thus $f \in L^2(T)$. The proof of the second statement is similar. \square

The homogeneity of behavior of lacunary series also appears in the following theorem. We shall denote the length of a subarc Γ of T by $|\Gamma|$.

THEOREM 1.5.3. *Suppose that (λ_j) is lacunary with constant q . For every $\delta > 0$ there exists $j_0 \in \mathbb{N}$ such that for all lacunary $f \in L^2(T)$, $f(t) = \sum_{-\infty}^{\infty} c_j e^{i\lambda_j t}$ with $c_j = 0$ for $|j| < j_0$, the following inequality holds for every subarc Γ of T :*

$$\left(\frac{1}{2\pi}|\Gamma| - \delta\right)\|f\|^2 \leq \int_{\Gamma} |f|^2 \frac{dt}{2\pi} \leq \left(\frac{1}{2\pi}|\Gamma| + \delta\right)\|f\|^2.$$

PROOF. It suffices to prove the theorem for trigonometric polynomials. We shall give a proof for $q \geq 3$. For the general case, see exercise 1.7.7. We have

$$(1.5.9) \quad \int_{\Gamma} |P|^2 \frac{dt}{2\pi} = \int_{\Gamma} \sum_{n,m} c_n \bar{c}_m e^{i(\lambda_n - \lambda_m)t} \frac{dt}{2\pi} = \frac{1}{2\pi} |\Gamma| \|P\|^2 + \sum_{n \neq m} c_n \bar{c}_m \int_{\Gamma} e^{i(\lambda_n - \lambda_m)t} \frac{dt}{2\pi}.$$

Let h be the characteristic function of Γ . Then $\hat{h}(n) = \int_{\Gamma} e^{-int} dt / 2\pi$, therefore (1.5.9) can be rewritten as

$$(1.5.10) \quad \frac{1}{2\pi} |\Gamma| \|P\|^2 + \sum_{n \neq m} c_n \bar{c}_m \hat{h}(\lambda_m - \lambda_n).$$

Now because $q \geq 3$, $\lambda_n - \lambda_m$ assumes any integer j at most twice (solutions occur in pairs: (n, m) and $(-m, -n)$, and compare the beginning of this section). For general $q > 1$ there exists $K = K(q)$ such that there are at most K solutions, Ex. 1.7.7. Moreover, $\min_{m, n \geq j_0} |\lambda_m - \lambda_n|$ tends to ∞ with j_0 . Thus frequencies j of \hat{h} occur at most twice in the sum in (1.5.10). We apply Cauchy-Schwarz to the sum and obtain that the norm of the sum is less than

$$\left(\sum_{n \neq m} |c_n c_m|^2 \sum_{|j| \geq \inf_{n \neq m} \{|\lambda_n - \lambda_m|\}} |\hat{h}(j)|^2 \right)^{\frac{1}{2}} \leq \|P\|^2 \delta,$$

if we choose j_0 so large that the 2-norm of the tail of the series of h is less than $\delta/2$. It follows from the explicit form of $\hat{h}(j)$ that this can be done independent of Γ . Combining this with (1.5.9) and (1.5.10) we obtain

$$\left(\frac{1}{2\pi}|\Gamma| - \delta\right)\|P\|^2 \leq \int_{\Gamma} |f|^2 dt \leq \left(\frac{1}{2\pi}|\Gamma| + \delta\right)\|P\|^2.$$

□

1.6. Sidon sets

DEFINITION 1.6.1. Let $E \subset \mathbb{Z}$. A function f (or measure, or distribution) on T is called *E-spectral* if $\hat{f}(n) = 0$ if $n \notin E$. Denote by C_E , L_E^p , M_E the respective subspaces of $C(T)$, $L^p(T)$, $M(T)$ consisting of *E-spectral* elements. These are closed subspaces. A subset E of \mathbb{Z} is called a *Sidon set* if $f \in C_E$ implies $\hat{f} \in l^1(E)$.

EXAMPLE 1.6.2. Of course every finite set is a Sidon set. By Theorem 1.5.2 every lacunary set is a Sidon set.

THEOREM 1.6.3. *The following are equivalent:*

1. E is a Sidon set.
2. There exists $K > 0$ such that $\|\hat{f}\|_1 \leq K\|f\|_{\infty}$ for all *E-spectral* trigonometric polynomials f .
3. $\|\hat{f}\|_1$ is bounded for every $f \in L_E^{\infty}$.
- 3.a There exists a K such that $\|\hat{f}\|_1 \leq K\|f\|_{\infty}$ for every $f \in L_E^{\infty}$.
4. $\widehat{M}|_E = l^{\infty}(E)$.
5. $\widehat{L}^1|_E = c_0(E)$.

PROOF. (1 \implies 2) If E is Sidon, then the map $f \mapsto \hat{f}$ is linear bijective from C_E to $l^1(E)$. Also its inverse (as a linear map) is continuous. Indeed

$$\|f\|_\infty = \sup_t \left| \sum \hat{f}(n)e^{int} \right| \leq \sum |\hat{f}(n)|.$$

Hence, by the Open Mapping Theorem, $\hat{f} \mapsto f$ is open and thus $f \mapsto \hat{f}$ is continuous, which means that for some $K > 0$ one has $\|\hat{f}\|_1 \leq K\|f\|_\infty$. In particular, this is true for E -spectral trigonometric polynomials.

(2 \implies 3(a)) If $f \in L_E^\infty$, then $\sigma_N(f)$ is an E -spectral trigonometric polynomial with $|\sigma_N(f)|_\infty \leq \|f\|_\infty$, independent of N . Thus there is a constant K such that $\|\hat{\sigma}_N(f)\|_1 \leq K\|f\|_\infty$. As in the proof of Theorem 1.5.2 we conclude that $\|\hat{f}\|_1 \leq K\|f\|_\infty$.

(3 \implies 1) is trivial.

Item 4 and 5 both say that a certain bounded linear transformation is surjective. Now recall that the Open Mapping Theorem implies that a surjective bounded linear transformation F between Banach spaces X and Y is *open*. Thus $F(\{\|x\| < 1\})$ contains an open neighborhood of $0 \in Y$. By linearity we obtain that there is a constant $C > 0$ such that for every $y \in Y$ there exist $x \in X$ with $Fx = y$ and $\|x\| < C\|y\|$. This will be used in the last two steps of the proof.

(3(a) \implies 4) Let $(d_j)_j \in l^\infty(E)$. Then $f \mapsto \sum_j \hat{f}(j)d_j$ is a continuous linear functional on C_E which by Hahn-Banach can be extended to $C(T)$. By the Riesz Representation Theorem there exists a complex regular Borel measure μ which represents this functional. Thus for $f \in C_E$ we have $\int f d\mu = \sum_j \hat{f}(j)d_j$. We choose $f(t) = e^{i\lambda_j t}$, $\lambda \in E$ and find $\hat{\mu}(-\lambda_j) = d_j$. Thus $\hat{\mu}(\lambda_j) = \bar{d}_j$. Replacing d_j by \bar{d}_j we obtain our result.

(4 \implies 5) First observe that modification of the Fejér kernels yields that if Λ is a finite set of integers and $\varepsilon > 0$, then there exists a trigonometric polynomial $P = P_{\Lambda, \varepsilon}$ such that $\hat{P}(j) = 1$ and $\|P\|_1 \leq 1 + \varepsilon$. We will use this with $\varepsilon = 1$. Now let $(d_j)_j \in c_0(E)$. We may assume that $|d_j| \leq 1$. Put

$$E_k = \{n : 2^{-k} < |d_n| \leq 2^{-k+1}\}.$$

By 4. there exist measures μ_k such that $\hat{\mu}_k(j) = d_j$ if $j \in E_k$, while $\hat{\mu}_k(j) = 0$ if $j \in E \setminus E_k$; moreover $\|\mu_k\| \leq C2^{-k}$. Let T_k be trigonometric polynomials with $\hat{T}_k(j) = 1$ on E_k and $\|T_k\|_1 \leq 2$. Then $T_k * \mu_k$ is a trigonometric polynomial of L^1 norm less than $2C2^{-k}$ and with Fourier coefficients $\hat{T}_k(j)\hat{\mu}_k(j) = \hat{\mu}_k(j)$ on E . The conclusion is that

$$f(t) = \sum_{k=1}^{\infty} T_k * \mu_k$$

is in $L^1(T)$ because the series converges in $L^1(T)$ and has $f(j) = d_j$ on E .

(5 \implies 2) Let g be an E -spectral trigonometric polynomial. Define $d_n = |\hat{g}(n)|/\bar{\hat{g}}(n)$ if $\hat{g}(n) \neq 0$ and $d_n = 0$ elsewhere. $(d_n)_n \in c_0(E)$ of norm 1, hence there exists $f \in L^1(T)$ with $\hat{f}|_E = d_n$ and $\|f\|_1 \leq C$, where C is a constant only depending on E . Then

$$\sum |\hat{g}(n)| = \int_{-\pi}^{\pi} f(t)\bar{g}(t) dt \leq \|g\|_\infty \|f\|_1 \leq C\|g\|_\infty.$$

□

REMARKS 1.6.4. The smallest constant K in Theorem 1.6.3, such that 3a holds, is called the *Sidon constant* of E .

In retrospect, we can understand the (limits of the) Riesz products in Lemma (1.5.1) as explicitly constructed measures of Theorem (1.6.3), statement 4 and 5.

COROLLARY 1.6.5 (to Theorem 1.5.2). *Suppose that $E = \{\lambda_j\}$ is lacunary and that $L = (d_j) \in l^2(E)$. Then there exists a bounded function f on T such that $\hat{f}(\lambda_j) = d_j$.*

PROOF. In view of Riesz Representation Theorem, L is in $l^2(E)^* = (L_E^2)^*$. In fact, $L\varphi = \sum \hat{\varphi}(\lambda_j)\bar{d}_j$. By Theorem 1.5.2 L is also a continuous linear functional on L_E^1 . Now L_E^1 is a closed subspace of $L_1(T)$. By Hahn Banach we can extend L to all of $L^1(T)$. We denote the extension again by L . Recalling, c.f. [13][Ch 7], that the dual space of $L^1(T)$ is $L^\infty(T)$, we find that there exists a function $f \in L^\infty$ such that $L\varphi = \int_0^{2\pi} \varphi \bar{f} \frac{dt}{2\pi}$, for all $\varphi \in L^1(T)$. We now apply this with $\varphi(t) = e^{i\lambda_j t}$ and obtain $\hat{f}(\lambda_j) = d_j$. \square

REMARK 1.6.6. One can do the same for general Sidon sets by proving the analogue of Theorem 1.5.3. This requires considerably more effort, cf. [2].

1.7. Exercises

1.7.1. Let $(b_n)_{n \in \mathbb{Z}}$ be a sequence of positive numbers such that $b_n \rightarrow 0$ as $|n| \rightarrow \infty$. Show that there exists a sequence (a_n) with $a_n \geq b_n$ which satisfies the convexity condition of Theorem 3.1.

1.7.2. Suppose that $f \in C_{2\pi}$ is Lipschitz continuous with exponent α , i.e. there exists $B > 0$ such that for all x , $|f(x+t) - f(x)| \leq B|t|^\alpha$. Show that $|\hat{f}(n)| \leq Cn^{-\alpha}$. Hint: Write the Fourier coefficients as

$$\hat{f} = \frac{1}{2} \left(\int_0^{2\pi} f(s)e^{-ins} \frac{ds}{2\pi} - \int_0^{2\pi} f(s - \pi/n)e^{-ins} \frac{ds}{2\pi} \right).$$

1.7.3. Let (λ_n) be lacunary. Suppose that $f \in L^1(T)$ has Fourier series $\sum a_n \cos(\lambda_n t)$. Show that if f is Hölder continuous at one point $t_0 \in T$, then $a_n = O(\lambda_n^{-\alpha})$. Next show that f is Hölder continuous on T .

1.7.4. With the assumptions as in exercise 1.7.3, show that if f equals 0 on a small interval in T , then $f \in C^\infty(T)$. Can you relax the condition that f be 0 on an interval and still reach the same conclusion?

1.7.5. With the same assumptions as in exercise 1.7.3, $f(t) = \sum c_n e^{i\lambda_n t}$, show that if f is (real) analytic on a subarc Γ of T , then f is real analytic on T by completing the following outline.

- (1) Observe that f is C^∞ because of exercise 1.7.4 extended.
- (2) Choose a clever interval Γ , a suitable modification \tilde{f} of f and show that there is a $C > 0$ such that for every k, l

$$|c_k|^2 |\lambda_k|^{2l} \leq C \int_{\Gamma} |\tilde{f}^{(l)}(t)|^2 dt.$$

- (3) Use the Cauchy estimates to make the integrals in ii. bounded by a constant times $(l!\delta^{-l})^2$.
- (4) Make a favourable choice of l to have $\limsup_{k \rightarrow \infty} |c_k|^{\frac{1}{\lambda_k}} < 1$.
- (5) Finish it off!

1.7.6. Prove *Hadamard's Theorem*: If $f(z) = \sum_{n=0}^{\infty} c_n z^{\lambda_n}$ is a lacunary power series with radius of convergence R then $f(z)$ has no analytic continuation to a domain larger than the disc with radius R . ($C(0, R)$ is a natural boundary.)

1.7.7. Prove Theorem 1.5.3 for general q by showing that now there exists $K > 0$ such that every $j \in \mathbb{Z} \setminus \{0\}$ is assumed at most K times as a value of $\lambda_n - \lambda_m$.

1.7.8. Under the assumptions of Corollary 1.6.5, show that f can in fact be chosen continuous, by completing the following steps.

- (1) Find sequences $(a_n), (b_n)$ such that $d_n = a_n b_n$, with $a_n \in l^2(E)$ and $b_n \in c_0(E)$.
- (2) Show that the convolution of a bounded and an integrable function is continuous.
- (3) Prove the assertion.

1.7.9. Suppose that $f \in C(T)$ has positive Fourier coefficients. Prove that $\hat{f} \in l^1$.

1.7.10. Suppose that for every $d = (d_n)_n \in l^\infty(E)$ with $|d_n| = 1$, there exists a measure $\mu \in M(T)$ with

$$|d_n - \hat{\mu}(n)| \leq 1 - \delta.$$

Prove that E is Sidon. (Think of the previous exercise and let $d_n = \frac{|\hat{f}(n)|}{\hat{f}(n)}$.)

1.8. Final remarks, notes, and references

The classical book on trigonometric series is [14]. Section 1. is in [7], but also can be found e.g. in [6].

Material concerning section 3 can be found in any reasonable book on Functional Analysis, e.g. [9], see [13] for a more extensive list of references.

Section 2. has been taken from [6], but is similarly treated in [14].

Section 4,5 and 6 are also mostly taken from [6]; [2] has a more comprehensive treatment and was also used. Many things can be found in [14] too.

Distributions and their Fourier Series

2.1. Introduction

Consider the wave equation on \mathbb{R}^2

$$(2.1.1) \quad \frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial t^2} = 0.$$

Introducing new coordinates $x' = x + t$, $y' = x - t$, we obtain the equation

$$(2.1.2) \quad \frac{\partial^2 u}{\partial x' \partial y'} = 0,$$

which has the classical solution

$$u(x', y') = f(x') + g(y'), \quad f, g \in C^2(\mathbb{R}).$$

Thus (2.1.1) has as classical solutions $f(x+t) + g(x-t)$. Classically, C^2 is required because 2 differentiations are performed on u . Physically, however, there is no reason to ask much more than continuity. Also, from (2.1.2) we see that $\frac{\partial u}{\partial y'} = \tilde{g}(y')$, where the only thing that matters is that this function doesn't depend on x' . If $\tilde{g} \in L^1(\mathbb{R})$ we get a solution $u(x', y') = f(x') + g(y')$ with no stronger conditions on f and g than continuity. There are of course problems with changing the variables and we have a solution which is not symmetric in x and y . The point is that it is at least inconvenient not to be able to differentiate continuous functions.

As far as Fourier analysis is concerned, we know that $f \in C_{2\pi}^1$ has the property that $\hat{f}'(n) = in\hat{f}(n)$. We can formally write down this sequence of Fourier coefficients also if f is no longer differentiable. Can we give meaning to it as the Fourier series of something interesting? Moreover, consider the Fourier transform $\mathcal{F}f$ of $f \in L^1(\mathbb{R})$. If f is C^1 we know that

$$(2.1.3) \quad \mathcal{F}f'(\xi) = i\xi\mathcal{F}f(\xi).$$

Multiplication with ξ is a well defined operation on functions, the righthand side of (2.1.3) is always well defined. A meaningful lefthand side, that is, indefinite differentiability of L^1 functions, is desirable.

One way out is the concept of weak solution. Notice that if $u \in C^2$ solves (2.1.1), then for every compactly supported $\varphi \in C^\infty(\mathbb{R}^2)$ we have

$$(2.1.4) \quad \iint \left(\frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial t^2} \right) \varphi(x, t) dx dt = \iint u(x, t) \left(\frac{\partial^2 \varphi}{\partial x^2} - \frac{\partial^2 \varphi}{\partial t^2} \right) dx dt = 0.$$

If the last equality in (2.1.4) holds for all compactly supported $\varphi \in C^\infty(\mathbb{R}^2)$ and $u \in C^2$, then u satisfies (2.1.1). However the last integral in (2.1.4) makes sense for locally integrable u . Thus one calls a locally integrable (sometimes only a continuous) u a weak solution of (2.1.1) if u satisfies

$$\iint \left(\frac{\partial^2 \varphi}{\partial x^2} - \frac{\partial^2 \varphi}{\partial t^2} \right) u(x, t) dx dt = 0.$$

We will use this idea to differentiate locally integrable functions arbitrarily often. In fact we will go one step further. Observing that

$$u \mapsto \iint u\varphi \, dxdt$$

is a linear functional on compactly supported C^∞ functions, we will explain how to “differentiate” a large class of such linear functionals, which is determined by a continuity condition, and identify the ones originating from locally integrable functions with a subclass.

2.2. Smooth functions on T

Recall, see [13][Chapter 6], that a *seminorm* on a (complex) vector space X is a real valued function p such that

$$(2.2.1) \quad \begin{aligned} p(x) &\geq 0, \\ p(x_1 + x_2) &\leq p(x_1) + p(x_2), \\ p(\lambda x) &= |\lambda|p(x). \end{aligned}$$

Here $x, x_i \in X$, $\lambda \in \mathbb{C}$.

On $C^\infty(T) = C_{2\pi}^\infty$ we define seminorms

$$(2.2.2) \quad P_j(\varphi) = \|\varphi^{(j)}\|_\infty,$$

the maximum of the j -th derivative. These seminorms can be used to define a topology on $C^\infty(T)$ by requiring that they are *continuous*, just as weak topologies were introduced in Chapter 1. Section 1.3 Thus, a local subbasis at 0 consists of sets

$$V_{j,\varepsilon} = \{\varphi : P_j(\varphi) < \varepsilon\}, \quad j \in \mathbb{N}, \varepsilon > 0.$$

The space $C^\infty(T)$ endowed with this topology is called $\mathcal{D}(T)$. From Chapter 1, Section 1.3 we see that

$$(2.2.3) \quad f_j \in \mathcal{D}(T) \rightarrow f \in \mathcal{D}(T) \iff \forall k \in \mathbb{N} \, f_j^{(k)} \rightarrow f^{(k)} \text{ uniformly.}$$

So it is pretty hard for functions to converge in $\mathcal{D}(T)$, cf. Exercise 2.7.1. However, Fourier series behave nice in $\mathcal{D}(T)$.

LEMMA 2.2.1. *Let $\varphi \in \mathcal{D}(T)$. Then $S_N[\varphi] \rightarrow \varphi$ in $\mathcal{D}(T)$ as $N \rightarrow \infty$.*

PROOF. Because φ is smooth, $S_N[\varphi] \rightarrow \varphi$ uniformly, but also $S_N[\varphi]' = S_N[\varphi'] \rightarrow \varphi'$ uniformly. The same is true for higher derivatives. By (2.2.3) we are done. \square

LEMMA 2.2.2. *The space $\mathcal{D}(T)$ is metrizable and complete in the metric*

$$(2.2.4) \quad d(\varphi, \psi) = \sum_{j=0}^{\infty} 2^{-j} \frac{P_j(\varphi - \psi)}{1 + P_j(\varphi - \psi)}.$$

PROOF. See Exercise 2.7.2. \square

Observe that the metric d is *translation invariant*, that is $d(x, y) = d(x - z, y - z)$ for all $x, y, z \in X$

However, $\mathcal{D}(T)$ with the present topology cannot be turned into a Banach space because of the following Lemma, which says that $\mathcal{D}(T)$ has the *Heine-Borel property*. A Banach space can only possess this property if it is finite dimensional. Recall that a set X in a topological vector space is *bounded* if it has the property that for every open neighborhood U of 0, there exists $N > 0$ such that $X \subset NU$. If the topology is determined by seminorms, this just means that every seminorm is bounded on X .

LEMMA 2.2.3. *Every bounded sequence in $\mathcal{D}(T)$ has a convergent subsequence; A closed bounded set in $\mathcal{D}(T)$ is compact.*

PROOF. This is a simple consequence of the Arzelà-Ascoli Theorem, cf. Exercise 2.7.3. \square

It will be convenient to replace P_n by an equivalent set of seminorms

$$\tilde{P}_n = \sum_{j=0}^n P_j.$$

This doesn't change the topology, but it has the advantage that a local basis at 0 of $\mathcal{D}(T)$ is now given by $\{\tilde{P}_n(\varphi) < 1/n\}$, $n \in \mathbb{N}$.

REMARK 2.2.4. A vector space with topology induced by a complete invariant metric, like we have met here, is called a *Fréchet space*. (Sometimes local convexity is required, but that can be shown to hold here too.)

2.3. Distributions on T

Let $\mathcal{D}' = \mathcal{D}'(T)$ denote the dual space of $\mathcal{D}(T)$, that is the space of continuous linear functionals on $\mathcal{D}(T)$. The notation is classical. Elements of \mathcal{D}' are called (periodic) *distributions* or *generalized functions*. The space \mathcal{D}' will naturally be equipped with the *weak-** topology. We denote the action of $L \in \mathcal{D}'(T)$ on $\varphi \in \mathcal{D}(T)$ usually by

$$\langle L, \varphi \rangle (= L\varphi).$$

The following lemma is an extension of a familiar result for Banach spaces, cf. [13][thm. 3.2].

LEMMA 2.3.1. *Let X be a vector space with a topology induced by a countable set of seminorms $\{p_1, p_2, \dots\}$ and let L be a linear functional on X . The following are equivalent.*

- i. L is continuous on X ,
- ii. L is continuous at $x_0 \in X$,
- iii. There exist $C > 0$, $K \in \mathbb{N}$ such that

$$|Lx| \leq C \max_{i=1, \dots, K} p_i(x).$$

PROOF. (i \implies ii) is trivial.

(ii \implies iii). If L is continuous at x_0 , then for every $\varepsilon > 0$ there exist $K \in \mathbb{N}$ and $\delta_j > 0$, ($j = 1, \dots, K$), such that $p_j(x - x_0) < \delta_j$, ($j = 1, \dots, K$) implies $|L(x - x_0)| < \varepsilon$. Let $\delta = \min_j \{\delta_j\}$ and denote for $y \in X$, $M_y = \max_{i=1, \dots, K} p_i(y)$. Then

$$|Ly| = \left| \frac{M_y}{\delta} L\left(\frac{\delta y}{M_y}\right) \right| < \frac{M_y}{\delta} \varepsilon,$$

which proves iii, with $C = \varepsilon/\delta$.

(iii \implies i). Let $x, y \in X$.

$$|Lx - Ly| = |L(x - y)| < C \max_{i=1, \dots, K} p_i(x - y).$$

This is less than ε if $p_i(x - y) < \varepsilon/KC$. Thus we described a small neighborhood of x which is mapped in an ε -neighborhood of Lx . \square

Apparently a linear functional L on $\mathcal{D}(T)$ is continuous, i.e. a *distribution*, if and only if there exist $n \in \mathbb{N}$ and $C > 0$ such that

$$(2.3.1) \quad |\langle L, \varphi \rangle| \leq C \tilde{P}_n(\varphi), \quad \forall \varphi \in \mathcal{D}(T).$$

The smallest n that is possible in (2.3.1) is called the *order* of the distribution. Of course the zero distribution has order $-\infty$ assigned to it.

REMARK 2.3.2. If the underlying space is not compact, e.g. \mathbb{R} instead of T , distributions may have infinite order.

2.3.1. Examples of distributions.

- (1) Every $u \in L^1_{2\pi}$ defines a distribution L_u via

$$\langle L_u, \varphi \rangle = \int_{-\pi}^{\pi} u \varphi \frac{dt}{2\pi}.$$

This functional is indeed continuous: $|\langle L_u, \varphi \rangle| \leq \|u\|_1 \|\varphi\|_{\infty}$.

- (2) We denote the set of Borel measures on T by $M(T)$. Every measure $\mu \in M(T)$ defines in the same way a distribution L_{μ} via

$$\langle L_{\mu}, \varphi \rangle = \int_{-\pi}^{\pi} \varphi d\mu(t).$$

Both examples are distributions of order 0.

Abusing the language we will drop the L and identify a function or measure with the associated distribution, writing e.g. $\langle u, \varphi \rangle$.

- (3) The delta-distribution $\delta = \delta_0$ is defined by

$$\langle \delta, \varphi \rangle = \varphi(0).$$

The delta distribution originates from point mass at 0.

REMARK 2.3.3. The name test functions for the elements of \mathcal{D} is now understandable, these functions are used to test the action of an L^1 function or distribution.

2.4. Operations on Distributions

- 2.4.1. Differentiation.** Let $L \in \mathcal{D}'(T)$. Define its derivative $L' \in \mathcal{D}'(T)$ by

$$\langle L', \varphi \rangle = -\langle L, \varphi' \rangle.$$

Observe that L' is well-defined because $\varphi' \in \mathcal{D}(T)$; L' is also continuous. Indeed, there exist $C > 0$, $n \in \mathbb{N}$ such that $|\langle L, \varphi \rangle| \leq C \tilde{P}_n(\varphi)$. Then

$$|\langle L', \varphi \rangle| = |\langle L, \varphi' \rangle| \leq C \tilde{P}_n(\varphi') \leq C \tilde{P}_{n+1}(\varphi).$$

We conclude that every distribution is infinitely often differentiable. In general, differentiation increases the order of a distribution by 1.

EXAMPLES 2.4.1. Let $f(x)$ be the characteristic function of $(0, \pi)$ viewed as element of $L^1_{2\pi}$. We compute its distributional derivative f' . Let $\varphi \in \mathcal{D}(T)$, then

$$\langle f', \varphi \rangle = -\langle f, \varphi' \rangle = -\int_{-\pi}^{\pi} f(t) \varphi'(t) \frac{dt}{2\pi} = -\int_0^{\pi} \varphi'(t) \frac{dt}{2\pi} = \frac{\varphi(0) - \varphi(\pi)}{2\pi}.$$

We conclude that $f' = \frac{\delta_0 - \delta_{\pi}}{2\pi}$.

The function $f(x) = \log|x|$ is in $L^1_{2\pi}$. Its distributional derivative is determined by

$$\begin{aligned} \langle f', \varphi \rangle &= -\langle f, \varphi' \rangle = -\int_{-\pi}^{\pi} \log|t| \varphi'(t) \frac{dt}{2\pi} = -\lim_{\varepsilon \rightarrow 0} \int_{-\pi}^{-\varepsilon} \log|t| \varphi'(t) \frac{dt}{2\pi} \\ &+ \int_{\varepsilon}^{\pi} \log|t| \varphi'(t) \frac{dt}{2\pi} \\ (2.4.1) \quad &= \frac{-1}{2\pi} \lim_{\varepsilon \rightarrow 0} \left(\log \varepsilon \varphi(-\varepsilon) - \log \pi \varphi(-\pi) - \log \varepsilon \varphi(\varepsilon) + \log \pi \varphi(\pi) - \right. \\ &\left. \left(\int_{-\pi}^{-\varepsilon} + \int_{\varepsilon}^{\pi} \frac{\varphi(t)}{t} dt \right) \right) = \text{p(ri)ncipal v(alue)} \int \frac{\varphi(t)}{t} \frac{dt}{2\pi}. \end{aligned}$$

Here we have used that φ is periodic, making the stock terms cancel in the limit. The last equality is the definition of the *principal value integral*. We have shown that $f' = \text{p.v.} \frac{1}{x}$.

PROPOSITION 2.4.2. *Differentiation is a continuous operation: If $L_j \rightarrow L$ in $\mathcal{D}'(T)$ then $L'_j \rightarrow L'$ in $\mathcal{D}'(T)$.*

PROOF. For any $\varphi \in \mathcal{D}(T)$, we have $\langle L_j, \varphi \rangle \rightarrow \langle L, \varphi \rangle$ by definition of weak-* convergence. Hence $\langle L_j, \varphi' \rangle \rightarrow \langle L, \varphi' \rangle$, which by definition of differentiation gives $\langle L'_j, \varphi \rangle \rightarrow \langle L', \varphi \rangle$. \square

2.4.2. Restricted Multiplication. The price one pays for infinite differentiability is that multiplication of distributions is in general not possible. However, if $f \in C^\infty(T)$, $L \in \mathcal{D}'$ then fL can be defined by

$$\langle fL, \varphi \rangle = \langle L, f\varphi \rangle.$$

This is indeed continuous: if $\varphi_n \rightarrow \varphi$ in $\mathcal{D}(T)$ then $f\varphi_n \rightarrow f\varphi$ in $\mathcal{D}(T)$, therefore

$$\langle fL, \varphi_n \rangle = \langle L, f\varphi_n \rangle \rightarrow \langle L, f\varphi \rangle = \langle fL, \varphi \rangle.$$

If $\mu \in M(T)$ and $f \in C(T)$, then $f\mu \in M(T)$. Hence multiplication of a continuous function with a distribution associated with a measure or an L^1 function is also possible. More generally one can prove that a distribution of order k can be multiplied with a function in C^k , cf. ex. 2.7.5.

LEMMA 2.4.3 (Product rule). *If $f \in C^\infty(T)$ and $L \in \mathcal{D}'$, then*

$$(fL)' = f'L + fL'.$$

PROOF. Let $\varphi \in \mathcal{D}$. Then

$$\begin{aligned} (2.4.2) \quad \langle (fL)', \varphi \rangle &= -\langle fL, \varphi' \rangle = -\langle L, f\varphi' \rangle \\ &= -\langle L, (f\varphi)' - f'\varphi \rangle = \langle L', f\varphi \rangle + \langle f'L, \varphi \rangle. \end{aligned}$$

\square

2.4.3. Local Equality. The fact that $L_{2\pi}^1$ is identified with a subset of \mathcal{D}' makes it clear that the “value of a distribution in a point” makes no sense. However, on open sets equality makes sense! Recall that the *support* of a continuous function φ is the closed set $\text{Supp } \varphi = \text{cl}\{t : \varphi(t) \neq 0\}$.

DEFINITION 2.4.4. Two distributions L_1, L_2 on T are called *equal on an open subset* $\Gamma \subset T$ if for every $\varphi \in \mathcal{D}(T)$ with support in Γ , one has $\langle L_1, \varphi \rangle = \langle L_2, \varphi \rangle$.

The *support* of a distribution $L \in \mathcal{D}'$ is the complement of the union of the open sets Γ with $L = 0$ on Γ .

EXAMPLES 2.4.5. The support of δ is $\{0\}$. If L originates from a continuous function then the two notions of support coincide.

REMARK 2.4.6. Multiplication can be localized: If $L \in \mathcal{D}'$ has order k , and on an open $\Gamma \subset T$ is equal to a distribution of order $j < k$ then L can be multiplied with C^j functions, that are C^k in a neighborhood of the complement of Γ . See Exercise 2.7.11. Moreover it turns out that (especially in higher dimensions) a further refinement is possible. This is based on local Fourier analysis of the distribution and takes into account the directions in which the singularities occur. We deal with this topic in a later chapter, but to get a flavour of the problem, consider the distribution L on \mathbb{R}^2 given by

$$\langle L, \varphi \rangle = \frac{\partial \varphi}{\partial x_1}(0, 0)$$

This has order 1, so we can multiply with $f \in C^1$ but a closer analysis gives that we only need to require something like f continuous and differentiable with respect to x_1 .

2.5. Fourier Series of Periodic Distributions

In analogy with $L_{2\pi}^1$ we can define for $L \in \mathcal{D}(T)$ the *Fourier coefficients*

$$\hat{L}(n) = \frac{1}{2\pi} \langle L, e^{-int} \rangle.$$

For example, as we know already $\hat{\delta}(n) = 1/2\pi$.

As one expects, we put

$$S_N[L](t) = \sum_{-N}^N \hat{L}(n) e^{int}.$$

For example, $S_N[\delta] = D_N$, the Dirichlet kernel. Let $\varphi \in \mathcal{D}(T)$

$$\langle S_N[\delta], \varphi \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \sum_{n=-N}^N e^{int} \varphi(t) dt = D_N * \varphi(0) \rightarrow \varphi(0), \quad \text{as } N \rightarrow \infty.$$

We see that the Dirichlet kernels tend weakly to δ .

LEMMA 2.5.1. *If $L \in \mathcal{D}'$ and $\varphi \in \mathcal{D}(T)$ then*

$$\langle L, \varphi \rangle = 2\pi \sum_{-\infty}^{\infty} \hat{L}(-n) \hat{\varphi}(n).$$

PROOF. Using Lemma 2.2.1 we see that

$$\langle L, \varphi \rangle = \lim_{N \rightarrow \infty} \langle L, S_N[\varphi] \rangle = \lim_{N \rightarrow \infty} 2\pi \sum_{-N}^N \hat{\varphi}(n) \hat{L}(-n).$$

This proves the Lemma. □

THEOREM 2.5.2. *If $L \in \mathcal{D}'$, then the Fourier series of L tends weak-* to L .*

PROOF. For every $\varphi \in \mathcal{D}(T)$ we find by Lemma 2.5.1, if $N \rightarrow \infty$,

$$\langle S_N[L], \varphi \rangle = 2\pi \sum_{-N}^N \hat{L}(n) \hat{\varphi}(-n) \rightarrow \langle L, \varphi \rangle.$$

□

COROLLARY 2.5.3. *The map $L \rightarrow \hat{L}$ is injective on \mathcal{D}' .*

PROOF. If $\hat{L}(n) = 0$ for all $n \in \mathbb{Z}$, then $L = 0$ by Theorem 2.5.2. □

COROLLARY 2.5.4. *Let $L \in \mathcal{D}'$. If $L' = 0$ on T , then L is a constant.*

PROOF. The Fourier coefficients of L' are $0 = \widehat{L'}(n) = in\hat{L}(n)$. It follows that the Fourier series of L consists only of the constant term. Theorem 2.5.2 gives that L is a constant. □

THEOREM 2.5.5. *The following are equivalent.*

- i. $\sum_{n=-\infty}^{\infty} c_n e^{int}$ is the Fourier series of a periodic distribution.
- ii. There exist constants N, C such that $|c_n| \leq Cn^N$.

PROOF. If $\sum_{n=-\infty}^{\infty} c_n e^{int}$ is the Fourier series of a periodic distribution L then $\sum_{n=-N}^N c_n e^{int}$ tends weakly to L by Theorem 2.5.2. It follows that for every k , if $N > k$

$$\left\langle \sum_{-N}^N c_n e^{int} - L, e^{-ikt} \right\rangle = c_k - \hat{L}(k)$$

tends to 0 if $N \rightarrow \infty$. Hence $c_k = \hat{L}(k)$. Now L has finite order, say N . Then $|c_n| = |\langle L, e^{-int} \rangle| \leq C\tilde{P}_N(e^{-int}) \leq CNn^N$.

In the other direction, if $|c_n| \leq Cn^N$, then $(c_n/n^{N+1})_n \in l^2(\mathbb{Z})$. Parseval gives

$$f(t) = \sum_{-\infty}^{\infty} \frac{c_n}{n^{N+1}} e^{int} \in L^2(T).$$

It follows that $f^{(N+1)}$ is a distribution of at most order $N+1$. It has Fourier coefficients $i^{N+1}c_n$ for $n \neq 0$. Dividing by i^{N+1} and adding c_0 we have found a distribution with the prescribed Fourier series. \square

COROLLARY 2.5.6. *Every periodic distribution is of the form $F^{(N)} + c$ with F continuous (or in L^2).*

PROOF. This is in the proof of the second part of Theorem 2.5.2 for F in L^2 . If F is not continuous we consider

$$F(t) = \sum_{-\infty}^{\infty} \frac{c_n}{n^{N+1}} e^{int} \in C(T),$$

because the series is uniformly convergent and proceed as in the last part of the proof of the Theorem. \square

2.6. Convolution and Multiplication

From [7] we know that if $f, g \in C_{2\pi}$, then $f * g := \int_{-\pi}^{\pi} f(x-y)g(y) \frac{dy}{2\pi}$ and fg are also in $C_{2\pi}$ and have Fourier series given by

$$(2.6.1) \quad (f * g)^{\wedge}(n) = \iint f(x-y)g(y) e^{-in(x-y)} e^{-iny} \frac{dy}{2\pi} \frac{dx}{2\pi} = \hat{f}(n)\hat{g}(n)$$

and

$$(2.6.2) \quad (fg)^{\wedge}(n) = \sum_{-\infty}^{\infty} \hat{f}(n-j)\hat{g}(j) =: \hat{f} * \hat{g}(n).$$

2.6.1. We can use (2.6.1) to define the convolution $L_1 * L_2$ for periodic distributions $L_i \in \mathcal{D}$:

$$(2.6.3) \quad L_1 * L_2 \stackrel{\text{def}}{=} \sum_{-\infty}^{\infty} \hat{L}_1(n)\hat{L}_2(n) e^{inx}.$$

Application of Theorem 2.5.5 gives that there exist $C, k > 0$ such that $|\hat{L}_1(n)\hat{L}_2(n)| < C|n|^k$, ($n \neq 0$), and another application of Theorem 2.5.5 shows that the series represents a distribution.

Formula (2.6.2) shows once more why it is difficult to multiply distributions: The Fourier coefficients $(fg)^{\wedge}(n)$ have to be finite. Suppose f is a distribution of order k . Then $\hat{f}(n)$ behaves like n^k . For (2.6.2) to converge, the $\hat{g}(n)$ have to be something like n^{-k} , i.e. g is fairly smooth.

2.7. Exercises

2.7.1. Prove that, as $n \rightarrow \infty$, the sequence $\sin(nt)/n^3$ converges to 0 uniformly, but it does not converge in $\mathcal{D}(T)$.

2.7.2. Prove that (2.2.4) defines a metric and complete the proof of Lemma 2.2.2.

2.7.3. Recall the Arzelà-Ascoli theorem: If $\{f_\alpha\}_\alpha$ is an equicontinuous family of pointwise bounded continuous functions on a separable compact metric space, then $\{f_\alpha\}_\alpha$ has a uniformly convergent subsequence. Equicontinuous means

$$\forall \varepsilon \exists \delta \text{ such that } |x-y| < \delta \implies |f_\alpha(x) - f_\alpha(y)| < \varepsilon,$$

independent of α .

Prove the following. If $\{f_\alpha\}_\alpha$ is bounded in the C^1 -norm on T , that is, the norm $\|\cdot\|_\infty + \|\cdot'\|_\infty$ on T , then $\{f_\alpha\}_\alpha$ is equicontinuous and pointwise bounded. (Apply the mean value theorem.)

2.7.4. Prove the continuity in example 2.3.1 by showing that $\varphi_n \rightarrow \varphi$ in $\mathcal{D}(T)$ implies $\langle L_u, \varphi_n \rangle \rightarrow \langle L_u, \varphi \rangle$.

2.7.5. Let $L \in \mathcal{D}'$ be of order k and let $f \in C^l(T)$. For which k and l can you give a definition of fL by using (2.6.1) ?

2.7.6. Compute the distribution sum of

$$(2.7.1) \quad \sum_{n=-\infty}^{\infty} ne^{inx},$$

$$\sum_{n=1}^{\infty} \sin nx.$$

2.7.7. Compute the distributional derivative of

$$(2.7.2) \quad f(x) = \begin{cases} x + \pi & \text{if } -\pi < x < 0, \\ x - \pi & \text{if } 0 < x < \pi. \end{cases}$$

2.7.8. Let $\varphi \in \mathcal{D}(T)$, $L \in \mathcal{D}$. Put $\varphi^x(y) = \varphi(x - y)$. Show that

$$L * \varphi(x) \stackrel{\text{def}}{=} \langle L, \varphi^x \rangle$$

is well-defined and coincides with definition (2.6.3). Prove that $L * \varphi$ is a smooth function.

2.7.9. Let $L \in \mathcal{D}'$ have support $\{0\}$.

- (1) Use Corollary 2.5.6 to express L as the j 'th derivative of a continuous function f on $(-\pi, \pi)$.
- (2) Show that there are polynomials P_1, P_2 of degree at most $j - 1$ such that $f = P_1$ if $x > 0$ and $f(x) = P_2(x)$ if $x < 0$.
- (3) Conclude that we may take $f = PU$ with P a polynomial of degree $j - 1$ and U the characteristic function of $(0, \pi)$.
- (4) Prove that L is a finite linear combination of derivatives of δ .

2.7.10. For $L \in \mathcal{D}$, let L_h be defined by $L_h \varphi := L\varphi(\cdot - h)$. Compute the distributional limit

$$\lim_{h \rightarrow 0} \frac{L_h - L}{h}.$$

2.7.11. Suppose that $L \in \mathcal{D}'$ has order k , and on an open $\Gamma \subset T$ is equal to a distribution \tilde{L} of order $j < k$. Prove that L can be multiplied with any C^j function F , that is C^k in a neighborhood of the complement of Γ . [First prove the result for $L = 0$ on Γ . Then write $L = (L - \tilde{L}) + \tilde{L}$.]

CHAPTER 3

Distributions on \mathbb{R}^n

As before, distributions on \mathbb{R}^n will be continuous linear functionals on test spaces of smooth functions. However, because \mathbb{R}^n is not compact, there is some freedom in the choice of the test space, which leads to different classes of distributions. Also the topology on the test space is more complicated. Therefore we will base our treatment on the semi-norm approach, (2.3.1) of the previous chapter. In particular this makes it clear that advanced knowledge of topological vector spaces is not necessary to work fruitfully with distributions.

We will mention some useful facts about smooth functions in Section 3.1. The proofs will be rather sketchy. Details may be found in [5]. In what follows U will be open in \mathbb{R}^n and K will be compact in \mathbb{R}^n . Variables in \mathbb{R}^n will generally be written as $x = (x_1, \dots, x_n)$, and $dx = dx_1 dx_2 \dots dx_n$ will denote the corresponding Lebesgue measure on \mathbb{R}^n . Also $B(x, r)$ will be the open ball with radius r centered at x . We write $C_0^k(X) \subset C^k(X)$ for the class of k -times differentiable functions that are supported on compact subsets of X ; k may of course be ∞ . The space $C_0^\infty(U)$ will be our *test space* and following Schwartz we denote it by $\mathcal{D}(U)$ or \mathcal{D} if no confusion is possible. As usual the elements of $\mathcal{D}(U)$ are called *test functions*. The reader is warned that this notation is **not** the same as in [7]. Differential operators will be denoted using multi-index notation: with $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$, we put $D^\alpha = D_1^{\alpha_1} \dots D_n^{\alpha_n}$, it is a partial derivative of order $|\alpha| = \alpha_1 + \dots + \alpha_n$; also $\alpha! := \alpha_1! \alpha_2! \dots \alpha_n!$.

3.1. Smooth Functions

LEMMA 3.1.1. *Let $p \in U$. There exists a non-negative test function φ on \mathbb{R}^n with $\varphi(p) > 0$ and $\text{Supp } \varphi$ is a compact subset of U .*

PROOF. Without loss of generality, $p = 0 \in \text{cl } B(0, 1) \subset U$. Let

$$(3.1.1) \quad f(t) = \begin{cases} e^{-1/t} & \text{if } t > 0, \\ 0 & \text{elsewhere.} \end{cases}$$

The function f is smooth on \mathbb{R} . Now take $\varphi(x) = f(1 - \|x\|^2)$ on \mathbb{R}^n , this function is smooth and its support is $\text{cl } B(0, 1)$ □

LEMMA 3.1.2. *There exists a smooth, compactly supported approximate identity on \mathbb{R}^n .*

PROOF. Let φ be as in Lemma 3.1.1 with $p = 0$. Multiplying with a positive constant if necessary, we can assume that $\int_U \varphi(x) dx = 1$. Let

$$\varphi_\varepsilon(x) \stackrel{\text{def}}{=} \frac{1}{\varepsilon^n} \varphi\left(\frac{x}{\varepsilon}\right).$$

Then $\varphi_\varepsilon \geq 0$, $\int \varphi_\varepsilon = 1$ and $\varphi_\varepsilon \rightarrow 0$ uniformly on $\|x\| \geq \delta > 0$, hence $\{\varphi_\varepsilon\}$ is an approximate identity consisting of compactly supported functions. □

THEOREM 3.1.3. *Let $f \in C_0^k(U)$ and let $\{\varphi_\varepsilon\}$ be an approximate identity consisting of compactly supported smooth functions. Then $f * \varphi_\varepsilon$ is C^∞ and for $0 \leq |\alpha| \leq k$ the convolutions $D^\alpha f * \varphi_\varepsilon$ will tend to $D^\alpha f$ uniformly on U if $\varepsilon \rightarrow 0$.*

PROOF. Writing $f * \varphi_\varepsilon(x) = \int f(y) \varphi_\varepsilon(x - y) dy$, an application of Lebesgue's dominated convergence theorem allows us to differentiate under the integral sign infinitely often. On

the other hand we may write $f * \varphi_\varepsilon(x) = \int f(x-y)\varphi_\varepsilon(y)dy$ which yields that for $|\alpha| \leq k$, $D^\alpha(f * \varphi_\varepsilon) = (D^\alpha f) * \varphi_\varepsilon$. Hence it suffices to prove that $f * \varphi_\varepsilon \rightarrow f$ uniformly if $f \in C_0(U)$. Indeed,

$$|f * \varphi_\varepsilon - f(x)| = \left| \int (f(x-y) - f(x))\varphi_\varepsilon(y) dy \right| \leq \delta \int \varphi_\varepsilon \leq \delta,$$

if ε is so small that $|f(x) - f(x-y)| < \delta$ for $y \in \text{Supp } \varphi_\varepsilon$. \square

LEMMA 3.1.4. *Let K be compact in \mathbb{R}^n , $\varepsilon > 0$ and let $K_\varepsilon = \{y : d(y, K) < \varepsilon\}$. There exists $\psi_\varepsilon \in C_0^\infty(K_\varepsilon)$ such that $\psi_\varepsilon = 1$ on K . Moreover the derivatives satisfy*

$$(3.1.2) \quad |D^\alpha \psi_\varepsilon(x)| \leq C_\alpha \varepsilon^{-|\alpha|},$$

for certain constants C_α depending on α but not on ε .

PROOF. Let $\{\varphi_\varepsilon\}$ be the compactly supported approximate identity constructed in Lemma 3.1.2. Then φ_ε is supported in $B(0, \varepsilon)$ and its derivatives satisfy $D^\alpha(\varphi_\varepsilon)(x) = \frac{1}{\varepsilon^{|\alpha|}}(D^\alpha \varphi)\left(\frac{x}{\varepsilon}\right)$. Let χ be the characteristic function of $K_{\varepsilon/2}$ and set

$$\psi_\varepsilon = \chi * \varphi_{\varepsilon/2}.$$

As in the proof of Theorem 3.1.3 we obtain that ψ_ε is smooth. It is obviously supported in K_ε and its derivatives are $\psi * D^\alpha(\varphi_{\varepsilon/2})$ which is bounded by a $\varepsilon^{-|\alpha|}C_\alpha$ by our previous estimate. \square

We have used the straightforward but noteworthy fact that for L^1 functions f and g , $\text{Supp}(f * g) \subset \text{Supp } f + \text{Supp } g$, where for $A, B \subset \mathbb{R}^n$ we write $A+B = \{a+b : a \in A, b \in B\}$.

LEMMA 3.1.5. *Every open U can be written as $U = \cup_1^\infty K_j$ with K_j compact and K_j a subset of the interior of K_{j+1} . In fact we can make the convenient choice*

$$(3.1.3) \quad K_j = \{x \in U : \|x\| \leq j \text{ and } d(x, \partial U) \geq 2^{-j}\}.$$

PROOF. Immediate. \square

DEFINITION 3.1.6. A *partition of unity* of an open set U subordinate to an open cover $\mathcal{U} = \{U_\tau\}_\tau$ of U , (i.e., $U = \cup_\tau U_\tau$), is a countable set of nonnegative smooth functions $\{\varphi_j, j = 1, \dots\}$ on \mathbb{R}^n with the following properties:

1. $\forall j \exists \tau_j$ such that $\text{Supp } \varphi_j \subset U_{\tau_j}$.
2. If K is a compact subset of U , then only finitely many of the φ_j are non-zero on K .
3. On U we have $\sum_j \varphi_j \equiv 1$.

Notice that the sum in 3. is well-defined because on every compact K we only have to take into account the finitely many φ_j that are positive on K . We keep the notation of the definition.

THEOREM 3.1.7. *For every open cover \mathcal{U} of U , there exists a partition of unity subordinate to \mathcal{U} .*

PROOF. First remark that every compact $K \subset U$ is contained in one of the K_j of Lemma 3.1.5. Thus we only have to check 2. of Definition 3.1.6 for these K_j .

The important observation is as follows. If $x \notin K_j$, then $d(x, \partial U) < 2^{-j}$ or $\|x\| > j$. In the first case, $d(x, K_{j-1}) > 2^{-j}$, in the latter case $d(x, K_{j-1}) > 1$. The ball $B(x, r)$ with $r = \frac{1}{2} \min\{1, d(x, \partial U)\}$ will have empty intersection with K_{j-1} .

We now define for $x \in U_\tau$ the ball $B_\tau(x) = B(x, t)$ with $t = \frac{1}{2} \min\{1, d(x, \partial U_\tau)\}$. By Lemma 3.1.1 we can find non-negative $\varphi_{\tau, x} \in \mathcal{D}(B_\tau(x))$ with $\varphi_{\tau, x}(x) = 1$. Keeping in mind that \mathcal{U} covers U we introduce for every $x \in U$ (at least one) open neighborhood $D_\tau(x) = \{y \in B_\tau(x) : \varphi_{\tau, x}(y) > 1/2\}$.

We will select a good covering of U . First $K_1 \subset \cup_{x \in K_1} D_\tau(x)$. We can find a finite subcover labeled $D_{1,1}, \dots, D_{1,j_1}$. Next, $K'_2 = K_2 \setminus (\cup_{j=1}^{j_1} D_{1,j})$ is compact; again $K'_2 \subset \cup_{x \in K'_2} D_\tau(x)$. We can find a finite subcover labeled $D_{2,1}, \dots, D_{2,j_2}$. The $D_{1,j}$ and $D_{2,j}$ cover

K_2 . Continuing in this way, covering each time $K'_i = K_i \setminus \cup_{l < i} D_{l,j}$ with finitely many $D_{i,j}$, and relabeling, we obtain a countable covering D_1, D_2, \dots of U . Call the corresponding $B_\tau(x)$'s: B_1, B_2, \dots and the corresponding $\varphi_{\tau,x}$'s: $\varphi_1, \varphi_2, \dots$

Now every φ_i has support in some $B_i = B_\tau(x) \subset U_\tau$, therefore it satisfies 1. For condition 2 notice that there are only finitely many B_i of the form $B(x)$ with x in K_j . By our first observation, if $x \notin K_j$ then $B(x) \cap K_{j-1} = \emptyset$. Thus for every j , K_{j-1} is covered by finitely many B_i and condition 2 is satisfied.

To finish the proof we have to modify φ_i a little. Observe that

$$\varphi = \sum_1^\infty \varphi_i$$

is a well-defined smooth function on U , because on compact sets only finitely many terms are non-zero, and since every $y \in U$ is in some D_i , the sum is at least $1/2$. Now introduce

$$\psi_i = \frac{\varphi_i}{\varphi}.$$

Clearly $\psi_i \in \mathcal{D}(U_\tau)$ and the ψ_i satisfy 1, 2, and 3 and form a partition of unity. \square

COROLLARY 3.1.8. *For every $\psi \in C_0^\infty(U)$ and every cover \mathcal{U} of U there exists a finite subcollection U_i of \mathcal{U} such that we can write $\psi = \sum_{i=1}^N \psi_i$ with $\psi_i \in \mathcal{D}(U_i)$.*

PROOF. Let K be the support of ψ . Let φ_j , $j = 1, \dots, N$ be the finitely many φ 's that are non zero on K . Then $\psi \varphi_j \in \mathcal{D}(\mathbb{R}^n)$ has compact support in $U_{\tau_j} \cap K$. The U_{τ_j} constitute a finite set $\{U'_i \mid i = 1, \dots, N\}$. We have

$$\psi = \psi 1 = \sum_j \psi \varphi_j, \quad \text{on } K.$$

Assembling the φ_j which have support in the same U_i , i.e., setting $\psi_i = \sum_{\tau_j=U_i} \varphi_{\tau_j}$, we are done. \square

REMARK 3.1.9. It is possible to put a topology on the test space $\mathcal{D}(U)$ that makes it a complete locally convex topological vector space. This is not so easy, because we have tried to keep $\mathcal{D}(U)$ as small as possible in order to have as many continuous linear functionals as possible on it. However, occasionally we don't need so many distributions and deal with two "easier" spaces of test functions.

We introduce seminorms $Q_{K,\alpha}$ on $C^\infty(U)$ as follows

$$(3.1.4) \quad Q_{K,\alpha}(\varphi) = \sup_{x \in K} |D^\alpha \varphi|,$$

where K runs over the compacts in U and α over all multiindices. $C^\infty(U)$ equipped with the topology induced by $Q_{K,\alpha}$ is denoted by $\mathcal{E}(U)$ or simply \mathcal{E} .

3.1.1. The Schwartz Class. In [7] the class \mathcal{S} of Laurent Schwartz was introduced on \mathbb{R} . For \mathbb{R}^n the definition is similar

$$(3.1.5) \quad \mathcal{S} = \mathcal{S}(\mathbb{R}^n) = \{f \in C^\infty(\mathbb{R}^n) : \text{for every } \alpha \in \mathbb{N}^n \text{ and } k \in \mathbb{N}, \|x\|^k |D^\alpha(f)| \text{ is bounded}\}.$$

\mathcal{S} is topologized by the seminorms $P_{k,\alpha}$ defined by

$$(3.1.6) \quad P_{k,\alpha}(f) = \sup_{\mathbb{R}^n} \|x\|^k |D^\alpha(f)|, \quad a \in \mathbb{N}^n, k \in \mathbb{N}.$$

We say that $f \in C^\infty$ is rapidly decreasing if for every k and every α , $|D^\alpha(f(x))| \|x\|^k$ tends to 0 if $\|x\| \rightarrow \infty$. Replacing k by $k+k'$ in the definition we see that in fact \mathcal{S} is the collection of rapidly decreasing functions.

PROPOSITION 3.1.10. *$\mathcal{D}(\mathbb{R}^n)$ is a dense subspace of \mathcal{S} .*

PROOF. Let $f \in \mathcal{S}$. We have to show that given $k, l \in \mathbb{N}$, $\varepsilon > 0$, there exists $\varphi \in \mathcal{D}$ with $P_{k,\alpha}(f - \varphi) < \varepsilon$ for $|\alpha| \leq l$. Let $\psi \in \mathcal{D}$, $0 \leq \psi \leq 1$, with $\psi = 1$ on $B(0, 1)$. Put $\psi_r(x) = \psi(x/r)$, then $\psi_r = 1$ on $B(0, r)$ and $|D^\alpha(\psi_r)| \rightarrow 0$ for $1 \leq |\alpha| \leq k$ if $r \rightarrow \infty$. Put $\varphi_r = f\psi_r$. Also $(1 - \psi_r)f \rightarrow 0$ if $r \rightarrow \infty$. We have

$$D^\alpha(f - \varphi_r) = \sum_{p+q=\alpha} c_{p,q} D^p(f) D^q(1 - \psi_r),$$

for suitable (fixed) constant $c_{p,q}$ and multiindices p and q . Using that $D^p(f) = O(1/\|x\|^k)$ and that for $|q| > 1$ all $D^q(1 - \psi_r)$ can be made arbitrary small by choosing r large enough, while in case $|q| = 0$ this can be done on compact sets, we can bound $D^\alpha(f - \varphi_r)$ by $\varepsilon\|x\|^{-k}$. The proposition follows. \square

PROPOSITION 3.1.11. \mathcal{D} is dense in \mathcal{E} .

We leave the proof as an exercise.

3.2. Distributions

DEFINITION 3.2.1. A *distribution* L on U is a linear functional on $\mathcal{D}(U)$ such that for every K compact in U there exist C, N such that

$$(3.2.1) \quad |L\varphi| = |\langle L, \varphi \rangle| \leq C \max_{|\alpha| < N} \|D^\alpha(\varphi)\|, \quad \text{for } \varphi \in \mathcal{D}(U) \text{ supported in } K.$$

The set of distributions on U is denoted by $\mathcal{D}'(U)$. If N can be chosen independent of K , the smallest N such that (3.2.1) holds for all K is called the *order* of L .

We give $\mathcal{D}'(U)$ the weak-* topology, that is,

$$L \mapsto L\varphi$$

is a continuous map for every $\varphi \in \mathcal{D}(U)$. Equivalently $L_j \rightarrow L$ in $\mathcal{D}'(U)$ if and only if for every test function φ we have

$$(3.2.2) \quad L_j\varphi \rightarrow L\varphi.$$

LEMMA 3.2.2. Let φ and $(\varphi_j)_j$ be in $\mathcal{D}(U)$ and have support in a compact set $K \subset U$. Suppose that for every α , $D^\alpha\varphi_j \rightarrow D^\alpha\varphi$ uniformly on K . Then for every $L \in \mathcal{D}'(U)$ we have $L\varphi_j \rightarrow L\varphi$.

PROOF. $L(\varphi - \varphi_j) \rightarrow 0$ in view of (3.2.1). \square

Differentiation, local equality and support of distributions are defined just like in Chapter 2. E.g.,

$$\langle D^\alpha L, \varphi \rangle \stackrel{\text{def}}{=} (-1)^{|\alpha|} \langle L, D^\alpha \varphi \rangle, \quad L \in \mathcal{D}', \varphi \in \mathcal{D}.$$

EXAMPLES 3.2.3. Let δ_j denote point mass at $j \in \mathbb{R}$. The following distributional limit exists $L = \sum_{j=1}^{\infty} \delta_j^{(j)}$. It has infinite order.

Let $u_t(x) = t^N e^{ixt}$, $x \in \mathbb{R}$, $N \in \mathbb{N}$, then as distributions the limit for $t \rightarrow \infty$ exists: using integration by parts,

$$\langle u_t, \varphi \rangle = \int_{\mathbb{R}} t^N e^{itx} \varphi(x) dx = i^{N+1} t^{-1} \int_{\mathbb{R}} e^{itx} \varphi^{(N+1)}(x) dx \rightarrow 0, \quad \text{as } t \rightarrow \infty, \quad \varphi \in \mathcal{D}.$$

In addition we define

DEFINITION 3.2.4. The *singular support* of $L \in \mathcal{D}'$ is the complement (in U) of the union of the open sets where L is equal to a C^∞ -function.

THEOREM 3.2.5. Let $V \subset \mathbb{R}^m$ be open. Let $L \in \mathcal{D}'(U)$. If $\varphi \in C^\infty(U \times V)$ is such that $\varphi(\cdot, v)$ has support inside a compact $K \subset U$ for all $v \in V$, then $L\varphi(\cdot, v) \in C^\infty(V)$. If $\varphi \in \mathcal{D}(U \times V)$, then $L\varphi(\cdot, v) \in \mathcal{D}(V)$.

PROOF. We write down the second order Taylor development of φ with respect to the second variable at $v \in V$:

$$\varphi(u, v + h) = \varphi(u, v) + \nabla_v \varphi(u, v)h + \psi(u, v, h),$$

where, by compactness of K for every α

$$(3.2.3) \quad \max_{u \in K} |D_u^\alpha \psi(u, v, h)| < C_\alpha \|h\|^2.$$

We conclude

$$(3.2.4) \quad \frac{L\varphi(\cdot, v + h) - L\varphi(\cdot, v) - L(\nabla_v \varphi(\cdot, v))h}{\|h\|} = \frac{1}{\|h\|} |L\psi(\cdot, v, h)|.$$

Using that $|L\psi(\cdot, v, h)|$ is bounded by a constant times the maximum over finitely many α 's of left hand sides of (3.2.3), we see that (3.2.4) tends to 0 with $\|h\|$. We conclude that $L\varphi$ is C^1 , its partial derivatives (in V) are given by $LD_j \varphi(\cdot, v)$. By what we have proved, they are C^1 ! Continuing in this way we obtain the first statement of the theorem. For the second statement, if $\varphi \in \mathcal{D}(U \times V)$ then $\varphi(\cdot, v)$ will vanish identically for v outside a suitable compact set in V , hence so will $L\varphi(\cdot, v)$. \square

THEOREM 3.2.6. *Let $V \subset \mathbb{R}^m$ and let $\varphi = \varphi(u, v) \in \mathcal{D}(U \times V)$. If $L \in \mathcal{D}'(U)$ then $L \int \varphi(\cdot, v) dv = \int L\varphi(\cdot, v) dv$.*

PROOF. We use Riemann sums. For every α we have

$$\int D_u^\alpha \varphi(u, v) dv = \lim_{h \rightarrow 0} \sum_{t \in \mathbb{Z}^m} D_u^\alpha \varphi(u, ht) h^m,$$

uniformly on a compact set in U , because of compactness. Thus Lemma 3.2.2 implies that

$$L \int \varphi(\cdot, v) dv = L \left(\lim_{h \rightarrow 0} \sum_{t \in \mathbb{Z}^m} \varphi(\cdot, ht) h^m \right) = \lim_{h \rightarrow 0} \sum_{t \in \mathbb{Z}^m} L(\varphi(\cdot, ht)) h^m = \int L\varphi(\cdot, v) dv.$$

\square

THEOREM 3.2.7. *Every distribution $L \in \mathcal{D}'(U)$ of order k can be extended to a linear form on $C_0^k(U)$, in such a way that (3.2.1) holds for all $\varphi \in C_0^k$.*

PROOF. Let $f \in C_0^k(U)$. By Theorem 3.1.3 there exists a sequence of functions $\varphi_j \in \mathcal{D}(U)$ that approximate f in the sense that the partial derivatives of $f - \varphi_j$ of order $\leq k$ tend to 0 uniformly. It follows that for $|\alpha| \leq k$, $D^\alpha(\varphi_j)$ form a Cauchy sequence in the uniform norm on U . The same is then true for $L\varphi_j$, because of the estimate (3.2.1). By usual arguments this implies that

$$Lf = \lim_{j \rightarrow \infty} L\varphi_j.$$

is well-defined, independent of the choice of φ_j , and that the estimate (3.2.1) also holds for f . \square

COROLLARY 3.2.8. *A distribution of order 0 is a complex Borel measure.*

PROOF. Recalling that complex Borel measures on U are precisely the linear forms that satisfy (3.2.1) for all $\varphi \in C_0(U)$, this is immediate from the theorem. \square

A distribution $L \in \mathcal{D}'(U)$ is called *positive* if $L\varphi > 0$ for every non-negative test function φ .

THEOREM 3.2.9. *Positive distributions are measures.*

PROOF. Let K be compact in U and let $0 \leq \psi \leq 1$ be a test function that equals 1 on K . If $\varphi \in C_0^\infty(K)$ is real valued, then

$$\|\varphi\|_\infty \psi \pm \varphi \geq 0.$$

Applying L to these relations, it follows that for real valued test functions supported in K

$$(3.2.5) \quad |L\varphi| \leq \|\varphi\|_\infty L\psi.$$

For complex valued testfunctions we write $\varphi = \varphi_1 + i\varphi_2$. Applying (3.2.5) we obtain in this case

$$|L\varphi| \leq 2\|\varphi\|_\infty L\psi, \quad \varphi \in \mathcal{D}(K).$$

This means that L has order 0 and thus it is a measure. \square

3.3. Distributions with Compact Support

Recall that we introduced \mathcal{E} as the space of smooth functions on U equipped with the topology determined by (3.1.4). Its dual space is denoted by $\mathcal{E}'(U)$. As in Chapters 1 and 2 we see that $L \in \mathcal{E}'$ if and only if there exist a compact set K in U , and constants C, N with

$$(3.3.1) \quad |L\varphi| \leq C \max_{x \in K, |\alpha| \leq N} |D^\alpha(\varphi)|, \quad (\varphi \in \mathcal{E}).$$

THEOREM 3.3.1. *Let $L \in \mathcal{D}'(U)$ be compactly supported with $\text{Supp } L = K$. There exists precisely one linear functional \tilde{L} on $C^\infty(U)$ with the following properties:*

- i. $\tilde{L} = L$ on $\mathcal{D}(U)$;
- ii. $\tilde{L}\varphi = 0$ for all φ in C^∞ with support in the complement of K .

Moreover, $\tilde{L} \in \mathcal{E}'(U)$.

PROOF. Let $0 \leq u \leq 1 \in \mathcal{D}$ be identically 1 on a neighborhood of K and let $S = \text{Supp } u$. We define

$$\tilde{L}\varphi = L(u\varphi), \quad \varphi \in \mathcal{E}.$$

Observe that for $\varphi \in \mathcal{D}$ we have

$$L\varphi = L(u\varphi) + L((1-u)\varphi) = \tilde{L}(\varphi),$$

because $(1-u)\varphi \in \mathcal{D}$ has its support in the complement of K , thus is annihilated by L . Next if $\varphi \in \mathcal{E}$ and $\text{Supp } \varphi$ is in the complement of K , then $u\varphi$ has compact support in the complement of K , so $L(u\varphi) = 0$.

Suppose L_1 is another linear functional satisfying i. and ii. Let $\varphi \in \mathcal{E}$. We write $\varphi = u\varphi + (1-u)\varphi$. The first term in the sum is in \mathcal{D} , the second has support in the complement of K , hence is annihilated by L_1 . Then

$$L_1\varphi = L_1(u\varphi + (1-u)\varphi) = L_1(u\varphi) + L_1((1-u)\varphi) = L(u\varphi) + 0 = \tilde{L}\varphi.$$

Finally observe that by (3.2.1)

$$(3.3.2) \quad |\tilde{L}\varphi| = |L(u\varphi)| \leq C \max_{x \in S, |\alpha| \leq N} |D^\alpha(u\varphi)|.$$

This implies by Leibnitz rule that

$$|\tilde{L}\varphi| \leq C' \max_{x \in S, |\alpha| \leq N} |D^\alpha(\varphi)|,$$

where the norms of the derivatives of the (fixed) function u are absorbed in C' . In other words, $\tilde{L} \in \mathcal{E}'$. \square

The proof of Theorem 3.3.1 shows that for a distribution supported on a compact set K , we can take the maximum in (3.3.1) over an arbitrarily small compact neighborhood of K . Can't we just take K ? The following example shows that the answer is in general No. It is known that the supports for which the answer is Yes, are finite unions of compact connected sets such that any two points x, y in the same component can be joined by a rectifiable arc of length $C|x - y|$, cf. [5], Th. 2.3.10.

EXAMPLE 3.3.2. Consider the sequence X with elements $x_n = 1/n^3$ in \mathbb{R} . Let $K = X \cup \{0\}$ be the closure of X . Define $L \in \mathcal{D}'(\mathbb{R})$ by

$$(3.3.3) \quad \langle L, \varphi \rangle = \sum_{n=1}^{\infty} n(\varphi(x_n) - \varphi(0)).$$

Observe that L is compactly supported (on K). Let us show that it has order 1. By the mean value theorem $\varphi(x_n) - \varphi(0) = 1/n^3 \varphi'(\xi_n)$ for certain $\xi_n \in (0, x_n)$. We substitute this in (3.3.3) and find

$$(3.3.4) \quad |\langle L, \varphi \rangle| \leq C \max_{0 \leq x \leq 1} |\varphi'(x)|.$$

However, we can not replace this maximum by a maximum over K , even if we were willing to increase the order of L . To see this, let $\varphi_m \in \mathcal{D}(\mathbb{R})$ satisfy

$$(3.3.5) \quad \varphi_m(x) = \begin{cases} 0 & \text{for } x \leq x_{m+1}, \\ 1 & \text{for } x_m \leq x \leq 2. \end{cases}$$

We have $|\varphi_m| \leq 1$ on K and all derivatives of φ vanish on K . If (3.3.4) would hold with the max taken over K , then it would follow that $\langle L, \varphi_m \rangle$ were uniformly bounded. However, $\langle L, \varphi_m \rangle = m$, a contradiction!

REMARK 3.3.3. This example easily generalizes to more general compacts having infinitely many connected components.

THEOREM 3.3.4. Let $L \in \mathcal{E}'$ be of order k and supported on a compact K . Suppose $\varphi \in \mathcal{E}$ has the property that $D^\alpha \varphi \equiv 0$ on K for all $|\alpha| \leq k$. Then $L\varphi = 0$.

PROOF. Let $u_\varepsilon \in \mathcal{D}$ equal 1 on K and 0 outside K_ε as constructed in Lemma 3.1.4. Then in view of Theorem 3.3.1 we have $L\varphi = Lu_\varepsilon\varphi$ for all ε . There exist C and k such that

$$|Lu_\varepsilon\varphi| \leq C \max_{x \in K_\varepsilon, |\alpha| \leq k} |D^\alpha(u_\varepsilon\varphi)|.$$

Using (3.1.2) we see that

$$|D^\alpha(U_\varepsilon\varphi)| \leq C'_\alpha \sum_{|\beta| \leq |\alpha|} \varepsilon^{|\beta| - |\alpha|} |D^\beta\varphi|.$$

Therefore, it suffices to show that for all $|\beta| \leq k$

$$(3.3.6) \quad \varepsilon^{|\beta| - k} D^\beta\varphi \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

If $|\beta| = k$, then this follows from continuity of $D^\beta\varphi$ and $D^\beta\varphi \equiv 0$ on K . For the general case we apply Taylor's formula to $D^\beta\varphi$ on an interval $[x, y]$, $x \in K$, $y \in K_\varepsilon$:

$$(3.3.7) \quad |D^\beta(\varphi(t(y-x) + x))| \leq C|t|^j \|y-x\|^j \sum_{|\beta'|=j} |D^{\beta'+\beta}\varphi(\xi)|.$$

where $j = k - |\beta|$, $\xi \in [x, y]$. We take $t = 1$ and use again that for $|\alpha| = k$, we have $D^\alpha\varphi(x) = o(1)$ on K_ε for $\varepsilon \rightarrow 0$. Hence (3.3.7) becomes

$$|D^\beta(\varphi(y))| = \varepsilon^j o(1) \quad \text{on } K_\varepsilon \quad \text{for } \varepsilon \rightarrow 0.$$

Inserting this in (3.3.6) we are done. \square

COROLLARY 3.3.5. *If $L \in \mathcal{D}'$ has order N and $\text{Supp } L = \{0\}$, then*

$$L\varphi = \sum_{|\alpha| \leq N} c_\alpha D^\alpha \varphi(0).$$

That is, L is a linear combination of derivatives of δ .

PROOF. Let $\varphi \in \mathcal{E}$. Write $\varphi(x) = \sum_{|\alpha| \leq N} \frac{D^\alpha \varphi(0)}{\alpha!} x^\alpha + \psi(x)$, where the derivatives of ψ up to order N at 0 vanish. By Theorem 3.3.4 $L\psi = 0$, hence

$$L\varphi = \sum_{|\alpha| \leq N} L(x^\alpha) \frac{D^\alpha \varphi(0)}{\alpha!}.$$

□

3.4. Convolutions and Product Spaces

It is convenient to have some notation for translated and reflected functions. Let φ be a function on \mathbb{R}^n .

- Translation: $(\tau_x \varphi)(y) = \varphi_x(y) = \varphi(y - x)$;
- Reflection: $\check{\varphi}(y) = \varphi(-y)$.

Recalling the definition of convolution we see that

$$(3.4.1) \quad f * g(x) = \int_{\mathbb{R}^n} f(y)g(x - y) dy = \int f(y)\tau_x \check{g}(y) dy, \quad f, g \in L^1(\mathbb{R}^n)$$

It is now natural to make the following

DEFINITION 3.4.1. For $L \in \mathcal{D}'$, $\varphi \in \mathcal{D}$ the *convolution*

$$L * \varphi(x) \stackrel{\text{def}}{=} \langle L, \tau_x \check{\varphi} \rangle (= \langle L, \varphi(x - \cdot) \rangle).$$

By Theorem 3.2.5, $L * \varphi$ is a smooth function. In view of Theorem 3.3.1, if L has compact support, then $L * \varphi$ is defined for $\varphi \in \mathcal{E}$. Note that $L * \varphi(0) = L\check{\varphi}$.

THEOREM 3.4.2. *If $L \in \mathcal{D}'$, and $\varphi, \psi \in \mathcal{D}$ then*

$$(L * \varphi) * \psi = L * (\varphi * \psi).$$

PROOF. We use Theorem 3.2.6.

$$(3.4.2) \quad \begin{aligned} (L * \varphi) * \psi(t) &= \int (L\tau_x \check{\varphi})\psi(t - x) dx = L \int \varphi(x - \cdot)\psi(t - x) dx \\ &= L \int \varphi(x)\psi(t - \cdot - x) dx = L(\varphi * \psi(t - \cdot)) = L * (\varphi * \psi)(t). \end{aligned}$$

□

Convolution with a distribution is a translation invariant operator from \mathcal{D} to \mathcal{E} . Indeed,

$$\tau_x(L * \varphi)(t) = \tau_x(L(\tau_t \check{\varphi})) = L(\tau_{t-x} \check{\varphi}) = L(\tau_t(\tau_x \varphi)^\sim) = L * (\tau_x \varphi)(t).$$

It also has some continuity properties, in the sense that, of course, $\varphi \mapsto L * \varphi(t)$ is for every t a distribution. The converse is also true:

THEOREM 3.4.3. *Suppose that $T : C_0^\infty \rightarrow C(\mathbb{R}^n)$ is a translation invariant operator with the property that $\varphi \mapsto T\varphi(0)$ is an element of \mathcal{D}' . Then there exists a unique $L \in \mathcal{D}'$ such that $T\varphi = L * \varphi$. In particular, T maps C_0^∞ into C^∞ .*

PROOF. Let $L\varphi \stackrel{\text{def}}{=} T\check{\varphi}(0)$. Then L is a distribution. We compute

$$L * \varphi(x) = L * (\tau_{-x} \varphi)(0) = L(\tau_{-x} \varphi)^\sim = T(\tau_{-x} \varphi)(0) = \tau_{-x} T\varphi(0) = T\varphi(x).$$

Thus we have found L . If $\tilde{L} * \varphi = L * \varphi$ for all $\varphi \in \mathcal{D}$ then for test functions φ

$$\langle \tilde{L}, \varphi \rangle = \tilde{L} * \varphi(0) = L * \varphi(0) = \langle L, \varphi \rangle,$$

and $\tilde{L} = L$ by the very definition of distribution. □

Now let $L_1 \in \mathcal{D}'$, $L_2 \in \mathcal{E}'$. Then for $\varphi \in \mathcal{D}$ we find that

$$L_2 * (L_1 * \varphi) \quad \text{and} \quad L_1 * (L_2 * \varphi)$$

are well defined – for the second expression observe that $L_2 * \varphi$ has compact support. Moreover, both operators are translation invariant. Therefore by Theorem 3.4.3 we can make the following

DEFINITION 3.4.4. $L_1 * L_2$ is the unique distribution L with the property that $L * \varphi = L_1 * (L_2 * \varphi)$. One similarly defines $L_2 * L_1$.

One can show that $L_1 * L_2 = L_2 * L_1$, cf. exercise 3.6.7.

There is another way of looking at convolutions. We start with (3.4.1) and wish to view that as a distribution, thus let it act on $\varphi \in \mathcal{D}$.

$$\langle f * g, \varphi \rangle = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} f(y)g(x-y) dy \varphi(x) dx = \iint_{\mathbb{R}^n \times \mathbb{R}^n} f(y)g(t)\varphi(y+t) dy dt.$$

This explains commutativity of convolution as a direct consequence of commutativity in \mathbb{R}^n . To make it work we have to make sense of $u(x)v(y)$ for distributions $u \in \mathcal{D}'(\mathbb{R}^n)$, $v \in \mathcal{D}'(\mathbb{R}^m)$.

Let $X_j \subset \mathbb{R}^{n_j}$, $j = 1, 2$. For $f_j \in C(X_j)$ we define the *tensor product* as follows

$$f_1 \otimes f_2(x_1, x_2) = f_1(x_1)f_2(x_2).$$

What we need is an analogous tensor product of distributions. It should be defined on $X_1 \times X_2$ and it is clear how it must act on tensor products of test functions of the form $\varphi_1 \otimes \varphi_2$.

THEOREM 3.4.5. Suppose that X_j are open sets in \mathbb{R}^{n_j} ($j = 1, 2$) and that $u_j \in \mathcal{D}'(X_j)$. There exists a unique distribution $u \in \mathcal{D}'(X_1 \times X_2)$ which has the property that

$$u(\varphi_1 \otimes \varphi_2) = u_1(\varphi_1)u_2(\varphi_2) \quad \varphi_j \in \mathcal{D}(X_j).$$

Moreover,

$$(3.4.3) \quad u(\varphi) = u_{1x}[u_{2y}\varphi(x, y)] = u_{2y}[u_{1x}\varphi(x, y)], \quad \varphi \in \mathcal{D}(X_1 \times X_2).$$

We write $u = u_1 \otimes u_2$, the *tensor product* of u_1 and u_2 .

Notice that the expressions in (3.4.3) are well-defined in view of Theorem 3.2.5 L_x is a sloppy notation, indicating that the distribution L acts on the test functions viewed as a function of x , while the other variables are viewed as parameters.

PROOF. Uniqueness follows once we can show that for $L \in \mathcal{D}'(X_1 \times X_2)$ the assumption $L(\varphi_1 \otimes \varphi_2) = 0$ for all $\varphi_j \in \mathcal{D}(X_j)$, implies that $L = 0$. Thus let $\varphi \in \mathcal{D}(X_1 \otimes X_2)$. Choose $\psi_i \in \mathcal{D}(X_i)$ positive with integral equal to 1. Then $\psi_\varepsilon(x, y) = \varepsilon^{-n_1-n_2}\psi_1(\varepsilon^{-n_1}x)\psi_2(\varepsilon^{-n_2}y)$ is a tensor product, as well as an approximate identity on $\mathbb{R}^{n_1} \times \mathbb{R}^{n_2}$. Therefore

$$L(\varphi * \psi_\varepsilon) = L * (\varphi * \psi_\varepsilon)(0) = ((L * \psi_\varepsilon) * \varphi)(0) = 0,$$

by the assumption. Letting $\varepsilon \rightarrow 0$ uniqueness follows.

Next we show that

$$(3.4.4) \quad \varphi \mapsto u_{1x}[u_{2y}(\varphi(x, y))], \quad \varphi \in \mathcal{D}(X_1 \times X_2)$$

is in $\mathcal{D}'(X_1 \times X_2)$. We estimate, using Theorem 3.2.5,

$$(3.4.5) \quad \begin{aligned} |u_{1x}[u_{2y}(\varphi(x, y))]| &\leq C_1 \sup_{\substack{|\alpha| \leq m_1 \\ x \in K_1}} |D_x^\alpha(u_{2y}(\varphi(x, y)))| = C_1 \sup_{\substack{|\alpha| \leq m_1 \\ x \in K_1}} |(u_{2y}(D_x^\alpha \varphi(x, y)))| \\ &\leq C_1 C_2 \sup_{\substack{|\alpha| \leq m_1 \\ x \in K_1}} \sup_{\substack{|\beta| \leq m_2 \\ y \in K_2}} |D_y^\beta D_x^\alpha \varphi(x, y)|. \end{aligned}$$

Hence u is in $\mathcal{D}'(X_1 \times X_2)$. Since u has the required behavior on tensor products of smooth functions, the unicity part completes the proof. \square

REMARK 3.4.6. We mention without proof the important *kernel theorem*:

THEOREM 3.4.7 (Laurent Schwartz). *There is a one one correspondence between distributions K on $X_1 \times X_2$ and continuous linear maps T from $\mathcal{D}(X_1)$ to $\mathcal{D}'(X_2)$ given by*

$$K \in \mathcal{D}'(X_1 \times X_2) \quad \leftrightarrow \quad T_K, \quad \langle T_K(\varphi_1), \varphi_2 \rangle = \langle K, \varphi_1 \otimes \varphi_2 \rangle.$$

Here continuity means $T\varphi_j \rightarrow 0$ in \mathcal{D}' if $\varphi_j \rightarrow 0$ in \mathcal{D} . A proof can be found in [5].

3.5. Tempered distributions

The dual space \mathcal{S}' of \mathcal{S} is called the space of *tempered distributions* (or temperate). It turns out that this is the “best” space to do Fourier analysis on. In this section we only give a few properties of \mathcal{S}' . In view of (3.1.6) a linear functional L on \mathcal{S} is in \mathcal{S}' if and only if there exist $C > 0$, $k, N \in \mathbb{N}$, such that

$$|L\varphi| \leq C \sup_{x \in \mathbb{R}^n} \|x\|^k |D^\alpha \varphi(x)|, \quad \text{for all } |\alpha| \leq N.$$

EXAMPLES 3.5.1. The distribution associated to a locally integrable function f with the property that

$$\int (1 + \|x\|^2)^{-N} |f(x)| dx \quad \text{is bounded,}$$

for N sufficiently large, is tempered. This explains the word tempered: moderate growth at infinity is allowed. If P is a polynomial and $L \in \mathcal{S}'$ then $PL \in \mathcal{S}'$; also $D^\alpha L \in \mathcal{S}'$; compactly supported distributions are in \mathcal{S}' .

3.6. Exercises

3.6.1. Determine the distributional limit of any approximate identity φ_ε .

3.6.2. Suppose that $\varphi \in \mathcal{D}(\mathbb{R}^n)$ and $\varphi(0) = 0$. Prove that there exist $\varphi_j \in \mathcal{D}$ such that $\varphi(x) = \sum_{j=1}^n x_j \varphi_j(x)$. Hint: At 0 write

$$\varphi(x) = \int_0^{x_1} D_1 \varphi(t, x_2, \dots, x_n) dt + \varphi(0, x_2, \dots, x_n).$$

Continue with respect to the other variables and change the interval of integration to $[0, 1]$. get everything in \mathcal{D} by writing far away from 0

$$\varphi = \sum \frac{x_i^2 \varphi(x)}{\|x\|^2},$$

and patch things together with a partition of unity.

3.6.3. Let $u_t(x) = t^N e^{itx}$, for $x \geq 0$ and $u_t(x) = 0$ elsewhere. Determine the distributional limit $\lim_{t \rightarrow \infty} u_t$.

3.6.4. Let $u_t(x) = t^{1/k} e^{itx^k}$, $x \in \mathbb{R}$ and k an integer > 1 . Determine the distributional limit $\lim_{t \rightarrow \infty} u_t$. Hint: Using integration by parts, write

$$\langle u_t, \varphi \rangle = - \int F(x) t^{1/k} \varphi'(x) dx = (F(\infty) - F(-\infty)) \varphi(0),$$

with $F(x) = \int_0^x e^{it^k} dt$. Next apply Cauchy's theorem to compute

$$\begin{aligned} \lim_{x \rightarrow \infty} F(x) &= e^{\pi i/2k} \int_0^\infty e^{-t^k} dt, \\ \lim_{x \rightarrow -\infty} F(x) &= -e^{(-1)^k \pi i/2k} \int_0^\infty e^{-t^k} dt. \end{aligned}$$

Compare with the case $k = 1$ which is in Example 3.3.2.

3.6.5. Show that distributions are locally determined, i.e., if $L \in \mathcal{D}'$ has the property that for every x there is a neighborhood B_x such that $L = 0$ on B_x , then $L \equiv 0$.

3.6.6. (continued) Suppose that U_τ is an open cover of U and that on every U_τ there is given $L_\tau \in \mathcal{D}(U_\tau)$. Show that there exists $L \in \mathcal{D}(U)$ with $L = L_\tau$ on U_τ if and only if $L_\tau = L_\sigma$ on $U_\tau \cap U_\sigma$ whenever this intersection is non empty. Show also that L is unique.

3.6.7. If u_1, u_2 are distributions one of them with compact support, then $u_1 * u_2 = u_2 * u_1$. Prove this by showing first that for any $\varphi_1, \varphi_2 \in \mathcal{D}$ the equality

$$u_1 * u_2 * \varphi_1 * \varphi_2 = u_2 * u_1 * \varphi_1 * \varphi_2$$

holds and that this implies the equality that you are looking for.

3.6.8. (continued) Show that $\text{Supp } u_1 + \text{Supp } u_2$ contains $\text{Supp } u_1 * u_2$.

3.6.9. (continued) Prove that $(u_1 * u_2)(\varphi) = (u_1 \otimes u_2)(\varphi(x + y))$.

3.6.10. Suppose $L \in \mathcal{D}'$ has the property that $D_j L = 0$ for $j = 1, \dots, n$. Prove that L is a constant.

3.6.11. Let $u, v \in \mathcal{D}'$. Show that $u * \delta = u$. Show that $u * D^\alpha \delta = D^\alpha u$. Finally show that $D^\alpha(u * v) = (D^\alpha u) * v = u * (D^\alpha v)$.

3.6.12. Show that for every compactly supported distribution u there exists a sequence $(\varphi_j)_j$ in \mathcal{D} such that $\varphi_j \rightarrow u$ in distribution sense. (Use smart convolutions).

3.6.13. (continued) Show that every $u \in \mathcal{D}'$ is a distributional limit of distributions with compact support. Conclusion?

3.6.14. (Riesz) Let $a \in \mathbb{C}$ and let

$$(3.6.1) \quad \chi_a(x) = \begin{cases} x^a, & \text{if } x \geq 0; \\ 0, & \text{if } x \leq 0 \end{cases}$$

i. Show that for $\mathbf{Re } a > -1$ the function χ_a is in $L^1_{loc}(\mathbb{R})$, i.e. integrable over compact sets in \mathbb{R} , and satisfies $(a + 1)\chi_a = \frac{d}{dx}\chi_{a+1}$.

ii. Let I_a denote the distribution associated to χ_a . Show that it satisfies

$$(3.6.2) \quad I_a \varphi' = -a I_{a-1}(\varphi), \quad \text{if } \mathbf{Re } a > 0, \quad \varphi \in \mathcal{D}(\mathbb{R}).$$

iii. Observe that for fixed φ the function $I_a(\varphi)$ is analytic in a for $\mathbf{Re } a > -1$ and use (3.6.2) to extend I_a analytically to $\mathbb{C} \setminus \{-1, -2, \dots\}$. Show that

$$(3.6.3) \quad \langle I_a, x\varphi \rangle = \langle I_{a+1}, \varphi \rangle \quad \text{if } a \in \mathbb{C} \setminus \{-1, -2, \dots\}.$$

iv. Try to compute the residues of I_a in the negative integers. (These must be distributions, of course).

v. Try to compute the distribution I_a minus its principal part in a , where a is a negative integer. Show that (6.2) holds for this distribution too.

3.6.15. Show that the distribution on \mathbb{R}^n induced by $f(x) = e^{|x|^2}$ is not tempered. Construct a function g on \mathbb{R} such that g induces a tempered distribution and

$$\limsup_{t \rightarrow \infty} g(t)e^{-t^2} = 1.$$

3.6.16. Show that the compactly supported distributions are dense in \mathcal{S}' .

3.7. Final remarks

Distributions were introduced in the 1940's by Laurent Schwartz. His book [10] is a classic. Already around 1930 Sobolev came very close to the concept while the physicist Dirac worked with distribution ideas informally. An interesting book about the history of distribution theory is Lutzens [8]; Dutch readers may be happily surprised by the relatively large Dutch role in it. Another classical reference is [3].

We have followed Hörmander [5] quite closely. Because this book is aimed at partial differential operators, it selects the parts of the theory that are really important for other subjects in a way that can be appreciated beyond abstract functional analysis. It is an excellent book anyway and highly recommended.

The Fourier Transform

We begin this chapter by reviewing the *Fourier transform* on $\mathcal{S}(\mathbb{R}^n)$. The case $n = 1$ is treated in [7][part 2] and also in [12]; this part of the theory is virtually the same if $n > 1$. Proofs that are analogous to the one variable case that is treated in [7, 12] will often be omitted.

In the later sections we will deal with Fourier transforms of tempered distributions and prove the *Paley-Wiener Theorem*, which characterizes Fourier transforms of distributions with compact support. We will also study Fourier transform in relation with other spaces.

Notation in this and later chapters: For $x, y \in \mathbb{R}^n$ (or \mathbb{C}^n) we will write the bilinear form

$$x \cdot y \stackrel{\text{def}}{=} \sum_{i=1}^n x_i y_i.$$

Also, from now on

$$D_j \stackrel{\text{def}}{=} -i \frac{\partial}{\partial x_j}.$$

This absorbing of i in the operator is convenient in connection with Fourier transform.

4.1. Fourier Transform on \mathcal{S}

The *Fourier transform* $\mathcal{F}f$ of $f \in L^1(\mathbb{R}^n)$ is defined as

$$(\mathcal{F}f)(\xi) = \hat{f}(\xi) = \int_{\mathbb{R}^n} f(x) e^{-ix \cdot \xi} dx.$$

The function \hat{f} is bounded and continuous. In fact, the *Riemann-Lebesgue Lemma* states that $\hat{f} \rightarrow 0$ as $|\xi| \rightarrow \infty$.

THEOREM 4.1.1. *Let $f \in L^1(\mathbb{R}^n)$. Then*

$$\lim_{|\xi| \rightarrow \infty} \mathcal{F}f(\xi) = 0.$$

An instructive proof is indicated in exercise 4.5.1. If f and \hat{f} are both in L^1 , then the *inversion formula* holds:

$$f(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \hat{f}(\xi) e^{ix \cdot \xi} d\xi.$$

In particular, \mathcal{F} is a continuous injective map from \mathcal{S} onto \mathcal{S} and the inversion formula may be written as

$$\mathcal{F}^2 \varphi(-x) = (2\pi)^n \varphi(x).$$

See exercise 4.5.2 for a proof different from [7]. Other properties are

(4.1.1)

$$\mathcal{F}(D_j \varphi)(\xi) = \xi_j (\mathcal{F}\varphi)(\xi), \quad \mathcal{F}(x_j \varphi)(\xi) = (-D_j \mathcal{F}\varphi)(\xi), \quad \mathcal{F}(\varphi(\cdot - x))(\xi) = e^{ix \cdot \xi} \mathcal{F}\varphi(\xi)$$

The map \mathcal{F} is extendible to an isometry from $L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$, thus

$$\int_{\mathbb{R}^n} f(t) \bar{g}(t) dt = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \hat{f}(\xi) \bar{\hat{g}}(\xi) d\xi.$$

The following related formulas are valid, first on \mathcal{S} and then continuously extended to L^2 :

$$(4.1.2) \quad \int \hat{f}g = \int f\hat{g}, \quad (g * f)^\wedge = \hat{f}\hat{g}, \quad \widehat{fg} = (2\pi)^{-n} \hat{f} * \hat{g}.$$

Proofs of these formula are indicated in exercise 4.5.3

4.2. Fourier Transform on \mathcal{S}'

We will use the first formula in (4.1.2) and the fact that $\mathcal{F}\mathcal{S} = \mathcal{S}$ to define the Fourier transform of a tempered distribution. In fact if in this first formula $g \in \mathcal{S}$, then $\hat{g} \in \mathcal{S}$ so the righthand side of the formula has a meaning for $f \in \mathcal{S}'$ and can be used to define the lefthand side.

DEFINITION 4.2.1. For $L \in \mathcal{S}'$ we define its *Fourier transform* $\mathcal{F}L = \hat{L} \in \mathcal{S}'$ by

$$\langle \hat{L}, \varphi \rangle = \langle L, \hat{\varphi} \rangle, \quad \varphi \in \mathcal{S}.$$

Fourier transform is continuous on \mathcal{S}' , see exercise 4.5.4. Because (4.1.2) is valid for $f \in L^1(\mathbb{R}^n)$ and $g \in \mathcal{S}$, the Fourier transform of a distribution stemming from an L^1 function f is the same as the distribution associated to \hat{f} . Most formulas for Fourier transform remain valid in the distribution setting. E.g.,

$$\langle \hat{\hat{L}}, \varphi \rangle = \langle \hat{L}, \hat{\varphi} \rangle = \langle L, \hat{\hat{\varphi}} \rangle = (2\pi)^n \langle L, \check{\varphi} \rangle,$$

Therefore, $\hat{\hat{L}} = (2\pi)^{2n} \check{L}$, which is the inversion formula for \mathcal{S}' . In particular \mathcal{F} is a bijection of \mathcal{S}' .

THEOREM 4.2.2. Let $u \in \mathcal{E}'$. Then \hat{u} can be represented by a smooth function (also denoted by \hat{u}) and

$$(4.2.1) \quad \hat{u}(\xi) = u_x(e^{-ix \cdot \xi}), \quad \xi \in \mathbb{R}^n.$$

Moreover, the righthand side of (4.2.1) defines a holomorphic function on \mathbb{C}^n .

PROOF. In view of Theorem 3.3.1 and 3.2.5 the righthand side of (4.2.1) is a smooth function of ξ on \mathbb{C}^n . It satisfies the *Cauchy-Riemann* equations, i.e.

$$\frac{\partial \hat{u}}{\partial \xi_j} = 0, \quad j = 1, \dots, n$$

because $e^{-ix \cdot \xi}$ satisfies these, and differentiation towards the parameter is allowed in view of Theorem 3.2.5.

This proves holomorphy. Now let $\varphi \in \mathcal{S}$. Using Theorem 3.4.5 we find

$$(4.2.2) \quad \begin{aligned} \hat{u}(\varphi) &= u(\hat{\varphi}) = u_x \int e^{-ix \cdot \xi} \varphi(\xi) d\xi \\ &= u_x \otimes \varphi_\xi(e^{ix \cdot \xi}) = \varphi_\xi[u_x(e^{-ix \cdot \xi})] = \int u_x(e^{-ix \cdot \xi}) \varphi(\xi) d\xi. \end{aligned}$$

□

EXAMPLES 4.2.3. The Fourier transform of δ equals

$$\hat{\delta} = \delta_x(e^{-ix\xi}) = 1.$$

Its derivatives have Fourier transform

$$(4.2.3) \quad \widehat{D_j \delta} = D_j \delta(e^{-ix\xi}) = \delta(\xi_j e^{-ix\xi}) = \xi_j.$$

THEOREM 4.2.4 (Convolutions). If $u_1 \in \mathcal{S}'$, $u_2 \in \mathcal{E}'$ then $u_1 * u_2 \in \mathcal{S}'$ and $(u_1 * u_2)^\wedge = \hat{u}_1 \hat{u}_2$.

PROOF. Recall that by the definition of convolution

$$(u_1 * u_2)(\varphi) = u_1 * u_2 * \check{\varphi}(0) = u_1(\check{u}_2 * \varphi), \quad \varphi \in \mathcal{D}.$$

The righthand side extends to a continuous form on \mathcal{S} : if u_2 has order k and $\varphi \in \mathcal{S}$, then using $\check{u}_s * \varphi = u_2(\varphi(x + \cdot))$, we check that

$$\sum_{|\alpha+\beta|\leq j} \sup |x^\beta D^\alpha(\check{u}_2 * \varphi)| \leq \sum_{|\alpha+\beta|\leq j+k} \sup |x^\beta D^\alpha(\varphi)|.$$

Now we compute the Fourier transform. For $\varphi \in \mathcal{S}$

$$\begin{aligned} (4.2.4) \quad (u_1 * u_2)(\hat{\varphi}) &= u_1(\check{u}_2 * \hat{\varphi}) = u_{1x}(\check{u}_{2y}(\hat{\varphi}(x - y))) = u_{1x}\left(\int u_{2y}e^{-i(x+y)t}\varphi(t) dt\right) \\ &= u_{1x}\left(\int \hat{u}_2(t)e^{-i(x)t}\varphi(t) dt\right) = u_{1x}((\hat{u}_2\varphi)\hat{\ }) = \hat{u}_1(\hat{u}_2\varphi) = (\hat{u}_1\hat{u}_2)(\varphi). \end{aligned}$$

□

COROLLARY 4.2.5. For $u \in \mathcal{S}'$ we have

$$\widehat{D}_j u = \xi_j \hat{u}, \quad \widehat{x}_j u = -D_j \hat{u}.$$

PROOF. This follows directly from (4.2.3) and the previous theorem, e.g.,

$$\widehat{D}_j u = \widehat{D}_j \delta * u = \xi_j \hat{u}(\xi).$$

One can also use that the corollary is known for \mathcal{S} a dense subset of \mathcal{S}' in combination with continuity of \mathcal{F} on \mathcal{S}' . □

REMARK 4.2.6 (Computational tricks). If $f \in \mathcal{S}'$ is associated to a locally integrable function, also denoted by f , its Fourier transform can be computed by approximation:

$$(4.2.5) \quad f = \lim_{A \rightarrow \infty} f_A \quad \text{in } \mathcal{S}' \text{ with } f_A(x) = \begin{cases} f(x) & \text{if } |x| \leq A; \\ 0 & \text{if } |x| > A. \end{cases}$$

This follows from Lebesgue's dominated convergence theorem. By continuity of \mathcal{F} we find

$$(4.2.6) \quad \hat{f} = \lim_{A \rightarrow \infty} \hat{f}_A = \lim_{A \rightarrow \infty} \int_{|x| \leq A} f(x) e^{-ix\xi} dx.$$

Similarly, again by Lebesgue's theorem

$$f = \lim_{\varepsilon \downarrow 0} f e^{-\varepsilon|x|} \quad \text{in } \mathcal{S}'$$

and therefore

$$(4.2.7) \quad \hat{f} = \lim_{\varepsilon \downarrow 0} \int_{\mathbb{R}^n} f(x) e^{-\varepsilon|x| - ix\xi} d\xi.$$

Of course, all limits are in \mathcal{S}' and will in general not exist pointwise or in L^1 sense.

EXAMPLES 4.2.7. Applying 4.2.6 and 4.2.7 to the tempered distribution 1 on \mathbb{R} yields

$$\mathcal{F}(1) = \lim_{A \rightarrow \infty} \int_{-A}^A e^{-ix\xi} dx = \lim_{A \rightarrow \infty} 2 \frac{\sin A\xi}{\xi} \quad (= 2\pi\delta).$$

$$\mathcal{F}(1) = \lim_{\varepsilon \downarrow 0} \int_0^\infty e^{-i\xi x - \varepsilon x} dx + \int_{-\infty}^0 e^{-i\xi x + \varepsilon x} dx = \lim_{\varepsilon \downarrow 0} i \left(\frac{1}{\xi - i\varepsilon} - \frac{1}{\xi + i\varepsilon} \right) \quad (= 2\pi\delta).$$

The limits as $\varepsilon \rightarrow 0$ of the fractions between the big brackets are usually denoted by

$$\frac{1}{\xi - i0} \quad \text{respectively} \quad \frac{1}{\xi + i0}.$$

These two are **not** the same! Indeed, with χ denoting characteristic function, we have by the above

$$\mathcal{F}(\chi_{(0,\infty)}) = \frac{i}{\xi - i0}, \quad \mathcal{F}(\chi_{(-\infty,0)}) = \frac{-i}{\xi + i0}.$$

In the same vein we find for the signum function Sign

$$\mathcal{F}(\text{Sign}) = \frac{i}{\xi - i0} - \frac{-i}{\xi + i0} = \lim_{\varepsilon \downarrow 0} i \left(\frac{1}{\xi - i\varepsilon} + \frac{1}{\xi + i\varepsilon} \right) = \lim_{\varepsilon \downarrow 0} \frac{2i\xi}{\xi^2 + \varepsilon^2} = 2iP.V. \frac{1}{\xi}.$$

The *Airy function* $Ai(x)$ is classically defined as the inverse Fourier transform of $e^{i\xi^3/3}$. Thus $Ai(x)$ is a tempered distribution. In fact, it is in C^∞ . We have in analogy with 4.2.7

$$(4.2.8) \quad Ai(x) = \frac{1}{2\pi} \lim_{\varepsilon \downarrow 0} \int e^{i(\xi+i\varepsilon)^3/3+i(\xi+i\varepsilon)x} d\xi \quad \text{in } \mathcal{S}'.$$

The integrals on the righthand side are (for a fixed positive ε) convergent and yield smooth functions of x . They can even be extended analytically to $x \in \mathbb{C}$, because the leading term in the exponent is $-\xi^2\varepsilon$, independent of $x \in \mathbb{C}$. On the other hand they are independent of ε in view of Cauchy's theorem.

In \mathbb{R}^n we will consider Gaussians. Let A be a positive definite symmetric matrix. Then clearly $e^{-Ax \cdot x}$ is in \mathcal{S} . It is called a *Gaussian*. We compute its Fourier transform. We can write $A = R^2$ with R again positive symmetric.

$$(4.2.9) \quad \int_{\mathbb{R}^n} e^{-Ax \cdot x - i\xi \cdot x} dx = \int e^{-Rx \cdot Rx - iR^{-1}\xi \cdot Rx} dx.$$

Now we change coordinates, $Rx = y$ and (4.2.9) equals

$$\int e^{-y \cdot y - iR^{-1}\xi \cdot y} dR^{-1}y = (\det A)^{-1/2} e^{-\frac{1}{4}A^{-1}\xi \cdot \xi} \int e^{-(y + \frac{i}{2}R^{-1}\xi) \cdot (y + \frac{i}{2}R^{-1}\xi)} dy.$$

The last integral is, by an application of *Cauchy's theorem* (compare the proof of the Paley-Wiener-Schwartz theorem), equal to

$$\int e^{-\|y\|^2} dy = \prod_{j=1}^n \int e^{-y_j^2} dy_j = (2\pi)^{n/2}.$$

The conclusion is that

$$\mathcal{F}(e^{-Ax \cdot x}) = (2\pi)^{n/2} (\det A)^{-1/2} e^{-\frac{1}{4}A^{-1}\xi \cdot \xi}.$$

4.3. The Paley-Wiener-Schwartz Theorem

Let K be a compact subset of \mathbb{R}^n . The *supporting function* of K is

$$H(x) = \sup_{y \in K} \{x \cdot y\}.$$

PROPOSITION 4.3.1. *Let $\mathcal{H}_x = \{y : y \cdot x \leq H(x)\}$. Then the convex hull of K is given by*

$$\text{ch } K = \bigcap_{x \in \mathbb{R}^n} \mathcal{H}_x.$$

PROOF. Recall that the convex hull of $K \subset \mathbb{R}^n$ is the intersection of all convex sets that contain K . Observe that $y \in K$ implies that for every x , $x \cdot y \leq H(x)$, therefore $K \subset \mathcal{H}_x$ for every x . Using the well-known corollary of the Hahn-Banach theorem, cf. [W], that for every closed convex G and $y \notin G$ there exists a hyperplane separating y from G , we find that $\text{ch } K$ equals the intersection of all half-spaces containing K . In particular, if $x_0 \notin \text{ch } K$ then we can find x such that $x \cdot x_0 > H(x)$, thus $x_0 \notin \mathcal{H}_x$. \square

THEOREM 4.3.2 (Paley-Wiener-Schwartz). *Let H be the supporting function of a compact set K in \mathbb{R}^n .*

1. *The following are equivalent:*

i. *The distribution $u \in \mathcal{E}'$ has order N and support with convex hull $K \subset \mathbb{R}^n$*

ii. There exists a constant C such that the Fourier transform \hat{u} is an entire function on \mathbb{C}^n that satisfies the inequality

$$(4.3.1) \quad |\hat{u}(z)| \leq C(1 + |z|)^N e^{H(\mathbf{Im} z)}, \quad (z \in \mathbb{C}^n).$$

2. The following are equivalent:

i. The function $u \in \mathcal{D}$ has compact support with convex hull $K \subset \mathbb{R}^n$.

ii. For every N there exists a constant C_N such that the Fourier transform \hat{u} is an entire function on \mathbb{C}^n that satisfies the inequality

$$(4.3.2) \quad |\hat{u}(z)| \leq C_N(1 + |z|)^{-N} e^{H(\mathbf{Im} z)}, \quad (z \in \mathbb{C}^n).$$

PROOF. We begin with 2. If $u \in \mathcal{D}$ then for every α , $D^\alpha u \in \mathcal{D}$ and

$$|z^\alpha \hat{u}(z)| = |\widehat{D^\alpha u}(z)| \leq \|D^\alpha u(z)\|_1 \max_{\xi \in \text{Supp } u} |e^{-iz \cdot \xi}| \leq C_\alpha e^{H(\mathbf{Im} z)}.$$

This shows that (4.3.2) is necessary. Next, suppose that (4.3.2) holds for some entire function U . Then U restricted to \mathbb{R}^n is the Fourier transform of some function u which is in C^∞ because of (4.3.2). We have

$$(4.3.3) \quad u(\xi) = (2\pi)^{-n} \int e^{i\xi \cdot x} U(x) dx.$$

Consider the integral in (4.3.3) as a repeated integral, say first integrating with respect to x_1 , the other variables being fixed. The integrand is a holomorphic function in $x_1 + iy_1$, and on the strip $\{z_1 : |y_1| \leq A_1\}$ it is bounded by a constant times $(1 + |z_1|)^{-N}$. Thus we may apply Cauchy's theorem and replace integration over \mathbb{R} by integration over $\mathbb{R} + iA_1$, for any A_1 in \mathbb{R} . We can also do this for the other variables, and obtain that for every $A = A_1, \dots, A_n$

$$(4.3.4) \quad u(\xi) = (2\pi)^{-n} \int e^{i\xi \cdot (x+iA)} U(x+iA) dx.$$

Using (4.3.2) with N sufficiently large, we infer from (4.3.4) that for every A

$$|u(\xi)| \leq (2\pi)^{-n} e^{-\xi \cdot A + H(A)}.$$

By considering tA , $t > 0$, instead of A , we infer that $u(\xi) \neq 0$ implies that $\xi \cdot A \leq H(A)$, that is, $\xi \in \mathcal{H}_A$ for all A . However, by Proposition 4.3.1, this implies that $\xi \in K$.

Next we turn to 1. and show that (4.3.1) is necessary. Suppose that u has order N and compact support in the convex set $K \subset \mathbb{R}^n$. Let $K_\delta = \{x + y : x \in K, |y| < \delta\}$ be the δ -neighborhood of K . Let $\chi = \chi_\delta \in \mathcal{D}(K_\delta)$ equal 1 on $K_{\delta/2}$. By Lemma 3.1.4 $|D^\alpha \chi_\delta| = O(\delta^{-|\alpha|})$. In view of Theorem 3.1.3 we estimate

$$(4.3.5) \quad \begin{aligned} |\hat{u}(z)| &= |u(e^{-iz \cdot \cdot})| = |u(\chi_\delta e^{-iz \cdot \cdot})| \leq \sum_{|\alpha| \leq n} \max_{x \in K_\delta} C_\alpha |D_\xi^\alpha (\chi_\delta(\xi) e^{\xi \cdot z})| \\ &\leq C e^{H(y) + \delta|y|} \sum_{|\alpha| \leq N} \delta^{-|\alpha|} (1 + |z|)^{N - |\alpha|}. \end{aligned}$$

For a fixed $|z|$, choose $\delta = \varepsilon(1 + |z|)^{-1}$ and necessity of (4.3.1) follows. Finally we show that (4.3.1) is sufficient. If U satisfies (4.3.1) then U restricted to \mathbb{R}^n is the Fourier transform of a distribution u , cf. Example 3.5.1. Let (φ_ε) be an approximate identity, which has $\text{Supp } \varphi_\varepsilon \subset B(0, \varepsilon)$. By 2. of the Theorem $(u * \varphi_\varepsilon)^\wedge = \hat{u} \hat{\varphi}_\varepsilon$, which in view of (4.3.2) and (4.3.1) satisfies

$$|(u * \varphi_\varepsilon)^\wedge(z)| \leq C_N (1 + |z|)^{-N} e^{H(y) + \varepsilon|y|}.$$

From 2. we conclude that $u * \varphi_\varepsilon$ has support in K_ε . Letting $\varepsilon \rightarrow 0$ we obtain that $\text{Supp } u \subset K$. \square

4.4. Fourier Transform and other Function Spaces

4.4.1. Sobolev Spaces. For positive integers s Sobolev introduced what is now known as the *Sobolev space* $H_{(s)}$. It is the completion of the space of compactly supported smooth functions on \mathbb{R}^n in the norm

$$(4.4.1) \quad \|f\|_{(s)} = \sum_{|\alpha| \leq s} \|D^\alpha f\|_2.$$

Obviously, this norm is induced by an inner product, namely, the sum of the L^2 inner products of the derivatives, and thus $H_{(s)}$ becomes a *Hilbert space*. In fact, a similar definition with L^p norms is possible, leading to spaces $H_{(s)}^p$. We will not pursue this here.

Using Fourier transform we can give an alternative definition which is meaningful for all $s \in \mathbb{R}$, thereby in some sense defining derivatives of fractional order.

DEFINITION 4.4.1. For $s \in \mathbb{R}$, let L_s^2 denote the *weighted L^2 space* $L^2\left(\mathbb{R}^n, (1 + |\xi|^2)^s \frac{d\xi}{(2\pi)^n}\right)$. Its norm is denoted $\|\cdot\|_s$. The *Sobolev space* $H_{(s)}(\mathbb{R}^n)$ consists of all $u \in \mathcal{S}'$ such that $\hat{u} \in L_s^2$. We put a norm on $H_{(s)}$ by

$$(4.4.2) \quad \|u\|_{(s)}'^2 = \|\hat{u}\|_s^2 = \int |\hat{u}(\xi)|^2 (1 + |\xi|^2)^s \frac{d\xi}{(2\pi)^n}.$$

This turns $H_{(s)}$ into a Hilbert space and Fourier transform is obviously an isometry from $H_{(s)}$ onto L_s^2 .

PROPOSITION 4.4.2. *In case s is a positive integer the norms defined by (4.4.1) and (4.4.2) are equivalent.*

PROOF. There exist positive integers C_α such that $(1 + |\xi|^2)^s = (1 + \xi_1^2 + \cdots + \xi_n^2)^s = \sum_{|\alpha| \leq s} c_\alpha \xi^{2\alpha}$, so that an application of Plancherel's theorem yields

$$\|u\|_{(s)}'^2 = \int |\hat{u}|^2 (1 + |\xi|^2)^s = \sum_{|\alpha| \leq s} c_\alpha \xi^\alpha \hat{u} \overline{\xi^\alpha \hat{u}} = \sum_{|\alpha| \leq s} \int D^\alpha u \overline{D^\alpha u} = \sum_{|\alpha| \leq s} c_\alpha \|D^\alpha u\|_2^2.$$

□

In fact, we don't need all monomials. For $0 < s < \infty$ there exists constants C_s with

$$(1 + |\xi|^2)^s \leq C_s (1 + |\xi|^{2s}) \leq 2C_s (1 + |\xi|^2)^s.$$

Therefore the norm (4.4.2) is equivalent to

$$(4.4.3) \quad (\|u\|_{(s)}'')^2 = \int |\hat{u}|^2 (1 + |\xi|^{2s}) \frac{d\xi}{(2\pi)^n}.$$

This may be used to derive *Sobolev's Lemma*

LEMMA 4.4.3. *Suppose that $f \in H_{(s)}(\mathbb{R}^n)$. If $p \geq 0$ and $s > p + n/2$, then $f \in C^p(\mathbb{R}^n)$.*

For a proof, see exercise 4.5.7.

Next we give a description of $H_{(s)}$ without Fourier transform.

PROPOSITION 4.4.4. *Let $0 \leq s \leq 1$. There exists a constant A_s such that*

$$(4.4.4) \quad (\|u\|_{(s)}')^2 = \int |u|^2 + A_s \iint |u(x) - u(y)|^2 |x - y|^{-n-2s} dx dy.$$

PROOF. We rewrite the double integral in (4.4.4) and apply the Plancherel theorem with respect to the variable y , using that $(u(x+y) - u(y))^\wedge = (e^{ix \cdot \xi} - 1)\hat{u}(\xi)$:

$$\iint |u(x+y) - u(y)|^2 |x|^{-n-2s} dx dy = \iint |e^{ix \cdot \xi} - 1|^2 |\hat{u}(\xi)|^2 |x|^{-n-2s} dx d\xi.$$

To prove the statement it is sufficient to show that

$$|\xi|^{-2s} \int |e^{ix \cdot \xi} - 1|^2 |x|^{-n-2s} dx = A(s),$$

where $A(s)$ only depends on s . This is indeed the case. The integral is invariant under rotation of ξ , (change the coordinates accordingly) so it is a function of $|\xi|$. Next replace ξ by $t\xi$ to see that it is constant.

Thus for $0 < s < 1$ $H_{(s)}$ is the space of L^2 functions such that (4.4.4) is finite. More generally for k a positive integer, $H_{(k+s)}$ is the space of functions f with $D^\alpha f \in L^2$ for $|\alpha| \leq k$ and (4.4.4) is finite for all derivatives of f of order k .

For negative s we can use duality to describe $H_{(s)}$ as follows. By the *Riesz representation theorem* the dual of L_s^2 can be identified with L_s^2 via the inner product of L_s^2 . However, we can make another identification with L_{-s}^2 . An $f \in L_{-s}^2$ acts on $g \in L_s^2$ via

$$(4.4.5) \quad \langle f, g \rangle = \int f(t)g(t)dt.$$

Observe that fg is integrable because $\int |fg| \leq \|f\|_{-s} \|g\|_s$, which also serves to show that f induces a continuous functional. Moreover, every $l \in L_s^{2*}$ may be represented, according to Riesz, by some $\tilde{f} \in L_s^2$. Then $f(x) = \tilde{f}(x)(1+|x|^2)^{-s}$ represents l in the sense of (4.4.5). In view of Fourier transform being an isometry from $H_{(s)}^2$ onto L_s^2 , we conclude that for $s < 0$, $H_{(s)}^2$ can be identified with the dual of $H_{(-s)}^2$.

4.4.2. L^p -spaces. Because

$$(4.4.6) \quad \begin{aligned} \mathcal{F} : L^2(\mathbb{R}^n) &\rightarrow L^2(\mathbb{R}^n), \\ \mathcal{F} : L^1(\mathbb{R}^n) &\rightarrow L^\infty(\mathbb{R}^n), \end{aligned}$$

it is reasonable to expect that for $1 < q < 2$ Fourier transform maps L^q to some space between L^2 and L^∞ . That this is indeed the case follows from the *Riesz-Thorin Convexity Theorem*.

THEOREM 4.4.5 (Riesz Thorin). *Let $1 \leq p_j, q_j \leq \infty$ and let T be a continuous linear mapping from $L^{p_1} \cap L^{p_2}$ to $L^{q_1} \cap L^{q_2}$, such that*

$$(4.4.7) \quad \|Tf\|_{q_j} \leq M_j \|f\|_{p_j}.$$

Put $\frac{1}{p} = t\frac{1}{p_1} + (1-t)\frac{1}{p_2}$ and $\frac{1}{q} = t\frac{1}{q_1} + (1-t)\frac{1}{q_2}$. Then

$$(4.4.8) \quad \|Tf\|_q \leq M_1^t M_2^{1-t} \|f\|_p.$$

COROLLARY 4.4.6 (Hausdorff-Young). *If $f \in L^p$, $1 \leq p \leq 2$, then $\hat{f} \in L^{p'}$, $1/p + 1/p' = 1$ and*

$$\|\hat{f}\|_{p'} \leq (2\pi)^{n/p'} \|f\|_p.$$

PROOF OF THE COROLLARY. \mathcal{F} satisfies the conditions of Riesz-Thorin, with $p_1 = 1$, $q_1 = \infty$, $M_1 = 1$ and $p_2 = q_2 = 2$, $M_2 = (2\pi)^{n/2}$. So the intermediate estimates follow from (4.4.8). \square

PROOF OF THE RIESZ-THORIN THEOREM. Recall from functional analysis that $(L^q)^* = L^{q'}$ with $1/q + 1/q' = 1$. Moreover, for any h that is locally integrable we can estimate by duality

$$\|h\|_q = \sup_{\|g\|_{q'}=1} |\langle h, g \rangle| = \sup_{\|g\|_{q'}=1} \left| \int h(t)g(t) dt \right|.$$

The sup is finite if and only if $h \in L^q$. We can now rewrite (4.4.7) as

$$(4.4.9) \quad |\langle Tf, g \rangle| \leq M_j \|f\|_{p_j} \|g\|_{q'_j} \quad \forall g \in L^{q'_j}.$$

To prove the theorem it suffices to prove

$$(4.4.10) \quad |\langle Tf, g \rangle| \leq M_1^t M_2^{1-t} \|f\|_p \|g\|_{q'},$$

which is the dual version of (4.4.8), for f of the form $f_0 F^{1/p}$ tested against g of the form $g_0 G^{1/q'}$, where f_0, g_0 are in absolute value ≤ 1 and, because step functions are dense in L^r if $r < \infty$, F, G are non negative step functions with integral equal to 1. Now consider, for fixed such f_0, F, g_0, G

$$\Phi(z) = \langle T(f_0 F^{z/p_1 + (1-z)/p_2}), g_0 G^{z/q_1 + (1-z)/q_2} \rangle M_1^{-z} M_2^{z-1}.$$

This function is continuous on $0 \leq \mathbf{Re} z \leq 1$ and analytic on the interior of this set (The reader may want to check this first for F, G being multiples of characteristic functions). Moreover it is bounded on the strip and has boundary values ≤ 1 on $\mathbf{Re} z \in \{0, 1\}$. The *Phragmén-Lindelöf Theorem* states that Φ is in absolute value ≤ 1 and the Theorem is proven. \square

4.5. Exercises

4.5.1. Prove the Riemann-Lebesgue Lemma 4.1.1 along the following lines

- i Write $f = f_1 + f_2$ with $f_1 = \sum_{j=1}^n c_j \chi_j$ and $\|f_2\|_1 \varepsilon$ small. Here, χ_j are indicator functions of blocks of the form $\{a_i < x_i < b_i, i = 1, \dots, n\}$. (This is essentially a fact from integration theory.)
- ii Prove the Riemann-Lebesgue Lemma for χ_j by direct integration.
- iii Conclude from i and ii that $\lim_{|\xi| \rightarrow \infty} |\mathcal{F}f(\xi)| < \varepsilon$ and complete the proof of the theorem.

4.5.2. Let $T\varphi = \mathcal{F}\mathcal{F}\varphi$. Prove that $(TD_j - D_jT)\varphi = 0$ and $T(x_j\varphi) - x_jT(\varphi) = 0$, for $\varphi \in \mathcal{S}$. Use exercise 3.6.2 and these relations to show that $(T\varphi)(x) = c(x)\varphi(x)$ for an appropriate function $c(x)$. Use the relations again to show that $c(x)$ is a constant c . Take $\varphi = e^{-|x|^2}$ to determine c and derive the inversion formula for \mathcal{S} .

4.5.3. Prove the formulas (4.1.2). Start with applying Fubini's theorem to the double integral of $f(x)g(y)e^{ix \cdot y}$. For the Fourier transform of a convolution, split the exponent in the exponential function in accordance with the convolution.

4.5.4. Prove that \mathcal{F} is a continuous map from \mathcal{S} to \mathcal{S} .

4.5.5. Suppose that f is an entire function on \mathbb{C}^n that satisfies the growth condition (4.3.1) for some N and that its restriction to \mathbb{R}^n is in $L^2(\mathbb{R}^n)$. Show that \hat{f} is a compactly supported L^2 -function.

4.5.6. Suppose that f is an entire function on \mathbb{C}^n that satisfies for every $\varepsilon > 0$ a growth condition

$$|f(z)| \leq C_\varepsilon (1 + |z|)^{N_\varepsilon} e^{\varepsilon |\mathbf{Im} z|}.$$

Show that f is a polynomial. (Hint study \hat{f} .)

4.5.7. Try to prove directly that on $C_0^\infty(\mathbb{R}^2)$ the following norms are equivalent

- a. $(\int (|f|^2 + |\frac{\partial^2 f}{\partial x^2}|^2 + |\frac{\partial^2 f}{\partial y^2}|^2)^{1/2})$;
- b. $(\int (\sum_{|\alpha| \leq 2} |D^\alpha f|^2)^{1/2})$.

Also give a proof using Fourier transform.

4.5.8. Show that $H_{(s)}(\mathbb{R}^n)$ consists of continuous functions if $s > n/2$. Can you now prove *Sobolev's Lemma* 4.4.3?

4.5.9. Show that if $s_1 > s_2$, then $H_{(s_1)}$ can be regarded as a dense subset of $H_{(s_2)}$. Can you give a description of

$$\bigcap_{s \in \mathbb{R}} H_{(s)} \quad \text{and} \quad \bigcup_{s \in \mathbb{R}} H_{(s)}.$$

4.5.10. (Sobolev for periodic functions) Let $H_{(s)}(T)$ denote the set of periodic distributions f with the property that $\sum |n|^{2s} |\hat{f}(n)|^2$ converges. Check that $H_{(s)}(T)$ is a Hilbert space with norm given by the root of the sum in (). Show that for s nonnegative integer, $H_{(s)}(T)$ coincides with L^2 functions with derivatives of order up to s in L^2 .

4.5.11. (continued.) For $f \in \mathcal{D}(T)$ define $f^{(s)} = \sum \hat{f}(n)n^s$ where $(-1)^s = e^{i\pi s}$. Show that for integer s this formula gives the ordinary derivative of sufficiently smooth functions. Show that $(f^{(s)})^{(t)} = f^{(s+t)}$. Can you make sense of this calculus for complex values of s .

4.5.12. Prove that the Airy function satisfies the differential equation

$$Ai''(x) - xAi(x) = 0.$$

Let ω be a third root of unity ($\omega^3 = 1$). Show that $Ai(\omega x)$ is an other solution of this equation. How many linearly independent solutions are there? Conclusion?

4.5.13. Prove that $Ai(0) = 3^{-1/6}\Gamma(1/3)/2\pi$ and $Ai'(0) = -3^{-1/6}\Gamma(2/3)/2\pi$. (Take $\varepsilon = 1$ in 4.2.8)

4.5.14. Let A be a symmetric matrix, possibly with complex coefficients. Show that $e^{-Ax \cdot x}$ is a tempered distribution if and only if $\mathbf{Re} A$ is positive semi-definite. Assuming in addition that A is non-singular, compute the Fourier transform of $e^{-Ax \cdot x}$.

4.5.15. Suppose that $f \in \mathcal{S}$ is radial, i.e., $f(Ux) = f(x)$ for rotations U , Prove that \hat{f} is also radial. Extend to temperate distributions.

4.5.16. Let χ_a and I_a be the distributions of exercise 3.6.14. Compute their Fourier transforms.

4.5.17. (continued) Give an interpretation of the previous exercise in terms of fractional derivatives.

(Micro) Local Analysis

In many applications Fourier transform and its inverse are used to analyze the frequencies that occur in a signal and to recover the signal from its frequencies. $\mathcal{F}f(\xi)$ can be viewed as the amplitude of the frequency ξ as occurring in the whole signal $f(t)$. Naturally, one wants to localize, that is, find out which frequencies are important at a small time interval. (Here one often makes the comparison with a music score which describes the frequencies that are prevailing at each beat, and certainly is not giving a global frequency analysis of the whole piece).

A theoretically very satisfactory localization, both in time and frequency, is obtained by means of the wave front set. We deal with it in section 1. For practical purposes it is not suitable and a weak version, the so called windowed Fourier transform is used. This is the topic of section 2. In section 3 we encounter the continuous wavelet transform. It is one of the several wavelet transforms that have been used successfully in the recent past in all kind of applications. In the later sections we will discuss discrete wavelets and multiresolution analysis.

5.1. The Wave Front Set

Consider a compactly supported distribution T on \mathbb{R}^n . According to the Paley-Wiener-Schwartz theorem, T can be distinguished from the compactly supported smooth functions by the fact that $\hat{T}(\xi)$ doesn't decrease rapidly if $|\xi| \rightarrow \infty$. We will now study more precisely the set of directions ξ where $\hat{T}(\xi)$ is not rapidly decreasing.

DEFINITION 5.1.1. A subset U of $\mathbb{R}^n \setminus \{0\}$ is called *conic* if it has the property: $\xi \in U$ implies $t\xi \in U$, for all positive t . We will call a conic set closed if it is closed in $\mathbb{R}^n \setminus \{0\}$.

For later use we remark that if a closed conic set is covered by a union of open conic sets, then it is covered by a finite subcollection, because the unit sphere is compact and the behavior of conic sets is completely determined by the behavior of their intersection with the unit sphere.

For a compactly supported distribution T we consider the set $\Sigma(T)$ consisting of all $\xi \in \mathbb{R}^n \setminus \{0\}$ with the property that there is no conic neighborhood U_ξ of ξ such that

$$(5.1.1) \quad |\hat{T}(\eta)| \leq C_N(1 + |\eta|)^{-N}, \quad N = 1, 2, \dots, \quad \eta \in U_\xi.$$

$\Sigma(T)$ is clearly a closed conic set.

While $\text{SingSupp}(T)$ describes the location of the singularities of T , $\Sigma(T)$ describes the directions of the high frequencies causing them. The following lemma shows that $\Sigma(T)$ can be localized.

LEMMA 5.1.2.

Suppose that $\varphi \in C_0^\infty(\mathbb{R}^n)$ and $T \in \mathcal{E}'(\mathbb{R}^n)$. Then $\Sigma(\varphi T) \subset \Sigma(T)$.

PROOF. We have

$$(5.1.2) \quad \widehat{\varphi T}(\xi) = \frac{1}{(2\pi)^n} \int \hat{\varphi}(\eta) \hat{T}(\xi - \eta) d\eta.$$

Here $\hat{\varphi} \in \mathcal{S}$ and $|\hat{T}(\xi)| \leq C(1 + |\xi|)^N$ for some constants $C, N > 0$. We want to show that if \hat{T} is rapidly decreasing in some conic neighborhood U_{ξ_0} of ξ_0 , then so is $\widehat{\varphi T}$ in a

possibly smaller conic neighborhood of ξ . It is natural to split the domain of integration into $\{|\eta| < c|\xi|\}$, where a favourable estimate must come from \hat{T} , and its complement, where $\hat{\varphi}$ must do the job. To wit:

$$(5.1.3) \quad \begin{aligned} |(5.1.2)| &\leq \frac{1}{(2\pi)^n} \left(\int_{|\eta| \leq c|\xi|} |\dots| + \int_{|\eta| \geq c|\xi|} |\dots| \right) \\ &\leq \frac{1}{(2\pi)^n} \left(\max_{|\eta| \leq c|\xi|} |\hat{T}(\xi - \eta)| \int_{\mathbb{R}^n} |\hat{\varphi}(\eta)| d\eta \right. \\ &\quad \left. + C_M \int_{|\eta| \geq c|\xi|} (1 + |\eta|)^{-M} (1 + |\xi - \eta|)^N d\eta \right). \end{aligned}$$

Now observe that $|\eta| \geq c|\xi| \implies |\xi - \eta| \leq (1 + \frac{1}{c})|\eta|$. Hence the last integral in (5.1.3) is bounded by

$$(5.1.4) \quad \begin{aligned} C_M \int_{|\eta| \geq c|\xi|} (1 + |\eta|)^{-M} (1 + \frac{1}{c}|\eta|)^N d\eta &\leq C'_M \int_{|\eta| \geq c|\xi|} (1 + |\eta|)^{-M+N} d\eta \\ &\leq C''_M |1 + |\xi||^{-M+N+n}. \end{aligned}$$

In other words, since M is arbitrary, this integral is rapidly decreasing.

To control the max in (5.1.3), we choose c so small that $(\xi - \eta) \in U_{\xi_0}$ if ξ is in a small conic neighborhood of ξ_0 . Then this max is also rapidly decreasing, which completes the proof. \square

We will use Lemma 5.1.2 to localize Σ .

DEFINITION 5.1.3. For $T \in \mathcal{D}'$ and $x \in \mathbb{R}^n$ define

$$\Sigma_x(T) = \cap_{\varphi} \Sigma(\varphi u); \quad \varphi \in C_0^\infty(\mathbb{R}^n), \quad \varphi(x) \neq 0.$$

The *wave front set* of a distribution $T \in \mathcal{D}'$ is the set

$$\text{WF}(T) = \{(x, \xi) : x \in \mathbb{R}^n, \xi \in \Sigma_x(T)\}.$$

Observe that for the projection on the first coordinate we have, $p_1(\text{WF}(T)) = \text{SingSupp}(T)$. For the projection on the second coordinate we get $p_2(\text{WF}(T)) = \Sigma(T)$ as Proposition 5.1.5 will show. Thus $\text{WF}(T)$ combines the information on the singularities of a distribution in a very efficient way. This is known as *micro localization*.

LEMMA 5.1.4. If $(\varphi_n)_n$ is a sequence in $\mathcal{D}(\mathbb{R}^n)$ with $\varphi_n(x) \neq 0$ and $\text{Supp } \varphi \rightarrow \{x\}$, then $\Sigma(\varphi_n T) \rightarrow \Sigma_x(T)$, that is $\cap \Sigma(\varphi_n T) = \Sigma_x(T)$.

PROOF. If V is an open conic neighborhood of $\Sigma_x(T)$ then its complement V^c is covered by the union of the open conic sets $\Sigma(\varphi T)^c$, taken over $\varphi \in \mathcal{D}$, $\varphi(x) \neq 0$. Thus, as remarked after Definition 5.1.1, there exist finitely many φ_i , $i = 1, \dots, m$, such that $V^c \subset \cup_j \Sigma(\varphi_j T)^c$. It follows that $\Sigma((\varphi_1 \cdots \varphi_m)T) \subset V$. Next, if ψ is any function such that $\text{Supp } \psi \subset (\text{Supp } \varphi_1 \cdots \varphi_m)^o$, then by Lemma 5.1.4 $\Sigma(\psi T) \subset \Sigma((\varphi_1 \cdots \varphi_m)T)$. \square

PROPOSITION 5.1.5. With the notation as above, $p_2(\text{WF}(T)) = \Sigma(T)$.

PROOF. Clearly $p_2(\text{WF}(T)) = \cup_x \Sigma_x(T) \subset \Sigma(T)$. To prove the converse inclusion, notice that $p_2(\text{WF}(T))$ is a closed cone, this follows by restricting to the unit sphere in ξ -space. Suppose that $\xi \notin p_2(\text{WF}(T))$. Then for every $x \in \text{Supp}(T)$ there exists $\varphi_x \geq 0$ with $\varphi_x(x) = 1$ such that $\widehat{(\varphi_x T)}(\eta)$ is rapidly decreasing for η in a conic neighborhood of ξ .

By compactness we can find finitely many φ_{x_i} such that $\sum_i \varphi_{x_i} > 1/2$ on $\text{Supp}(T)$. Then $(\sum_i \varphi_{x_i} u)^\wedge$ will be rapidly decreasing in a conic neighbourhood of ξ . By Lemma 5.1.4 this will also hold for $T = (\frac{1}{\sum \varphi_{x_i}}) \sum \varphi_{x_i} T$. \square

EXAMPLES 5.1.6.

1. $\text{WF}(\delta) = 0 \times \mathbb{R}^n$.

2. In \mathbb{R} , $\text{WF}(\frac{1}{x+i0}) = 0 \times \mathbb{R}_{>0}$.

3. Let $V = \{x : x_n = \dots = x_{k+1} = 0\} \subset \mathbb{R}^n$. Define a tempered distribution T_V (integration over V) by

$$T_V(\varphi) = \int_V f = \int_{\mathbb{R}^k} f(x_1, \dots, x_k, 0, \dots, 0) dx_1 dx_2 \dots dx_k.$$

The wave front set is computed in the next lemma.

LEMMA 5.1.7.

$$(5.1.5) \quad \text{WF}(T_V) = V \times (V^\perp \setminus \{0\}).$$

PROOF. Let $\xi \in C_0^\infty$. Then, writing $\xi = \xi' + \xi''$, $\xi' \in V$, $\xi'' \perp V$,

$$(5.1.6) \quad \widehat{\chi T_V} = \int_{\mathbb{R}^k} \chi(x, 0) e^{-i\xi' \cdot (x_1, \dots, x_k)} dx_1 \dots dx_k.$$

This is rapidly decreasing if ξ is in a conic neighborhood of ξ_0 with $\xi_0' \neq 0$, but if we take ξ in a conic neighborhood of ξ_0 with $\xi_0' = 0$, i.e. $\xi_0 \perp V$, then ξ' can be chosen to be bounded while $\xi \rightarrow \infty$, from which we conclude that (5.1.6) is not rapidly decreasing and the claim is proved. \square

REMARK 5.1.8. Let X be open in \mathbb{R}^n . It can be shown that any relatively closed subset of $X \times (\mathbb{R}^n \setminus \{0\})$ that is conic in the second variable, can occur as WF_L for some distribution L , cf. [5], Th. 8.1.4]

THEOREM 5.1.9. *Suppose that $u, v \in \mathcal{D}'(X)$ are distributions with the property that*

$$(5.1.7) \quad (x, \xi) \in \text{WF}(u) \implies (x, -\xi) \notin \text{WF}(v).$$

Then the product uv exists and may be defined as

$$(5.1.8) \quad \mathcal{F}^{-1}(\hat{u} * \hat{v}).$$

PROOF. It is sufficient to prove this locally. Thus, let $x \in X$ and let U be a neighborhood of x with the property that for all $y_1, y_2 \in U$ we have $(y_1, \xi) \in \text{WF}(u) \implies (y_2, -\xi) \notin \text{WF}(v)$. Such a U can be found because wave front sets are closed. We may suppose that u and v are supported in U and, for simplicity, that their Fourier transforms are bounded by $C(1+|\xi|)^N$ for some fixed N . Now we show that $\hat{u} * \hat{v}$ is in \mathcal{S}' . From the assumptions it follows that we can find a finite cover of $\mathbb{R}^n \setminus \{0\}$ with open cones Γ_j such that for slightly larger cones Γ'_j , $\hat{u}(\eta)$ is rapidly decreasing on Γ'_j or $\hat{v}(\eta)$ is rapidly decreasing on $-\Gamma'_j$ (or both). Let $\xi \in \mathbb{R}^n \setminus \{0\}$, $\eta \in \Gamma_j$ be fixed. If $\eta \in \Gamma_j$ is sufficiently large, then $\xi - \eta \in -\Gamma'_j$. Thus, assuming \hat{v} rapidly decreasing on $-\Gamma'_j$, we find for fixed N and any M a C_M with

$$(5.1.9) \quad |\hat{u}(\eta)\hat{v}(\xi - \eta)| \leq C_M(1 + |\eta|)^N(1 + |\xi - \eta|)^{-M}.$$

A similar estimate is obtained when \hat{u} is rapidly decreasing on Γ_j . We conclude that $\hat{u} * \hat{v}(\xi)$ is well defined for every ξ . Moreover, this convolution grows at most like a power of $(1 + |\xi|)$. Integrating $\hat{u}(\eta)\hat{v}(\xi - \eta)$ over cones Γ_j where $\hat{u}(\eta)$ is rapidly decreasing, this follows as the proof of Lemma 5.1.2. On the other cones we make a substitution (" $\xi - \eta = \eta$ ") to obtain integrals of the form

$$\int_{\eta \in -\Gamma_j + \xi} \hat{v}(\eta)\hat{u}(\xi - \eta) d\eta.$$

Now observe that the domain of integration is contained in the union of $-\Gamma'_j$ and a ball of radius $c|\xi|$ with c depending on the angle between Γ'_j and Γ_j . The integral over Γ'_j is treated as before, while the integral over the ball of radius $c|\xi|$ is bounded by a constant times $|\xi|^{2N+n}$. This proves that $\hat{u} * \hat{v}$ is a tempered distribution. \square

REMARK 5.1.10. In [5][Chapter 8.2] a proof can be found based on analysis of the wave front set of tensor products.

5.2. The windowed Fourier Transform on \mathbb{R}

In this and the next section, we will only consider L^2 and denote the 2-norm by $\|\cdot\|$. Choose a “good” function g on \mathbb{R} . Best is with most of its mass and most of the mass of its Fourier transform centered at 0, but for the time being think of a smoothened out characteristic function of an interval I . The *Windowed Fourier transform* (associated to g) is the following function defined on \mathbb{R}^2

$$(5.2.1) \quad \mathcal{F}_{WIN}f(\xi, t) = \int e^{-i\xi x} \overline{g(x-t)} f(x) dx = \langle f, g^{\xi, t} \rangle.$$

Here $g^{\xi, t}(x) = g(x-t)e^{i\xi x}$, the inner product is the usual one of $L^2(\mathbb{R})$. Basically $\mathcal{F}_{WIN}f(\cdot, t)$ is the Fourier transform of f restricted to the interval I shifted over t . In terms of the Fourier transform of f and g we find

$$\mathcal{F}_{WIN}f(\xi, t) = \frac{1}{2\pi} e^{i\xi t} \int_{-\infty}^{\infty} \hat{f}(\eta) \hat{g}(\eta - \xi) d\eta.$$

This formula expresses that $\mathcal{F}_{WIN}f(\xi, t)$ will be very much like $e^{i\xi t} \hat{f}(\xi)$, if \hat{g} has most of its mass at the origin. Unfortunately it is impossible to have the mass of g and \hat{g} concentrated in a small neighborhood of 0. This is the fundamental reason why windowed Fourier transform has only limited possibilities. However, all information of f is contained in $\mathcal{F}_{WIN}f$ as we shall prove now.

LEMMA 5.2.1. *for f_1, f_2 and g functions in $L^2(\mathbb{R})$ we have*

$$(5.2.2) \quad \iint \mathcal{F}_{WIN}f_1(\xi, t) \overline{\mathcal{F}_{WIN}f_2(\xi, t)} d\xi dt = 2\pi \|g\|^2 \langle f_1, f_2 \rangle.$$

PROOF. We use that the left hand side of (5.2.2) is an inner product of Fourier transforms which is integrated against t , and apply the Parseval Theorem. Then

$$(5.2.3) \quad \begin{aligned} \text{L.H.S of (5.2.2)} &= \iint g(x-t) f(x) \overline{g(x-t) f(x)} dx dt \\ &= \int |g(t)|^2 dt \int f_1(x) \overline{f_2(x)} dx = 2\pi \|g\|^2 \langle f_1, f_2 \rangle. \end{aligned}$$

The middle equality follows by a change of variables and Fubini’s Theorem. \square

COROLLARY 5.2.2. *In the weak L^2 sense we have*

$$f(x) = (2\pi \|g\|^2)^{-1} \iint \mathcal{F}_{WIN}f(\xi, t) g^{\xi, t}(x) d\xi dt.$$

PROOF. Integrating both sides against an L^2 function F should give a genuine equality:

$$(5.2.4) \quad \begin{aligned} \langle f, F \rangle &= \int f(t) \overline{F(t)} dt = (2\pi \|g\|^2)^{-1} \iint \mathcal{F}_{WIN}f(\xi, t) \overline{\int g^{\xi, t}(x) F(x) dx} d\xi dt \\ &= \iint \mathcal{F}_{WIN}f(\xi, t) g^{\xi, t}(x) d\xi dt \overline{F(x)} dx. \end{aligned}$$

\square

5.3. The Continuous Wavelet Transform

There is another way of windowing, which, in retrospect, is much more natural. Again let ψ be in $L^2(\mathbb{R})$. For reasons to become clear later, we require that

$$(5.3.1) \quad \int_{\mathbb{R}} \frac{|\hat{\psi}(\xi)|^2}{|\xi|} d\xi = C_\psi$$

is finite. We set $\psi^{a,b}(x) = |a|^{-1/2}\psi(\frac{x-b}{a})$. Then $\|\psi^{a,b}\| = \|\psi\|$ (these are again the L^2 norms). We will normalize ψ and assume from now on that $\|\psi\| = 1$. The *continuous wavelet transform* of $f \in L^2(\mathbb{R})$ is now defined as

$$(5.3.2) \quad T_{WAV}f(a,b) = \langle f, \psi^{a,b} \rangle = |a|^{-1/2} \int f(x)\psi\left(\frac{x-b}{a}\right) dx.$$

Thus both $T_{WAV}f$ and $\mathcal{F}_{WIN}f$ are function on \mathbb{R}^2 and both are square integrable, (with a suitable weight in the wavelet case, cf. Lemma 5.3.1 below). The difference with windowed Fourier transform is that there the size of the window is constant, independent of the frequency to be tested for, whereas in the case of wavelet transform the size of the window is proportional to the wavelength to be sampled, that is roughly equal to $|a|$. Short waves are tested over short intervals, long waves are tested over long ones. The shape of the testing function is not changed as a, b vary. The same is true for Fourier transform, but not for its windowed version.

As far as inversion is concerned we are in an analogous situation as in the previous section.

LEMMA 5.3.1. *Suppose that ψ satisfies (5.3.1). There exists a constant C_ψ such that if $f_1, f_2 \in L^2(\mathbb{R})$, then*

$$(5.3.3) \quad \iint T_{WAV}f_1(a,b)\overline{T_{WAV}f_2(a,b)}\frac{da db}{|a|^2} = C_\psi\langle f_1, f_2 \rangle = C_\psi \int f_1(x)\overline{f_2(x)} dx.$$

PROOF. Notice that, in view of Plancherel's Theorem, we have the following equalities:

$$\langle f, \psi^{a,b} \rangle = \langle \hat{f}, \hat{\psi}^{a,b} \rangle = \langle \hat{f}(\xi), |a|^{1/2}\hat{\psi}(a\xi)e^{-ib\xi} \rangle$$

We proceed as in Lemma 5.2.1. Recognize the two inner integrals in the following line as Fourier transforms, and use the Plancherel and Fubini theorems.

$$(5.3.4) \quad \begin{aligned} \text{L.H.S. of (5.3.3)} &= \iint \frac{da db}{|a|^2} \left(\int |a|^{1/2}\hat{f}_1(\xi)\overline{\hat{\psi}(a\xi)}e^{-ib\xi}d\xi \int |a|^{1/2}\overline{\hat{f}_2(\xi)}\hat{\psi}(a\xi)e^{ib\xi}d\xi \right) \\ &= \int \frac{da}{|a|} \left(\int \hat{f}_1(\xi)\overline{\hat{\psi}(a\xi)}\hat{f}_2(\xi)\hat{\psi}(a\xi)d\xi \right) \\ &= \int \hat{f}_1(\xi)\overline{\hat{f}_2(\xi)}d\xi \int |\hat{\psi}(a\xi)|^2\frac{da}{|a|} = C_\psi\langle f_1, f_2 \rangle. \end{aligned}$$

The constant C_ψ is defined by the last equality. □

COROLLARY 5.3.2. *In the weak L^2 sense we have*

$$f(x) = \frac{1}{C_\psi} \iint T_{WAV}f(a,b)\psi^{a,b}(x)\frac{da db}{a^2}.$$

PROOF. Copy the proof of Corollary 5.2.2 □

REMARK 5.3.3. One can show actually that

$$\lim_{A,B \rightarrow \infty} \left\| f - \frac{1}{C_\psi} \iint_{\substack{|a| < A \\ |b| < B}} T_{WAV}f(a,b)\psi^{a,b}(x)\frac{da db}{a^2} \right\| = 0.$$

Here the integral is viewed as the L^2 function which represents the continuous functional

$$Lg = \iint_{\substack{|a| < A \\ |b| < B}} T_{WAV}f(a,b)\left(\int \psi^{a,b}(x)g(x)dx\right)\frac{da db}{a^2}.$$

5.4. The mathematical zoom

Wavelets provide a good means for analyzing local smoothness as is shown by the following theorems, which are due to Holschneider and Tchamitchian [4], see also [1] who actually give stronger results.

In this section $\psi \in C_0^1$ is supported in $[-1, 1]$, $\hat{\psi}(0) = 0$, and f will always be a bounded square integrable function on \mathbb{R} . Recall that f is called *Hölder continuous* of order α , ($0 < \alpha < 1$) at x_0 if there is a positive C with

$$(5.4.1) \quad |f(x_0) - f(y)| \leq C|x_0 - y|^\alpha.$$

The function f is called Hölder continuous if (5.4.1) holds for all x_0 in \mathbb{R} and the constant C is independent of x_0 .

THEOREM 5.4.1. *The following are equivalent*

- a. f is Hölder continuous of order α .
- b. $|T_{WAV} f(a, b)| = |\langle f, \psi^{a,b} \rangle| \leq C|a|^{1/2+\alpha}$ for some $C > 0$.

THEOREM 5.4.2. *If f is Hölder α at x_0 then*

$$|\langle f, \psi^{a,x_0+b} \rangle| \leq C|a|^{1/2}(|a|^\alpha + |b|^\alpha).$$

THEOREM 5.4.3. *If f has the property that*

$$|\langle f, \psi^{a,x_0+b} \rangle| \leq C|a|^{1/2}(|a|^\alpha + \frac{|b|^\alpha}{\log |b|}),$$

then f is Hölder continuous at b .

These theorems should be compared to the situation in classical Fourier theory where such a characterization, especially the pointwise one, is not available. We will only prove Theorem 5.4.1 and 5.4.2.

PROOF. Theorem 5.4.1 (\implies)

$$(5.4.2) \quad \begin{aligned} |\langle f, \psi^{a,b} \rangle| &= \left| \int_{\mathbb{R}} f(x) \psi^{a,b}(x) dx \right| \stackrel{\left(\frac{x-b}{a}=t\right)}{=} |a|^{1/2} \left| \int_{-1}^1 f(at+b) \psi(t) dt \right| \\ &= |a|^{1/2} \left| \int_{-1}^1 (f(at+b) - f(b)) \psi(t) dt \right| \leq C|a|^{1/2}|a|^\alpha. \end{aligned}$$

We used that ψ is supported in $[-1, 1]$, and $\hat{\psi}(0) = \int_{\mathbb{R}} \psi = \int_{-1}^1 \psi = 0$. □

We continue with the proof of Theorem. 5.4.2, which is very similar.

(5.4.3)

$$\begin{aligned} |\langle f, \psi^{a,b+x_0} \rangle| &= |a|^{1/2} \left| \int_{-1}^1 (f(at+b+x_0) - f(x_0)) \psi(t) dt \right| \\ &\leq C|a|^{1/2} \int_{-1}^1 |at+b|^\alpha dt \leq C|a|^{1/2} \int_{-1}^1 |at|^\alpha + |b|^\alpha dt \leq 2C|a|^{1/2}(|a|^\alpha + |b|^\alpha). \end{aligned}$$

Theorem 5.4.1 (\impliedby) requires more effort. We will split f in a L(ow) F(requency) part and a H(igh) F(requency) part. We will show that both parts are bounded and next that both parts satisfy a uniform Hölder condition for $|x - y| \leq 1$. This will prove the theorem. In fact, it would be sufficient to show that only one of the two parts is bounded, because by boundedness of f the other part would be bounded too. However, the proof for each part is instructive in itself. We will write I_a for the interval $[-|a|, |a|]$.

We have

$$f = \frac{1}{C(\psi)} \iint (T_{WAV} f)(a, b) \psi^{a,b} \frac{da db}{a^2} = f_{HF} + f_{LF},$$

with $f_{HF} = \iint_{|a| \geq 1} \dots$, and $f_{LF} = \iint_{|a| \leq 1} \dots$

Boundedness of f_{LF} :

(5.4.4)

$$\begin{aligned} |f_{LF}| &= \frac{1}{C_\psi} \left| \int_{|a| \geq 1} \int_{b \in \mathbb{R}} \left(\int_{t \in \mathbb{R}} f(t) \psi^{a,b}(t) dt \right) \psi^{a,b}(x) \frac{da db}{a^2} \right| \\ &\leq \frac{1}{C_\psi} \int_{|a| \geq 1} \int_{b \in \mathbb{R}} \|f\|_2 \|\psi\|_2 \frac{|\psi^{a,b}(x)|}{a^2} da db \leq \frac{\|f\|_2 \|\psi\|_2}{C_\psi} \int_{|a| \geq 1} \int_{b \in \mathbb{R}} \frac{|\psi(\frac{x}{a} - \frac{b}{a})|}{a^{5/2}} db da \\ &\leq \frac{1}{C_\psi} \|f\|_2 \|\psi\|_2 \|\psi\|_1 \int_{|a| \geq 1} \frac{1}{|a|^{3/2}} da < \infty. \end{aligned}$$

Hölder estimate for f_{LF} : Let $|h| < 1$, keep in mind that here $|a| > 1$. We will use that ψ is supported in $[-1, 1]$, which tells us where $\psi^{a,b}$ is supported, and apply the mean value theorem on ψ .

(5.4.5)

$$\begin{aligned} &|f_{LF}(x+h) - f_{LF}(x)| \\ &= \frac{1}{C_\psi} \left| \int_{|a| \geq 1} \int_{b \in \mathbb{R}} \left(\int_{t \in \mathbb{R}} f(t) \psi^{a,b}(t) dt \right) (\psi^{a,b}(x+h) - \psi^{a,b}(x)) \frac{da db}{a^2} \right| \\ &\leq \frac{1}{C_\psi} \int_{|a| > 1} \int_{b \in x + I_{2a}} \left(\int_{t \in b + I_a} |f(t)| |\psi(\frac{t-b}{a})| dt \right) \left| \psi(\frac{x+h-b}{a}) - \psi(\frac{x-b}{a}) \right| \frac{da db}{|a|^3} \\ &\leq \frac{C_0}{C_\psi} |h| \int_{|a| > 1} \frac{da}{|a|^4} \int_{t \in b + I_a} |f(t)| \left(\int_{b \in \mathbb{R}} |\psi(\frac{t-b}{a})| db \right) dt \\ &\leq \frac{C_0}{C_\psi} |h| \int_{|a| > 1} \frac{da}{|a|^3} \|\psi\|_1 \int_{t \in b + I_a} |f(t)| dt \\ &\leq \frac{C}{C_\psi} |h| \|\psi\|_1 \|f\|_2 \int_{|a| > 1} \frac{da}{|a|^{5/2}} = C' |h|. \end{aligned}$$

Observe that this estimate did not use Hölder continuity of f ; F_{LF} is apparently always Hölder continuous!

Boundedness of f_{HF} : Starting from estimate $|\langle f, \psi^{a,b} \rangle| \leq C|a|^{1/2+\alpha}$ we find

$$\begin{aligned} |f_{HF}(x)| &\leq C \int_{|a| \leq 1} \frac{da}{a^2} \int_{b \in \mathbb{R}} |a|^{1/2+\alpha} |a|^{-1/2} \left| \psi\left(\frac{x-b}{a}\right) \right| db \\ (5.4.6) \quad &\leq C \int_{|a| \leq 1} \frac{|a|^\alpha da}{a^2} \int_{b \in \mathbb{R}} \left| \psi\left(\frac{x-b}{a}\right) \right| db = C \|\psi\|_1 \int_{|a| \leq 1} \frac{|a|^\alpha da}{|a|} \leq \infty. \end{aligned}$$

Hölder estimate for f_{HF} : Again fix $|h| < 1$,

(5.4.7)

$$\begin{aligned} |f_{HF}(x+h) - f_{HF}(x)| &= \left| \int_{|a| \leq 1} \frac{da}{a^2} \int_{b \in \mathbb{R}} T_{WAV}^{a,b}(f) \left(\psi^{a,b}(x+h) - \psi^{a,b}(x) \right) db \right| \\ &\leq \int_{|a| \leq 1} \frac{da}{a^2} \frac{|a|^{1/2+\alpha}}{|a|^{1/2}} \int_{b \in \mathbb{R}} \left| \psi\left(\frac{x+h-b}{a}\right) - \psi\left(\frac{x-b}{a}\right) \right| db = \int_{|a| < h} \dots + \int_{h < |a| < 1} \dots \end{aligned}$$

We estimate these integrals as follows. First for $|a| < h$. Note that $\int |\psi(\frac{x+h-b}{a})| db = |a| \|\psi\|_1$, hence

$$\int_{|a| < h} \dots \leq 2 \|\psi\|_1 \int_{|a| < h} \frac{|a|^\alpha}{|a|} da = C |h|^\alpha.$$

Next, for $|a| > h$, the mean value theorem gives

$$\left| \psi\left(\frac{x+h-b}{a}\right) - \psi\left(\frac{x-b}{a}\right) \right| \leq C \frac{|h|}{|a|}.$$

We only have to integrate with respect to b over the interval $x + I_{2a}$ therefore the inner integral in (5.4.7) is majorized by $C \frac{|h|}{|a|} 4|a|$. This leads to

$$\int_{h < |a| < 1} \dots \leq C \int_{h < |a| < 1} \frac{|a|^\alpha |h|}{a^2 |a|} 4|a| \leq C' h^\alpha.$$

□

5.5. Discrete Wavelets: Frames

In real life applications $T_{WAV} f(a, b)$ is only known in a discrete set of a 's and b 's. It is reasonable to let the a be of the form

$$a = a_0^m, \quad m \in \mathbb{Z}$$

and take

$$b = nb_0 a, \quad n \in \mathbb{Z}.$$

Thus we introduce

$$\psi_{n,m} = \psi^{a_0^{-m}, nb_0 a_0^{-m}} = a_0^{-m/2} \psi(a_0^m x - nb_0).$$

There are two things that we require from our discrete wavelets:

- i. It should be possible to reconstruct f from $\langle f, \psi_{n,m} \rangle$, $n, m \in \mathbb{Z}$, in a stable manner. That is, small changes of $\langle f, \psi_{n,m} \rangle$ should only come about from small changes of f .
- ii. Every reasonable f must have a representation as $f = \sum_{n,m} c_{n,m} \psi_{n,m}$. Of course the type of convergence is negotiable.

To make life easy on ourselves but without being really very demanding, we will assume that the map

$$(5.5.1) \quad F : f \mapsto ((\langle f, \psi_{m,n} \rangle)_{m,n})$$

is continuous from $L^2(\mathbb{R})$ to $l^2(\mathbb{Z} \times \mathbb{Z})$. Requirement i. is then equivalent with

$$(5.5.2) \quad \|(\langle f, \psi_{m,n} \rangle)\| \geq C \|f\|, \quad \text{for some } C > 0.$$

In particular (5.5.2) says that F is injective. In combination with the continuity of F we obtain the following condition: There exist constants $A, B > 0$ such that

$$(5.5.3) \quad A \|f\|^2 \leq \sum_{m,n \in \mathbb{Z}} |\langle f, \psi_{m,n} \rangle|^2 \leq B \|f\|^2.$$

DEFINITION 5.5.1. Let V be a (separable) Hilbert space. A *frame* in V is a subset $\{\psi_j : j \in J\} \subset V$ with the property that there are constants $C_1, C_2 > 0$ such that

$$(5.5.4) \quad C_1 \|f\|^2 \leq \sum_j |\langle f, \psi_j \rangle|^2 \leq C_2 \|f\|^2.$$

The C_i are called the *frame constants*. A frame is called *tight* if $C_1 = C_2$. The map F from V to $l^2(J)$

$$(5.5.5) \quad f \mapsto (\langle f, \psi_j \rangle)_j$$

is called the *frame map*. It is a continuous and injective. The dimension of V is less than or equal to the dimension of $l^2(J)$.

It is clear that the $\psi_{m,n}$ in (5.5.1)-(5.5.4) form a frame and that the map F in (5.5.1) is a frame map. Of course, if J is countable we may identify J with \mathbb{N} .

EXAMPLES 5.5.2. An orthonormal base of V is a tight frame with frame constants equal to 1. Frames have to be *spanning sets* for V , but the interesting case is when ψ_j are not linearly independent. The union of two orthonormal bases of V is a tight frame with frame constants equal to 2. The set $\{(0, 1), (-\sqrt{3}/2, 1/2), (\sqrt{3}/2, 1/2)\}$ is a tight frame in \mathbb{R}^2 with frame constants $3/2$.

Tight frames behave much like orthonormal basis; e.g., *polarization* gives

LEMMA 5.5.3. *Let ψ_j be a tight frame with constant A . Then*

$$(5.5.6) \quad A\langle f, g \rangle = \sum_j \langle f, \psi_j \rangle \overline{\langle g, \psi_j \rangle}.$$

PROOF. Using (5.5.4)

$$(5.5.7) \quad \begin{aligned} A(\|f\|^2 + \|g\|^2 + 2\mathbf{Re} \langle f, g \rangle) &= A\|f + g\|^2 = \sum_j |\langle f + g, \psi_j \rangle|^2 \\ &= A(\|f\|^2 + \|g\|^2) + 2\mathbf{Re} \sum_j \langle f, \psi_j \rangle \overline{\langle g, \psi_j \rangle}. \end{aligned}$$

This gives the real part of (5.5.6). A similar computation with f replaced by if gives the complete result. \square

As before f can be obtained from Ff in the weak sense.

THEOREM 5.5.4. *Suppose that $\{\psi_j\}$ is a tight frame with constant A . Then in the weak sense*

$$Af = \sum_j \langle f, \psi_j \rangle \psi_j.$$

PROOF. Take the inner product with a $g \in L^2(\mathbb{R})$ and apply the previous lemma. \square

PROPOSITION 5.5.5. *If $\{\psi_j\}$ is a tight frame with constant 1 and for all j , $\|\psi_j\| = 1$ then $\{\psi_j\}$ is an orthonormal basis.*

PROOF. We compute for a fixed j

$$\|\psi_j\|^4 = \left(\sum_k |\langle \psi_j, \psi_k \rangle|^2 \right)^2 = |\psi_j|^4 + \text{SUM}.$$

The SUM is a sum of products of $|\langle \psi_j, \psi_k \rangle|$, $k \neq j$, and has to equal 0. Therefore $|\langle \psi_j, \psi_k \rangle| = 0$ if $k \neq j$. We are done. \square

How to recover f from Ff if the frame is not tight? Well, F is an injective map into l^2 , but most likely not surjective. We want to invert F on its image in l^2 only, that is, we are looking for a right inverse $G : l^2 \rightarrow L^2(\mathbb{R})$, i.e. a map G with $GF = \text{id}$. From linear algebra we know the solution in the finite dimensional case; it is known as “least squares method” and G is constructed as orthogonal projection on the range of F followed by inversion of F as map onto its range. The formula in this case becomes

$$(5.5.8) \quad G = (F^*F)^{-1}F^*.$$

Here F^* is the adjoint of F .

LEMMA 5.5.6. *If F is a frame map then F^*F is invertible.*

PROOF. We have by (5.5.4) that

$$(5.5.9) \quad \langle F^*Fx, x \rangle = \langle Fx, Fx \rangle \geq C_1\|x\|^2.$$

Hence F^*F is injective. (This proves invertibility in the finite dimensional case.) Next we show that R , the range of F^*F , is closed. Let $f_n = F^*Fx_n$ form a Cauchy sequence in R . Then $\|f_n - f_m\|^2 = \|\langle F^*F(x_n - x_m), (x_n - x_m) \rangle\| \geq C_1\|x_n - x_m\|^2$, thus $\{x_n\}$ is a Cauchy sequence. Let its limit be x , then f_n converges to Fx . Finally, by functional analysis $R^\perp = \text{Ker } F^*F = 0$. This implies that $R = L^2(\mathbb{R})$ and the Lemma follows from the open mapping theorem. \square

Let us introduce

$$\tilde{\psi}_j = (F^*F)^{-1}\psi_j,$$

the *dual frame*. Using (5.5.4) it is easy to check that the dual frame corresponds to the frame map $F \circ (F^*F)^{-1}$. It satisfies the dual relation

$$C_2^{-1}\|f\|^2 \leq \sum_j |\langle f, \tilde{\psi}_j \rangle|^2 \leq C_1^{-1}\|f\|^2.$$

5.6. Multiresolution Analysis, Orthogonal Bases of Wavelets

DEFINITION 5.6.1. A *multiresolution analysis* of $L^2(\mathbb{R})$ is a sequence of subspaces $V_j \subset L^2(\mathbb{R})$ with the following properties.

1. $\dots \subset V_2 \subset V_1 \subset V_0 \subset V_{-1} \subset \dots$;
2. $\overline{\cup_{j \in \mathbb{Z}} V_j} = L^2(\mathbb{R})$;
3. $\cap_{j \in \mathbb{Z}} V_j = \emptyset$;
4. $f \in V_j \equiv f(2^j \cdot) \in V_0$, $j \in \mathbb{Z}$;
5. $f \in V_0 \implies f(\cdot - n) \in V_0$, $n \in \mathbb{Z}$;
6. $\exists \varphi \in V_0$ such that $\{\varphi(\cdot - n) : n \in \mathbb{Z}\}$ is an orthonormal basis of V_0 .

EXAMPLE 5.6.2. Take $V_j = \{f \in L^2(\mathbb{R}) : f|_{[2^j k, 2^j(k+1)]} \text{ is constant a.e.}\}$. Properties 1,3,4 and 5 are clear; 2 is an easy exercise and for 6, take φ the characteristic function of $[0, 1]$.

REMARK 5.6.3. Some further notation and simple consequences of the definition: Given a multiresolution analysis V_j , let P_j denote the orthogonal projection onto V_j and let W_j denote the orthoplement of V_j in V_{j-1} , so that $V_{j-1} = V_j \oplus W_j$, and more generally for $j < J$, $V_j = V_J \oplus_{k=0}^{J-j-1} W_{J-k}$. From this and 2. one readily infers $L^2(\mathbb{R}) = \oplus_{j \in \mathbb{Z}} W_j$. Moreover,

$$f \in W_j \iff (f \in V_{j-1} \text{ and } f \perp V_j \iff f(2^j \cdot) \in V_{-1} \text{ and } f(2^j \cdot) \in V_0 \iff f(2^j \cdot) \in W_0.$$

The main result is as follows:

THEOREM 5.6.4. *Keep the notation as above. To a multiresolution analysis of $L^2(\mathbb{R})$ as in 6.1 can be associated an orthonormal basis of wavelets given by $\psi_{jk}(x) = 2^{-j/2}\psi(2^{-j}x - k)$ such that for all $f \in L^2(\mathbb{R})$*

$$(5.6.1) \quad P_{j-1}f = P_j f + \sum_k \langle f, \psi_{jk} \rangle \psi_{jk}.$$

PROOF. Observe first that (5.6.1) is equivalent with: for every j an orthonormal basis of W_j is formed by $\{\psi_{jk}, k \in \mathbb{Z}\}$. Indeed, for $f \in W_j$ we have $f = P_{j-1}f = \sum_k \langle f, \psi_{jk} \rangle \psi_{jk}$. By scaling, it is now sufficient to find $\psi \in W_0$ such that the functions $\psi_{0k}(x) = \psi(x - k)$ form an orthonormal basis of W_0 .

The function ψ must be constructed from φ and belong to W_0 . Thus we will commence to derive some properties of φ and W_0 .

First $\varphi \in V_0 \subset V_{-1}$, while $\varphi_{-1,n}$ form an orthonormal basis of V_{-1} . Therefore

$$(5.6.2) \quad \varphi(x) = \sum_n \langle \varphi, \varphi_{-1,n} \rangle \varphi_{-1,n}(x) = \sqrt{2} \sum_n h_n \varphi(2x - n),$$

where the coefficients h_n are defined by (5.6.2). We introduce $m_0(\xi) = \frac{1}{\sqrt{2}} \sum_n h_n e^{-in\xi}$. Taking the Fourier transform of (5.6.2), we obtain

$$(5.6.3) \quad \hat{\varphi}(\xi) = \frac{1}{\sqrt{2}} \sum_n h_n \hat{\varphi}(\xi/2) e^{-in\xi/2} = m_0(\xi/2) \hat{\varphi}(\xi/2).$$

Also, by orthonormality, and Parseval's theorem

$$(5.6.4) \quad \begin{aligned} \delta_{k0} &= \int_{\mathbb{R}} dx \varphi(x) \overline{\varphi(x-k)} = \int_{\mathbb{R}} d\xi |\hat{\varphi}(\xi)|^2 e^{ik\xi} \\ &= \int_0^{2\pi} e^{ik\xi} \sum_{l \in \mathbb{Z}} |\hat{\varphi}(\xi + 2\pi l)|^2. \end{aligned}$$

It follows that

$$(5.6.5) \quad \sum_{l \in \mathbb{Z}} |\hat{\varphi}(\xi + 2\pi l)|^2 = \frac{1}{2\pi} \quad \text{a.e. for } \xi \in [0, 2\pi].$$

Plugging in (6.3) we conclude that

$$(5.6.6) \quad \sum_{l \in \mathbb{Z}} |m_0(\xi + \pi l)|^2 |\hat{\varphi}(\xi + \pi l)|^2 = \frac{1}{2\pi} \quad \text{a.e. for } \xi \in [0, \pi].$$

Now m_0 has period 2π . Thus we naturally split the sum in (6.5) into a sum over even l and one over odd l . This leads to

$$\frac{1}{2\pi} = \sum_{j \in \mathbb{Z}} |m_0(\xi)|^2 |\hat{\varphi}(\xi + 2\pi j)|^2 + \sum_{j \in \mathbb{Z}} |m_0(\xi + \pi)|^2 |\hat{\varphi}(\xi + \pi + 2\pi j)|^2.$$

Applying (6.4) yields

$$(5.6.7) \quad |m_0(\xi)|^2 + |m_0(\xi + \pi)|^2 = 1 \quad \text{a.e.}$$

This formula is at first only valid on $[0, \pi] \bmod 2\pi$; for $\xi \in [\pi, 2\pi] \bmod 2\pi$ replace ξ by $\tau + \pi$ and (6.6) becomes valid on \mathbb{R} .

Next let $f \in W_0$ that is $f \in V_{-1}$ and $f \perp V_0$. We will do a similar analysis as above. Letting $f_n = \langle f, \varphi_{-1n} \rangle$, we see that $f \in V_{-1}$ is equivalent with

$$f = \sum_n f_n \varphi_{-1n} \quad \text{or} \quad \hat{f}(\xi) = \frac{1}{2\pi} \sum f_n e^{-in\xi/2} \hat{\varphi}(\xi/2) = m_f(\xi/2) \hat{\varphi}(\xi/2).$$

Here $m_f(\xi) = \sum_{n \in \mathbb{Z}} f_n e^{-in\xi}$, so that $\|m_f\|_2 = \|f\|_2$. On the other hand, $f \perp \varphi_{0k}$ is equivalent with

$$0 = \int d\xi \hat{f}(\xi) e^{ik\xi} \overline{\hat{\varphi}(\xi)} = \int_0^{2\pi} \left(\sum_{l \in \mathbb{Z}} \hat{f}(\xi + 2\pi l) \hat{\varphi}(\xi + 2\pi l) \right) e^{ik\xi} d\xi.$$

Thus

$$\sum_l \hat{f}(\xi + 2\pi l) \overline{\hat{\varphi}(\xi + 2\pi l)} \equiv 0 \quad \text{a.e.}$$

Using (6.3) the left-hand side equals

$$\sum_l \hat{f}(\xi + 2\pi l) \overline{m_0(\xi/2 + \pi l) \hat{\varphi}(\xi/2 + \pi l)} \quad (= 0).$$

Again splitting in even and odd l for the second equality below, we find

$$(5.6.8) \quad 0 = \sum_{l \in \mathbb{Z}} m_f(\xi/2 + \pi l) m_0(\xi/2 + \pi l) |\hat{\varphi}(\xi/2 + \pi l)|^2 = m_f(\xi) \overline{m_0(\xi)} + m_f(\xi + \pi) \overline{m_0(\xi + \pi)}.$$

In view of (6.6) $m_0(\xi)$ and $m_0(\xi + \pi)$ can be 0 simultaneously on a set of measure 0 only. Thus, it follows that the function $\lambda(\xi) = m_f(\xi)/\overline{m_0(\xi)}$, which, using (6.7) equals $-m_f(\xi + \pi)/\overline{m_0(\xi)}$ is well defined a.e. and

$$\lambda(\xi) + \lambda(\xi + \pi) = 0, \quad \text{a.e.}$$

This is equivalent with the statement that λ can be written as

$$\lambda(\xi) = e^{i\xi} \nu(2\xi),$$

where $\nu = \nu[f]$ has period 2π . Our conclusion is that $f \in W_0$ if and only if $f \in L^2(\mathbb{R})$ and there exists a function $\nu = \nu[f]$ with period 2π such that

$$(5.6.9) \quad \hat{f}(\xi) = \nu(\xi)e^{i\xi/2}m_0(\xi/2 + \pi)\hat{\varphi}(\xi/2).$$

At this point let us look at where we are heading. We want to find ψ such that

$$f(x) = \sum c_k[f]\psi(x - k).$$

Taking the Fourier transform of this relation we find $\hat{f}(\xi) = \sum c_k e^{-ik\xi}\hat{\psi}(\xi)$. This looks much like (6.8). In fact, we have to put

$$\hat{\psi}(\xi) = e^{i\xi/2}m_0(\xi/2 + \pi)\hat{\varphi}(\xi/2)$$

and with $\nu(\xi) = \sum \nu_k e^{-ik\xi}$, just take $c_k = \nu_k[f]$.

To finish it up, we have to see that $\psi \in W_0$ and that, moreover, the functions $\psi(x - k)$, $k \in \mathbb{Z}$, are mutually orthogonal, of norm 1 and are spanning in W_0 . Since $|m_0|$ is bounded by 1 and $\hat{\varphi} \in L^2(\mathbb{R})$, we see that $\hat{\psi} \in L^2(\mathbb{R})$ and therefore $\psi \in L^2(\mathbb{R})$; (6.8) with $\nu \equiv 1$ yields that $\psi \in W_0$. Next

$$(5.6.10) \quad \int_{\mathbb{R}} \psi(x)\psi(x - k) dx = \int_{\mathbb{R}} |\hat{\psi}(\xi)|^2 e^{ikx} d\xi = \int_0^{2\pi} \left(\sum_{l \in \mathbb{Z}} |\hat{\psi}(\xi + 2\pi l)|^2 \right) e^{ik\xi} d\xi.$$

Now, with the definition of $\hat{\psi}$, by our usual splitting of the sum over even and odd l and (6.4),

$$(5.6.11) \quad \begin{aligned} \sum_{l \in \mathbb{Z}} |\hat{\psi}(\xi + 2\pi l)|^2 &= \sum_{l \in \mathbb{Z}} |m_0(\xi/2 + \pi(l + 1))|^2 |\hat{\varphi}(\xi/2 + \pi l)|^2 \\ &= |m_0(\xi/2)|^2 + |m_0(\xi/2 + \pi)|^2 = \frac{1}{2\pi}. \end{aligned}$$

So (6.9) is the k -th Fourier coefficient of the constant function $1/2\pi$ and vanishes unless $k = 0$. We conclude that $\{\psi(\cdot - k)\}$ is an orthonormal set in W_0 .

Finally we have to show that $\{\psi(\cdot - k)\}$ is a spanning set. We have already seen that there is an expansion $f = \sum \nu_k[f]\psi(x - k)$ for every $f \in W_0$, but this is only formal. We had better check that $\nu \in L^2_{2\pi}$. We have

$$(5.6.12) \quad \begin{aligned} \frac{\|f\|_2^2}{2} &= \int_0^{2\pi} |m_f(\xi)|^2 d\xi = \int_0^\pi |\lambda(\xi)|^2 (|m_0(\xi + \pi)|^2 + |m_0(\xi)|^2) \\ &= \int_0^\pi |\lambda(\xi)|^2 = 1/2 \int_0^{2\pi} |\nu(\xi')|^2 d\xi'. \end{aligned}$$

This finishes the proof of the theorem. □

Bibliography

- [1] I. Daubechies *Ten lectures on Wavelets* CBMS-NSF Regional conference series in Applied Math. SIAM, 1992.
- [2] R.E. Edwards *Fourier series*, Vol. 1 and 2, Springer GTM 1979 and 1981.
- [3] I. Gelfand and G.E. Shilov *Generalized functions 1-3* Academic press 1977 (translated from the Russian)
- [4] Holschneider and Tchamitchian
- [5] L. Hörmander, *The analysis of Linear Partial differential operators, 1*, Grundle. Math, Wiss., Springer 198
- [6] Y. Katznelson, *An introduction to harmonic analysis*, Dover, 1976
- [7] T. H. Koornwinder, *Fourier analysis*, UvA-Notes, 1996 (these notes has been reviewed since then)
- [8] J. Lützen, *The prehistory of the theory of distributions*. Studies in the History of Mathematics and Physical Sciences, 7. Springer-Verlag, 1982.
- [9] W. Rudin, *Functional Analysis*, McGraw-Hill 1991 (2d edition)
- [10] L. Schwartz, *Théorie des distributions* , Hermann 1966
- [11] E.M. Stein and G. Weiss *Introduction to Fourier analysis on Euclidean spaces*, Princeton UP 1971
- [12] E.M. Stein and R. Shakarchi, *Fourier analysis. An introduction*, Princeton Lectures in Analysis, 1. Princeton University Press, Princeton, NJ, 2003.
- [13] J. Wiegerinck, *Functionaal analyse* UvA-Notes, 1992 (in Dutch)
- [14] A. Zygmund, *Trigonometric series*, Cambridge Univ. Press 1959 (2d edition)

Index

- E*-spectral, 9
- supporting function, 36
- Airy function, 36
- approximate identity, 21
- Banach Alaoglu Theorem, 5
- basis, 2
- bounded set, 14
- Césaro sum, 2
- Cauchy's theorem, 36
- Cauchy-Riemann, 34
- conic, 43
- continuous wavelet transform, 47
- cosine series, 3
- Dirichlet kernels, 2
- distribution, 15, 24
 - derivative of, 16
 - local equality, 17
 - singular support, 24
 - support of, 17
- distributions, 15
- dual frame, 52
- even function, 3
- Fejér kernel, 2
- Fourier coefficients, 1
- Fourier series, 1
- Fourier transform, 33, 34
 - convolution, 34
 - distribution, 34
 - of Gaussian, 36
 - windowed, 46
- Fourier-Stieltjes series, 1
- Fréchet space, 15
- frame, 50
 - tight, 50
- frame constants, 50
- frame map, 50
- Gaussian, 36
- generalized functions, 15
- good kernels, 2
- Hölder continuous, 48
- Hadamard's Theorem, 11
- Hahn-Banach theorem, 36
- Hausdorff Young Theorem, 39
- Heine-Borel property, 14
- Hilbert space, 38
- inversion formula, 33
- kernel theorem, 30
- lacunary series, 5
 - hadamard, 5
- lacunary set, 9
- Lipschitz continuous, 3, 11
- micro localization, 44
- multiresolution analysis, 52
- nowhere differentiable function, 6
- odd function, 3
- order, 15
 - of a distribution, 24
- Paley-Wiener Theorem, 33
- Paley-Wiener-Schwartz Theorem, 36
- Parseval's formula, 2
- partition of unity, 22
- Phragmén-Lindelöf Theorem, 40
- Plancherel theorem, 38
- polarization, 51
- principal value integral, 16
- Riemann-Lebesgue Lemma, 1, 33, 40
- Riesz product, 6
- Riesz representation theorem, 39
- Riesz-Thorin Convexity Theorem, 39
- Schwartz Class, 23
- seminorm, 14
- Sidon constant, 10
- Sidon set, 9
- sine series, 3
- singular support, 24
- Sobolev space, 38
- Sobolev's Lemma, 38, 40
- spanning sets, 50
- summation by parts, 3
- support, 17
- tempered distributions, 30
- tensor product, 29
 - for distributions, 29
- test functions, 21
- test space, 21

tight frame, 50
topology, 4
 strong, 4
 weak, 4
 weak-*, 5

wave front set, 44
weak topology, 4
weak-* convergence, 4
weak-* topology, 5, 15
Weierstrass' function, 6
weighted L^2 space, 38
Windowed Fourier transform, 46