

Fluctuation of Goldstone modes and the chiral transition in QCD[†]

Frithjof Karsch, BNL& Bielefeld University

- Goldstone modes in 3 and 4 dimensions

$O(N)$ models

- (2+1)-flavor QCD below T_c

$$N_\tau = 4, 6 \text{ and } 8$$

- (2+1)-flavor QCD at T_c

$$N_\tau = 4$$

- Conclusions

[†] This talk is based on preliminary numerical results obtained by the hotQCD and RBC-Bielefeld collaborations

Chiral limit of finite-T QCD on the lattice

attempts to verify the universality conjecture for the chiral phase transition through lattice calculations so far are ambiguous; even the possibility of a 1st order transition is discussed

- while Wilson fermions recover chiral symmetry only in the continuum limit, staggered fermions have a $O(2)$ rather than the full $O(4)$ symmetry for finite lattice cut-off
- some features of $O(N)$ scaling have been observed in lattice calculations with staggered fermions, e.g. 'correct scaling' of the critical coupling as function of the quark mass
- Wilson fermions seem to yield the 'correct' magnetic equation of state
- lattice calculations so far fail to show the 'correct' $O(N)$ scaling of the chiral susceptibility

2 (+1)-flavor QCD and $O(N)$ spin models

physics of QCD at low energies as well as close to the chiral phase transition is described by effective, $O(N)$ symmetric spin models

- $T = 0$: chiral symmetry breaking at $T = 0$, $m_q = 0$ as well as leading temperature and quark mass dependent corrections are related to universal properties of 4-dimensional, $O(4)$ symmetric spin models
- $T \simeq T_c$: chiral symmetry restoration at $T = T_c$, $m_q = 0$ as well as leading temperature and quark mass dependent corrections are related to universal properties of 3-dimensional, $O(4)$ symmetric spin models

R. Pisarski and F. Wilczek, PRD29 (1984) 338

K. Rajagopal and F. Wilczek, hep-ph/0011333

A. Pelissetto and E. Vicari, Phys. Rept 368 (2002) 549

Spontaneous Symmetry Breaking

$O(N)$ spin models in d -dimensions

- non-vanishing expectation value, Σ , of the scalar field, $\Phi_{||}$, parallel to the symmetry breaking field H
- $(N - 1)$ transverse (Goldstone) modes give corrections for non-zero H (spin waves); controlled by Σ and the decay constant F for Goldstone modes

$$\Sigma_H = \Sigma_0 \left(1 - \frac{N - 1}{32\pi^2} \frac{\Sigma_0 H}{F_0^4} \ln (\Sigma_0 H / F_0^2 \Lambda_\Sigma) + \mathcal{O}(H^2) \right) , \quad d = 4$$

$$\Sigma_H = \Sigma_0 \left(1 + \frac{N - 1}{8\pi} \frac{(\Sigma_0 H)^{1/2}}{F_0^3} + \mathcal{O}(H) \right) , \quad d = 3$$

P. Hasenfratz and H. Leutwyler, NPB343, 241 (1990)

D.J. Wallace and R.K.P. Zia, PRB12, 5340 (1975)

Spontaneous Symmetry Breaking (cont.)

- (chiral) susceptibilities diverge below T_c for $H \rightarrow 0$

$$\chi_H = \frac{d\Sigma_H}{dH} \sim \langle \Phi_{\parallel}^2 \rangle - \langle \Phi_{\parallel} \rangle^2 \sim \begin{cases} H^{-1/2} & , d = 3 \\ -\ln H & , d = 4 \end{cases}$$

- divergence in the zero-field (chiral) limit

$$\chi_{H=0}(T) = \begin{cases} \infty & , T \leq T_c \\ A(T - T_c)^{-\gamma} & , T > T_c \end{cases}$$

- divergence at T_c

$$\chi_H(T = T_c) = H^{1/\delta-1} \quad , \quad T = T_c$$

crit. exp. O(2) [O(4)]: $\gamma = 1.32$ [1.45], $1 - 1/\delta = 0.79$ [0.79]

O(N) spin models in 3-dimensions

influence of Goldstone modes on spontaneous symmetry breaking below T_c and the consistency with critical behavior at T_c has been established in numerical simulations

J. Engels and T. Mendes, NP B572 (2000) 289

● $T < T_c$: $\Sigma_H = c_0(T) + c_1(T)H^{1/2} \Rightarrow \chi_H = \partial\Sigma/\partial H \sim H^{-1/2}$

● $T \simeq T_c$: scaling functions, e.g.

$$\Sigma_H = H^{1/\delta} f_s(t/H^{1/\beta\delta}), \quad t = |T - T_c|/T_c$$

$$\Rightarrow H = 0: \Sigma \sim t^\beta, \quad t = 0: \Sigma \sim H^{1/\delta}$$



● magnetic equation of state incorporates both features

D.J. Wallace and R.K.P. Zia, PRB12, 5340 (1975)

J.Engels and T. Mendes, NP Proc.Suppl. 83, 700 (2000)

$$\frac{1}{\beta} \Sigma^{\delta-1} \chi_H = \tilde{c}_0 + \tilde{c}_1 y^{-1/2}, \quad y = H/\Sigma^\delta$$

3-d, O(4) models close to T_c



J. Engels and T. Mendes 2000

- condensate shows \sqrt{H} dependence; scaling sets in for smaller H closer to T_c
- magnetic equation of state reflects $O(4)$ scaling

Goldstone modes in adjoint QCD

QCD with 2-flavor, adjoint (staggered) fermions


- QCD with fermions in the adjoint representation has two distinct phase transitions with $T_{deconf} < T_{chiral}$
- the intermediate deconfined phase shows chiral behavior as expected from 3-dimensional $O(N)$ models
- the disconnected part of the chiral susceptibility diverges with $1/\sqrt{m_a}$ for all $T_{deconf} < T < T_{chiral}$:

FK and M. Lütgemeier, NPB550, 449 (1999)

J, Engels, S. Holtmann and T. Schulze, NP B724, 357 (2005)



$$\chi_m = \frac{\partial \langle \bar{\psi} \psi \rangle}{\partial m_q a} \equiv \chi^{disc} + \chi^{con}$$
$$\chi^{disc} = \frac{1}{N_\sigma^3 N_\tau} \left(\langle (\text{Tr} M^{-1})^2 \rangle - \langle \text{Tr} M^{-1} \rangle^2 \right)$$

2-flavor adjoint QCD, $N_\tau = 4$





- $T_{\text{deconf}} \leq T \leq T_{\text{chiral}}$: weak T -dependence
- Goldstone modes lead to \sqrt{m} –terms in $\langle \bar{\psi}\psi \rangle$
 $\Rightarrow 1/\sqrt{m}$ singularity in χ_m
- $O(N)$ scaling at T_c barely visible; builds up for small m_q only

2-flavor adjoint QCD, $N_\tau = 4$



- $T_{deconf} \leq T \leq T_{chiral}$: weak T -dependence
- Goldstone modes lead to \sqrt{m} –terms in $\langle \bar{\psi}\psi \rangle$
 $\Rightarrow 1/\sqrt{m}$ singularity in χ_m
- $O(N)$ scaling at T_c barely visible; builds up for small m_q only

2-flavor adjoint QCD, $N_\tau = 4$



J. Engels et al (2005)

- $T_{deconf} \leq T \leq T_{chiral}$: weak T -dependence
- Goldstone modes lead to \sqrt{m} –terms in $\langle \bar{\psi}\psi \rangle$
 $\Rightarrow 1/\sqrt{m}$ singularity in χ_m
- $O(N)$ scaling at T_c barely visible; builds up for small m_q only

Goldstone modes in (2+1)-flavor QCD

(RBC-Bielefeld and hotQCD Collaborations)

QCD with 2 light and a 'physical' strange quark mass;
staggered fermions, p4 and asqtad actions, RHMC simulations

- calculations have been performed on $N_\sigma^3 N_\tau$ lattices for $N_\tau = 4, 6$ and 8
- check volume dependence: $4 \leq N_\sigma / N_\tau \leq 8$
- at present, most detailed analysis for $N_\tau = 4$:

$$0.0125 \leq m_l / m_s \leq 0.4$$



physical value: $m_l / m_s \simeq 0.05 \Rightarrow 70 \text{ MeV} \leq m_\pi \leq 320 \text{ MeV}$

\Rightarrow find evidence for $1/\sqrt{m_l}$ divergence in χ_{disc} in the symmetry broken phase

$N_\tau = 4$: chiral condensate

(RBC-Bielefeld collaboration, in preparation)

$$\langle \bar{\psi}\psi \rangle = \frac{1}{N_\sigma^3 N_\tau} \frac{n_f}{4} \frac{\partial \ln Z}{\partial m_l a}$$





- evidence for $\sqrt{m_l}$ term in $\langle \bar{\psi}\psi \rangle$

for orientation: $\beta = 3.28$ $T \simeq 188$ MeV, $\beta = 3.30$ $T \simeq 196$ MeV

$$N_\tau = 4: 0.0125 \leq m_l/m_s \leq 0.4$$

(RBC-Bielefeld collaboration, in preparation)

$$\chi_{disc} = \frac{1}{N_\sigma^3 N_\tau} \left(\frac{n_f}{4}\right)^2 \left(\langle (\text{Tr} M^{-1})^2 \rangle - \langle \text{Tr} M^{-1} \rangle^2 \right)$$



- evidence for $1/\sqrt{m_l}$ singularity in χ_{disc}



for orientation: $\beta = 3.28$ $T \simeq 188$ MeV, $\beta = 3.30$ $T \simeq 196$ MeV

- scaling for $m_l/m_s \lesssim 0.1$ all the way to the (pseudo-)critical temperature
- scaling sets in for smaller quark masses closer to T_c (similar to 3-d, O(N) models)

$$N_\tau = 8: 0.05 \leq m_l/m_s \leq 0.2$$

p4-data:



(RBC-Bielefeld and hotQCD collaborations)



- evidence for $1/\sqrt{m_l}$ singularity in a wide temperature range
- steep edge at high temperature is approximately quark mass independent
(as expected: $\langle \bar{\psi}\psi \rangle \sim c(T)m_l \Rightarrow \chi_m \sim c(T)$)

$N_\tau = 8$: p4 and asqtad



hotQCD and RBC-Bielefeld collaborations, preliminary



● p4 and asqtad calculation lead to similar quark mass dependence

$N_\tau = 8$: p4 and asqtad


hotQCD and RBC-Bielefeld collaborations, preliminary



- p4 and asqtad calculation lead to similar quark mass dependence
- the rapid drop at large temperature is consistent with the expected O(2) [O(4)] scaling; however no 'critical behavior' of peak heights
- to firmly establish these features requires a more thorough analysis of the (m_l, T) -dependence of $\chi_{l,disc}$

$N_\tau = 4$: $O(2)$ [$O(4)$] scaling at β_c

determining β_c using the χ^2 -method: J. Engels et al., PLB298 (1993) 154



fit: $\langle \bar{\psi}\psi \rangle = am_l^{1/\delta} + bm_l$, $1/\delta = 0.21$

minimize $\chi^2 \Rightarrow \beta_c$; in agreement with earlier determination:

M. Cheng et al, PRD74 (2006) 054507

error on β_c : $\chi^2 \leq 2 \Rightarrow$ narrow scaling window $\Delta T \simeq \pm 1.5$ MeV

evidence for $O(N)$ scaling at T_c for $m_l \lesssim m_{phys}$

Conclusions

- Goldstone modes control properties of the chiral condensate and its susceptibility in the confined phase
 - 3-d, $O(N)$ scaling close to T_c
- (2+1)-flavor QCD
 - the chiral susceptibility diverges like $1/\sqrt{m_l}$ for $T \lesssim T_c$;
this scaling sets already in the regime of physical quark mass values
- in order to disentangle the thermal critical behavior of the chiral condensate and its susceptibility from the singular behavior induced by Goldstone modes a detailed analysis of the temperature and quark mass dependence close to T_c is necessary