

2nd Homework sheet Model Theory

- Deadline: 18 February.
- Submit your solutions by handing them to the lecturer at the *beginning of the lecture at 15:00*.
- Good luck!

Exercise 1 (100 points) We will call a formula φ *positive*, if it does not contain any negations \neg or implications \rightarrow ; in other words, if it can be obtained from atomic formulas using only \wedge, \vee, \exists and \forall . In addition, we will call a homomorphism $f: M \rightarrow N$ of \mathcal{L} -structures *positive*, if

$$M \models \varphi(m_1, \dots, m_n) \Rightarrow N \models \varphi(f(m_1), \dots, f(m_n))$$

holds for all positive formulas $\varphi(x_1, \dots, x_n)$ and all $m_1, \dots, m_n \in M$.

- (a) (40 points) Let T be a consistent \mathcal{L} -theory and write

$$T_0 = \{\psi : \psi \text{ is a positive sentence and } T \models \psi\}.$$

Prove that for any model A of T_0 there is a diagram of \mathcal{L} -structures

$$\begin{array}{ccc} & & B \\ & & \downarrow l \\ A & \xrightarrow{k} & C \end{array}$$

such that: (1) B is a model of T , (2) k is an elementary embedding, (3) l is a positive homomorphism, and (4) the image of k is contained in the image of l .

- (b) (30 points) Let $f: D \rightarrow A$ be a positive homomorphism of \mathcal{L} -structures. Prove that there exists a commuting square of \mathcal{L} -structures

$$\begin{array}{ccc} D & \xrightarrow{k} & B \\ f \downarrow & & \downarrow g \\ A & \xrightarrow{l} & C \end{array}$$

in which the horizontal maps k and l are elementary embeddings, the vertical maps f and g are positive homomorphisms and the image of l is contained in the image of g .

- (c) (30 points) Let T be a consistent \mathcal{L} -theory whose models are closed under surjective images: so if $f: M \rightarrow N$ is a surjective homomorphism and M is a model of T , then so is N . Use parts (a) and (b) to prove that T can be axiomatised using positive sentences.