

Term extraction

One would expect that implicit in a constructive proof of an existential statement like $\exists x^\sigma \varphi(x)$ would be a method for finding a term t such that $\varphi(t)$ holds. Below we will show that this is correct for HA^ω . More generally, we will show that one can extract from a proof of

$$\forall x^\sigma \exists y^\tau \varphi(x, y)$$

a term t such that $\forall x^\sigma \varphi(x, tx)$ holds. In order to show this we need to modify modified realizability and that is what we will do in the first section.

In the second section we will show that similar results hold for PA^ω as well, provided φ is simple enough.

1. Modified realizability with truth

In order to show our term extract results we need to modify modified realizability in such a way that HA^ω believes that anything which is realized is also true. We have seen that for ordinary modified realizability this is not the case, because AC is both unprovable and provably *mr*-realized in HA^ω . To get what we want, we only have to make a small adaptation to the definition of *mr*: indeed, the notion of a type $\text{tp}(\varphi)$ of a formula φ is as before and we only add a clause saying that $\varphi \rightarrow \psi$ can only be realized if it is true. The result is called “modified realizability with truth” and is defined as follows:

DEFINITION 1.1. To any formula φ in the language of HA^ω we associate a new formula $x \text{ mrt } \varphi$ as follows, where $x \text{ mrt } \varphi$ is also a formula in the language of HA^ω whose free variables are those of φ plus possibly a variable x of type $\text{tp}(\varphi)$:

$$\begin{aligned} x \text{ mrt } \varphi &:= \varphi && \text{if } \varphi \text{ is atomic.} \\ x \text{ mrt } (\varphi \wedge \psi) &:= \mathbf{p}_0 x \text{ mrt } \varphi \wedge \mathbf{p}_1 x \text{ mrt } \psi \\ x \text{ mrt } (\varphi \rightarrow \psi) &:= (\varphi \rightarrow \psi) \wedge \forall y^{\text{tp}(\varphi)} (y \text{ mrt } \varphi \rightarrow x(y) \text{ mrt } \psi) \\ x \text{ mrt } \exists y^\sigma \varphi &:= \mathbf{p}_1 x \text{ mrt } \varphi(\mathbf{p}_0 x) \\ x \text{ mrt } \forall y^\sigma \varphi &:= \forall y^\sigma (x(y) \text{ mrt } \varphi) \end{aligned}$$

PROPOSITION 1.2. For any formula φ in the language of HA^ω we have

$$\text{HA}^\omega \vdash (x \text{ mrt } \varphi) \rightarrow \varphi.$$

PROOF. By induction on the structure of φ . □

THEOREM 1.3. Let φ be a formula in the language of HA^ω . If φ is provable in HA^ω , then one can find effectively from this proof a term t in the language of HA^ω such that:

- (1) any variables occurring freely in t also occur freely in φ , and
- (2) $\text{HA}^\omega \vdash t \text{ mrt } \varphi$.

The same statement holds for $\mathbf{E}\text{-HA}^\omega$.

PROOF. As for modified realizability. □

COROLLARY 1.4. (Term extraction for \mathbf{HA}^ω) *Consider a formula of the form $\forall x^\sigma \exists y^\tau \varphi(x, y)$. If this formula is provable in \mathbf{HA}^ω , then from this proof one can effectively extract a term t of type $\sigma \rightarrow \tau$ such that*

$$\mathbf{HA}^\omega \vdash \forall x^\sigma \varphi(x, tx).$$

In particular, if a formula of the form $\exists x^\sigma \varphi(x)$ is provable in \mathbf{HA}^ω , then one can find a term t of type σ such that $\varphi(t)$ is provable in \mathbf{HA}^ω as well. The same statements hold for $\mathbf{E}\text{-HA}^\omega$ as well.

PROOF. If $\forall x^\sigma \exists y^\tau \varphi(x, y)$ is provable, then it follows from the soundness of modified realizability with truth that there is a term s such that

$$\mathbf{HA}^\omega \vdash s \text{ mrt } \forall x^\sigma \exists y^\tau \varphi(x, y).$$

By definition of mrt this means that

$$\mathbf{HA}^\omega \vdash \forall x^\sigma \mathbf{p}_1(sx) \text{ mrt } \varphi(x, \mathbf{p}_0(sx)).$$

So if we put $t = \lambda x^\sigma. \mathbf{p}_0(sx)$, then

$$\mathbf{HA}^\omega \vdash \forall x^\sigma \mathbf{p}_1(sx) \text{ mrt } \varphi(x, tx).$$

But then Proposition 1.2 implies that

$$\mathbf{HA}^\omega \vdash \forall x^\sigma \varphi(x, tx),$$

as desired. □

2. Classical arithmetic

In this section we will show a term extraction result for classical arithmetic. This will be a consequence of the combination of two things: term extraction for \mathbf{HA}^ω and the fact \mathbf{PA}^ω is Π_2^0 -conservative over \mathbf{HA}^ω (that is, \mathbf{PA}^ω proves the same Π_2^0 -formulas as \mathbf{HA}^ω). Since we already proved the former, we only have to prove the latter. But before we can do that, we first have to define the notion of a Π_2^0 -formula and prove two lemmas.

DEFINITION 2.1. We will call a formula φ *simple* if it contains neither quantifiers nor equalities $=_\sigma$ with $\sigma \neq 0$ (this is not standard terminology). A formula of the form

$$\forall x^\sigma \exists y^\tau \varphi$$

where φ is simple is called a Π_2^0 -*formula*.

LEMMA 2.2. *For any simple formula φ with free variables among $x_1^{\sigma_1}, \dots, x_n^{\sigma_n}$ there is a closed term \mathbf{d} of type $\sigma \rightarrow \dots \rightarrow \sigma_n \rightarrow 0$ such that*

$$\mathbf{HA}^\omega \vdash \forall x_1^{\sigma_1}, \dots, \forall x_n^{\sigma_n} (\varphi \leftrightarrow \mathbf{d}x_1 \dots x_n =_0 0).$$

PROOF. Exercise! □

LEMMA 2.3. Let ∇ be a function mapping formulas in the language of HA^ω to formulas in the language in HA^ω , such that the following statements are provable in HA^ω :

$$\begin{aligned} \text{HA}^\omega &\vdash \varphi \rightarrow \nabla\varphi \\ \text{HA}^\omega &\vdash \nabla(\varphi \wedge \psi) \leftrightarrow (\nabla\varphi \wedge \nabla\psi) \\ \text{HA}^\omega &\vdash (\varphi \rightarrow \nabla\psi) \rightarrow (\nabla\varphi \rightarrow \nabla\psi) \end{aligned}$$

In addition, let φ^∇ be the formula obtained from φ by applying ∇ to each atomic subformula, disjunction and existentially quantified subformula. More precisely, φ^∇ is defined by induction on the structure of φ as follows:

$$\begin{aligned} \varphi^\nabla &:= \nabla\varphi && \text{if } \varphi \text{ is an atomic formula,} \\ (\varphi \wedge \psi)^\nabla &:= \varphi^\nabla \wedge \psi^\nabla, \\ (\varphi \vee \psi)^\nabla &:= \nabla(\varphi^\nabla \vee \psi^\nabla), \\ (\varphi \rightarrow \psi)^\nabla &:= \varphi^\nabla \rightarrow \psi^\nabla, \\ (\forall x \varphi(x))^\nabla &:= \forall x (\varphi(x))^\nabla, \\ (\exists x \varphi(x))^\nabla &:= \nabla\exists x (\varphi(x))^\nabla. \end{aligned}$$

Then $\text{HA}^\omega \vdash \varphi$ implies $\text{HA}^\omega \vdash \varphi^\nabla$.

PROOF. Since we already know this for predicate logic, it suffices to check that φ^∇ is provable in HA^ω if φ is a non-logical axiom of HA^ω . We leave this as an exercise. \square

In particular, we have that if φ is provable in PA^ω then $\varphi^{\neg\neg}$ (its double negation translation) is provable in HA^ω . But we also have the following:

THEOREM 2.4. Any Π_2^0 -formula provable in PA^ω is also provable in HA^ω .

PROOF. Let φ be a Π_2^0 -formula. In view of Lemma 2.2 we may assume that φ is of the form

$$\forall x^\sigma \exists y^\tau t(x, y) =_0 0.$$

We know that for any formula A the nucleus $\nabla\varphi := (\varphi \rightarrow A) \rightarrow A$ interprets classical logic. This means that if φ is provable in PA^ω , then

$$\text{HA}^\omega \vdash \forall x^\sigma \nabla(\exists y^\tau \nabla(t(x, y) = 0)).$$

In other words, we have

$$\text{HA}^\omega \vdash \forall x^\sigma (([\exists y^\tau ((t(x, y) = 0 \rightarrow A) \rightarrow A)] \rightarrow A) \rightarrow A)$$

for any formula A . But if we choose A to be $\exists y^\tau t(x, y) = 0$, then this is equivalent to

$$\text{HA}^\omega \vdash \forall x^\sigma \exists y^\tau t(x, y) = 0.$$

So φ is provable in HA^ω . \square

The proof method that we employed above is often called Friedman's trick (and is due to Harvey Friedman).

COROLLARY 2.5. (Term extraction for PA^ω) Consider a formula of the form $\forall x^\sigma \exists y^\tau \varphi(x, y)$, where φ is simple. If this formula is provable in PA^ω , then from this proof one can effectively extract a term t of type $\sigma \rightarrow \tau$ such that

$$\text{PA}^\omega \vdash \forall x^\sigma \varphi(x, tx).$$

In particular, if a formula of the form $\exists x^\sigma \varphi(x)$ with φ simple is provable in PA^ω , then one can find a term t of type σ such that $\varphi(t)$ is provable in PA^ω as well.

PROOF. Follows from Theorem 2.4 and Corollary 1.4. □

REMARK 2.6. Apart from Lemma 2.2 all the results in this section fail in the presence of extensionality. More precisely:

- (1) The double negation translation of the axiom of extensionality is not provable in E-HA^ω .
- (2) There are Π_2^0 -formulas provable in E-PA^ω which are not provable in E-HA^ω .
- (3) There are Π_2^0 -formulas provable in E-PA^ω from which no closed terms in the language of E-PA^ω can be extracted.

Proofs of these facts are beyond the scope of this course.