

THE ZEROS OF HERON'S FORMULA IN ORTHOCENTRIC TETRAHEDRA

Timothy F. Havel and Garret Sobczyk

MIT and UDLAP

August, 2024



TABLE OF CONTENTS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- 1 Low-dimensional Euclidean geometry is not so elementary
- 2 Heron's formula and its relation to incenter geometry
- 3 Extending Heron's formula to tetrahedra (and beyond ...)
- 4 But its zeros are **really** weird in 3 (& higher?) dimensions
- 5 Orthocentric tetrahedra as extensions of triangles to 3D
- 6 Orthocentric systems in 2 & 3D, and in/excenter geometry
- 7 Open Questions and Future Work



$\mathcal{G}_{3,1}$: A VECTOR SPACE MODEL WITH MANY DIFFERENT GEOMETRIC INTERPRETATIONS

- Even subalgebra of $\mathcal{G}_{3,1}$ is $O^+(3, 1) \approx O^+(1, 3) \approx SL(2, \mathbb{C})$
- So it represents the Lorentz group of space-time (like $\mathcal{G}_{1,3}$)
- And $\mathbb{R}^{3,1}$ contains models of several well-known geometries:
 - Lines inside the null cone: hyperbolic 3-space;
 - Lines on the null cone: the inversive (conformal) plane;
 - A parabolic section of the null cone: the Euclidean plane.

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



$\mathcal{G}_{3,1}$: A VECTOR SPACE MODEL WITH MANY DIFFERENT GEOMETRIC INTERPRETATIONS

- Even subalgebra of $\mathcal{G}_{3,1}$ is $O^+(3, 1) \approx O^+(1, 3) \approx SL(2, \mathbb{C})$
- So it represents the Lorentz group of space-time (like $\mathcal{G}_{1,3}$)
- And $\mathbb{R}^{3,1}$ contains models of several well-known geometries:
 - Lines inside the null cone: hyperbolic 3-space;
 - Lines on the null cone: the inversive (conformal) plane;
 - A parabolic section of the null cone: the Euclidean plane.

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra


Orthocentric
systems

Open
questions



$\mathcal{G}_{3,1}$: A VECTOR SPACE MODEL WITH MANY DIFFERENT GEOMETRIC INTERPRETATIONS

- Even subalgebra of $\mathcal{G}_{3,1}$ is $O^+(3, 1) \approx O^+(1, 3) \approx SL(2, \mathbb{C})$
- So it represents the Lorentz group of space-time (like $\mathcal{G}_{1,3}$)
- And $\mathbb{R}^{3,1}$ contains models of several well-known geometries:
 - Lines inside the null cone: hyperbolic 3-space;
 - Lines on the null cone: the inversive (conformal) plane;
 - A parabolic section of the null cone: the Euclidean plane.



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



$\mathcal{G}_{3,1}$: A VECTOR SPACE MODEL WITH MANY DIFFERENT GEOMETRIC INTERPRETATIONS

- Even subalgebra of $\mathcal{G}_{3,1}$ is $O^+(3, 1) \approx O^+(1, 3) \approx SL(2, \mathbb{C})$
- So it represents the Lorentz group of space-time (like $\mathcal{G}_{1,3}$)
- And $\mathbb{R}^{3,1}$ contains models of several well-known geometries:
 - Lines inside the null cone: hyperbolic 3-space;
 - Lines on the null cone: the inversive (conformal) plane;
 - A parabolic section of the null cone: the Euclidean plane.

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra


The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

Unfortunately, the latter is generally seen as "high-school" geometry, beneath the dignity of professional mathematicians!






$\mathcal{G}_{3,1}$: A VECTOR SPACE MODEL WITH MANY DIFFERENT GEOMETRIC INTERPRETATIONS

- Even subalgebra of $\mathcal{G}_{3,1}$ is $O^+(3, 1) \approx O^+(1, 3) \approx SL(2, \mathbb{C})$
- So it represents the Lorentz group of space-time (like $\mathcal{G}_{1,3}$)
- And $\mathbb{R}^{3,1}$ contains models of several well-known geometries:
 - Lines inside the null cone: hyperbolic 3-space;
 - Lines on the null cone: the inversive (conformal) plane;
 - A parabolic section of the null cone: the Euclidean plane.

Unfortunately, the latter is generally seen as “high-school” geometry, beneath the dignity of professional mathematicians!



But 2 & 3D Euclidean geometry enables one to visualize the null cone of a Lorentian vector space two dimensions higher (and so is maybe not so “elementary” after all).

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



TABLE OF CONTENTS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- 1 Low-dimensional Euclidean geometry is not so elementary
- 2 Heron's formula and its relation to incenter geometry
- 3 Extending Heron's formula to tetrahedra (and beyond ...)
- 4 But its zeros are **really** weird in 3 (& higher?) dimensions
- 5 Orthocentric tetrahedra as extensions of triangles to 3D
- 6 Orthocentric systems in 2 & 3D, and in/excenter geometry
- 7 Open Questions and Future Work



CASE IN POINT: HERON'S FORMULA FOR THE AREA OF A TRIANGLE

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions





CASE IN POINT: HERON'S FORMULA FOR THE AREA OF A TRIANGLE

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra


Orthocentric
systems

Open
questions

Although better known as a means of calculating the area of a triangle \overline{ABC} , Heron's formula is essentially a relation between the edge lengths a, b, c and the **squared** radius r of its incircle:

$$(2r)^2 = (-a + b + c)(a - b + c)(a + b - c)/(a + b + c)$$

Squared area $t^2 = r^2 s^2$ with *semi-perimeter* s defined as shown.



$$r = t/s \text{ where}$$

$$s := \frac{1}{2}(a + b + c) \\ = u + v + w$$

and

$$t := |\overline{ABC}|$$



CASE IN POINT: HERON'S FORMULA FOR THE AREA OF A TRIANGLE

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions




If $u = (-a + b + c)/2$, $v = (a - b + c)/2$, $w = (a + b - c)/2$ are the lengths of the segments shown in the figure, this may be written more compactly as

$$r^2 = \frac{1}{2s} \Omega \Leftrightarrow t^2 = \frac{s}{2} \Omega, \quad \Omega(u, v, w) := \det \begin{bmatrix} 0 & u & v \\ u & 0 & w \\ v & w & 0 \end{bmatrix},$$

wherein $\Omega = 2uvw$. Since $a = v + w$, $b = u + w$, $c = u + v$, the **Heron parameters** u, v, w determine the triangle up to isometry.

TRIGONOMETRIC VERSIONS OF THE HERON PARAMETERS & IN-TOUCH TRIANGLE EDGES



Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra

Orthocentric systems

Open questions



TRIGONOMETRIC VERSIONS OF THE HERON PARAMETERS & IN-TOUCH TRIANGLE EDGES


Basic trigonometry yields the following relations among the angles & distances in the triangle and its *in-touch triangle* \overline{JKL} :

$$\psi_J = 2\phi_J, \quad \psi_K = 2\phi_K, \quad \psi_L = 2\phi_L;$$

$$u = r \cot(\theta_A/2) = r \tan(\phi_J), \quad j = 2r \cos(\theta_A/2) = 2r \sin(\phi_J);$$


$$v = r \cot(\theta_B/2) = r \tan(\phi_K), \quad k = 2r \cos(\theta_B/2) = 2r \sin(\phi_K);$$

$$w = r \cot(\theta_C/2) = r \tan(\phi_L), \quad \ell = 2r \cos(\theta_C/2) = 2r \sin(\phi_L).$$





TRIGONOMETRIC VERSIONS OF THE HERON PARAMETERS & IN-TOUCH TRIANGLE EDGES



Eliminating r and applying the laws of sines & cosines then yields the algebraic relations between u, v, w and j, k, ℓ :

$$u = jk\ell / (-j^2 + k^2 + \ell^2), \quad j = 4u^2vw / (bc) ;$$

$$v = jk\ell / (j^2 - k^2 + \ell^2), \quad k = 4uv^2w / (ac) ;$$

$$w = jk\ell / (j^2 + k^2 - \ell^2), \quad \ell = 4uvw^2 / (ab)$$

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



GETTING THE IN-TOUCH TRIANGLE'S AREA WITHOUT USING HERON'S FORMULA


Since the vectors $\mathbf{j}, \mathbf{k}, \mathbf{l} \in \mathbb{R}^2$ representing the *in-touch points* $\bar{J}, \bar{K}, \bar{L}$ are barycentric sums of the triangle's vertices $\mathbf{a}, \mathbf{b}, \mathbf{c}$, i.e.

$$\mathbf{j} = \frac{w}{a} \mathbf{b} + \frac{v}{a} \mathbf{c}, \quad \mathbf{k} = \frac{w}{b} \mathbf{a} + \frac{u}{b} \mathbf{c}, \quad \mathbf{l} = \frac{v}{c} \mathbf{a} + \frac{u}{c} \mathbf{b},$$


the conformal blades of the triangle & in-touch triangle satisfy

$$abc \mathbf{n}_\infty \wedge \mathbf{j} \wedge \mathbf{k} \wedge \mathbf{l} = \Omega(u, v, w) \mathbf{n}_\infty \wedge \mathbf{a} \wedge \mathbf{b} \wedge \mathbf{c},$$

wherein $\mathbf{a} := \mathbf{n}_0 + \mathbf{a} + \mathbf{n}_\infty \mathbf{a}^2/2 \in \mathcal{G}_{3,1}$ etc. are **conformal** points.



GETTING THE IN-TOUCH TRIANGLE'S AREA WITHOUT USING HERON'S FORMULA



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



TABLE OF CONTENTS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- 1 Low-dimensional Euclidean geometry is not so elementary
- 2 Heron's formula and its relation to incenter geometry
- 3 Extending Heron's formula to tetrahedra (and beyond ...)**
- 4 But its zeros are **really** weird in 3 (& higher?) dimensions
- 5 Orthocentric tetrahedra as extensions of triangles to 3D
- 6 Orthocentric systems in 2 & 3D, and in/excenter geometry
- 7 Open Questions and Future Work



HERON FOR TETRAHEDRA, CONCEPT #1: MEDIAL PARALLELOGRAMS & OCTAHEDRON

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



NB: parallel line segments in space have been drawn in the same colors, and the medial parallelogram \overline{XUWZ} is drawn in **bold**; the diagonals of the octahedron \overline{UZ} , \overline{VY} , \overline{WX} , or the *bimedians* of the tetrahedron, are omitted for simplicity.

The **medial octahedron** of a tetrahedron \overline{ABCD} is spanned by the midpoints \bar{U} , ..., \bar{Z} of its edges. This octahedron's edges are pairwise parallel to those of the tetrahedron (in same color) but only half as long.

The octahedron's 3 diagonals intersect at its centroid \bar{G} , and any two of these diagonals are the diagonals of one of the tetrahedron's three **medial parallelograms** (also called "interior faces" in following).



HERON FOR TETRAHEDRA, CONCEPT #1: MEDIAL PARALLELOGRAMS & OCTAHEDRON

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



NB: parallel line segments in space have been drawn in the same colors, and the medial parallelogram \overline{XUWZ} is drawn in **bold**; the diagonals of the octahedron \overline{UZ} , \overline{VY} , \overline{WX} , or the *bimedians* of the tetrahedron, are omitted for simplicity.

The **medial octahedron** of a tetrahedron \overline{ABCD} is spanned by the midpoints \bar{U} , ..., \bar{Z} of its edges. This octahedron's edges are pairwise parallel to those of the tetrahedron (in same color) but only half as long.

The octahedron's 3 diagonals intersect at its centroid \bar{G} , and any two of these diagonals are the diagonals of one of the tetrahedron's three **medial parallelograms** (also called "interior faces" in following).



CONCEPT #1 (CONT): REPRESENTING MEDIAL PARALLELOGRAMS IN GEOMETRIC ALGEBRA

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



In \mathcal{G}_3 , the bivector of a medial parallelogram of \overline{ABCD} is e.g.

$$\begin{aligned} \mathbf{P} &:= \left(\frac{1}{2}(\mathbf{b} + \mathbf{c}) - \frac{1}{2}(\mathbf{a} + \mathbf{c})\right) \\ &\quad \wedge \left(\frac{1}{2}(\mathbf{d} + \mathbf{a}) - \frac{1}{2}(\mathbf{c} + \mathbf{a})\right) \\ &= \frac{1}{4}(\mathbf{b} - \mathbf{a}) \wedge (\mathbf{d} - \mathbf{c}) \end{aligned}$$

In the conformal model $\mathcal{G}_{4,1}$, its plane-bound bivector is related to the commutator product \bowtie of line-bound vectors of opposite pairs of edges, e.g. $\mathbf{L} := n_\infty \wedge \mathbf{a} \wedge \mathbf{b}$ and $\mathbf{M} := n_\infty \wedge \mathbf{c} \wedge \mathbf{d}$, as

$$n_\infty \wedge \mathbf{g} \wedge \mathbf{P} = n_\infty \wedge \mathbf{g} \wedge (\mathbf{L} \bowtie \mathbf{M})$$

with \mathbf{g} any point in the mid-plane.



CONCEPT #1 (CONT): REPRESENTING MEDIAL PARALLELOGRAMS IN GEOMETRIC ALGEBRA

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



In \mathcal{G}_3 , the bivector of a medial parallelogram of \overline{ABCD} is e.g.

$$\begin{aligned} \mathbf{P} &:= \left(\frac{1}{2}(\mathbf{b} + \mathbf{c}) - \frac{1}{2}(\mathbf{a} + \mathbf{c})\right) \\ &\quad \wedge \left(\frac{1}{2}(\mathbf{d} + \mathbf{a}) - \frac{1}{2}(\mathbf{c} + \mathbf{a})\right) \\ &= \frac{1}{4}(\mathbf{b} - \mathbf{a}) \wedge (\mathbf{d} - \mathbf{c}) \end{aligned}$$

In the conformal model $\mathcal{G}_{4,1}$, its plane-bound bivector is related to the commutator product \bowtie of line-bound vectors of opposite pairs of edges, e.g. $\mathbf{L} := \mathbf{n}_\infty \wedge \mathbf{a} \wedge \mathbf{b}$ and $\mathbf{M} := \mathbf{n}_\infty \wedge \mathbf{c} \wedge \mathbf{d}$, as

$$\mathbf{n}_\infty \wedge \mathbf{g} \wedge \mathbf{P} = \mathbf{n}_\infty \wedge \mathbf{g} \wedge (\mathbf{L} \bowtie \mathbf{M})$$

with \mathbf{g} any point in the mid-plane.



CONCEPT #2: TETRAHEDRON INEQUALITIES AMONG THE SEVEN FACIAL AREAS

Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles


Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra

Orthocentric systems

Open questions



Much as in Euler's triangle notation,

$$d := \|(\mathbf{b} - \mathbf{a}) \wedge (\mathbf{c} - \mathbf{a})\|$$

will be **twice** the area of the exterior face opposite \overline{D} , and similarly for the areas c, b, a opposite $\overline{C}, \overline{B}, \overline{A}$, while

$$e := \|(\mathbf{b} - \mathbf{a}) \wedge (\mathbf{d} - \mathbf{c})\|$$

will be **4 times** that of the indicated interior face, and similarly for f, g .

Then by the triangle inequality for Euclidean (bi)vectors, we have

$$\begin{aligned}
 (\mathbf{b} - \mathbf{a}) \wedge (\mathbf{d} - \mathbf{c}) &= (\mathbf{b} - \mathbf{a}) \wedge (\mathbf{d} - \mathbf{a}) - (\mathbf{b} - \mathbf{a}) \wedge (\mathbf{c} - \mathbf{a}) \\
 \implies e &\leq c + d \quad \text{as well as} \quad c \leq d + e, \quad d \leq c + e.
 \end{aligned}$$

All in all, we get 18 such **tetrahedron inequalities**, which are stronger than the better-known areal inequalities $a \leq b + c + d$ etc.



CONCEPT #2: TETRAHEDRON INEQUALITIES AMONG THE SEVEN FACIAL AREAS

Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles


Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra

Orthocentric systems

Open questions



Much as in Euler's triangle notation,

$$d := \|(\mathbf{b} - \mathbf{a}) \wedge (\mathbf{c} - \mathbf{a})\|$$

will be **twice** the area of the exterior face opposite \overline{D} , and similarly for the areas c, b, a opposite $\overline{C}, \overline{B}, \overline{A}$, while

$$e := \|(\mathbf{b} - \mathbf{a}) \wedge (\mathbf{d} - \mathbf{c})\|$$

will be **4 times** that of the indicated interior face, and similarly for f, g .

Then by the triangle inequality for Euclidean (bi)vectors, we have

$$\begin{aligned}
 (\mathbf{b} - \mathbf{a}) \wedge (\mathbf{d} - \mathbf{c}) &= (\mathbf{b} - \mathbf{a}) \wedge (\mathbf{d} - \mathbf{a}) - (\mathbf{b} - \mathbf{a}) \wedge (\mathbf{c} - \mathbf{a}) \\
 \implies e &\leq c + d \quad \text{as well as} \quad c \leq d + e, \quad d \leq c + e.
 \end{aligned}$$

All in all, we get 18 such **tetrahedron inequalities**, which are stronger than the better-known areal inequalities $a \leq b + c + d$ etc.



CONCEPT #3: THE AREAL LAWS OF SINES & COSINES; MCCONNELL'S RIGIDITY THEOREM

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



Dihedral Angle Bisectors

The plane bisecting the dihedral angle φ_{AB} divides \overline{CD} in the ratios shown, and similarly for the other dihedrals.

The Areal Law of Cosines

The cosine of φ_{AB} (etc.) satisfies:

$$cd \cos(\varphi_{AB}) = \frac{1}{2}(c^2 + d^2 - e^2)$$

The Areal Law of Sines

The **squared** sine of φ_{AB} (etc) is given by the Heron-like formula:

$$c^2 d^2 \sin(\varphi_{AB})^2 = \frac{1}{4}(c+d+e)(c+d-e)(c-d+e)(-c+d+e)$$

McConnell's Rigidity Theorem

The seven areas determine a non-degenerate tetrahedron up to isometry (proof: $cd \sin(\varphi_{AB}) = \|\mathbf{a} - \mathbf{b}\| t^2$ & $t^4 = T(a, \dots, g)$).



CONCEPT #3: THE AREAL LAWS OF SINES & COSINES; MCCONNELL'S RIGIDITY THEOREM

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



Dihedral Angle Bisectors

The plane bisecting the dihedral angle φ_{AB} divides \overline{CD} in the ratios shown, and similarly for the other dihedrals.

The Areal Law of Cosines

The cosine of φ_{AB} (etc.) satisfies:

$$cd \cos(\varphi_{AB}) = \frac{1}{2}(c^2 + d^2 - e^2)$$

The Areal Law of Sines

The **squared** sine of φ_{AB} (etc) is given by the Heron-like formula:

$$c^2 d^2 \sin(\varphi_{AB})^2 = \frac{1}{4}(c+d+e)(c+d-e)(c-d+e)(-c+d+e)$$

McConnell's Rigidity Theorem

The seven areas determine a non-degenerate tetrahedron up to isometry (proof: $cd \sin(\varphi_{AB}) = \|a - b\| t^2$ & $t^4 = T(a, \dots, g)$).



CONCEPT #3: THE AREAL LAWS OF SINES & COSINES; MCCONNELL'S RIGIDITY THEOREM

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



Dihedral Angle Bisectors

The plane bisecting the dihedral angle φ_{AB} divides \overline{CD} in the ratios shown, and similarly for the other dihedrals.

The Areal Law of Cosines

The cosine of φ_{AB} (etc.) satisfies:

$$cd \cos(\varphi_{AB}) = \frac{1}{2}(c^2 + d^2 - e^2)$$

The Areal Law of Sines

The **squared** sine of φ_{AB} (etc) is given by the Heron-like formula:

$$c^2 d^2 \sin(\varphi_{AB})^2 = \frac{1}{4}(c+d+e)(c+d-e)(c-d+e)(-c+d+e)$$

McConnell's Rigidity Theorem

The seven areas determine a non-degenerate tetrahedron up to isometry (proof: $cd \sin(\varphi_{AB}) = \|a - b\| t^2$ & $t^4 = T(a, \dots, g)$).



CONCEPT #3: THE AREAL LAWS OF SINES & COSINES; MCCONNELL'S RIGIDITY THEOREM

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



Dihedral Angle Bisectors

The plane bisecting the dihedral angle φ_{AB} divides \overline{CD} in the ratios shown, and similarly for the other dihedrals.

The Areal Law of Cosines

The cosine of φ_{AB} (etc.) satisfies:

$$cd \cos(\varphi_{AB}) = \frac{1}{2}(c^2 + d^2 - e^2)$$

The Areal Law of Sines

The **squared** sine of φ_{AB} (etc) is given by the Heron-like formula:

$$c^2 d^2 \sin(\varphi_{AB})^2 = \frac{1}{4}(c+d+e)(c+d-e)(c-d+e)(-c+d+e)$$

McConnell's Rigidity Theorem

The seven areas determine a non-degenerate tetrahedron up to isometry (proof: $cd \sin(\varphi_{AB}) = \|\mathbf{a} - \mathbf{b}\| t^2$ & $t^4 = T(a, \dots, g)$).

CONCEPT #4: THE NATURAL (ANALOGS OF THE HERON) PARAMETERS



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



The insphere “touches” the exterior faces at points \bar{J} , \bar{K} , \bar{L} , \bar{M} all at a distance r (the inradius) from its center \bar{I} . Thus the distances from each vertex to its three adjacent in-touch points are **equal**.

It follows that pairs of **contact triangles** sharing a common edge are congruent.

The **natural parameters** u, \dots, z of a tetrahedron are defined as twice the areas of these 6 congruent pairs (1 per edge).

Clearly the natural parameters determine the areas of the exterior faces (as seen on the left). It can be shown they also determine those of the interior, and hence the tetrahedron itself up to isometry.

CONCEPT #4: THE NATURAL (ANALOGS OF THE HERON) PARAMETERS



Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles


Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra

Orthocentric systems


Open questions



The insphere “touches” the exterior faces at points \bar{J} , \bar{K} , \bar{L} , \bar{M} all at a distance r (the inradius) from its center \bar{I} . Thus the distances from each vertex to its three adjacent in-touch points are **equal**.

It follows that pairs of **contact triangles** sharing a common edge are congruent.

The **natural parameters** u, \dots, z of a tetrahedron are defined as twice the areas of these 6 congruent pairs (1 per edge).



Clearly the natural parameters determine the areas of the exterior faces (as seen on the left). It can be shown they also determine those of the interior, and hence the tetrahedron itself up to isometry.

CONCEPT #4: THE NATURAL (ANALOGS OF THE HERON) PARAMETERS



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles



Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



The insphere “touches” the exterior faces at points \bar{J} , \bar{K} , \bar{L} , \bar{M} all at a distance r (the inradius) from its center \bar{I} . Thus the distances from each vertex to its three adjacent in-touch points are **equal**.

It follows that pairs of **contact triangles** sharing a common edge are congruent.

The **natural parameters** u, \dots, z of a tetrahedron are defined as twice the areas of these 6 congruent pairs (1 per edge).

Clearly the natural parameters determine the areas of the exterior faces (as seen on the left). It can be shown they also determine those of the interior, and hence the tetrahedron itself up to isometry.



FORMULAE FOR THE NATURAL PARAMETERS, AND THE INVERSE PARAMETERS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

With $s := a + b + c + d$ as twice the exterior surface area, we have:

$$u = r \|\mathbf{a} - \mathbf{b}\| \cot(\varphi_{AB}/2) = (c + d + e)(c + d - e) / (2s)$$

$$v = r \|\mathbf{a} - \mathbf{c}\| \cot(\varphi_{AC}/2) = (b + d + f)(b + d - f) / (2s)$$

$$w = r \|\mathbf{a} - \mathbf{d}\| \cot(\varphi_{AD}/2) = (b + c + g)(b + c - g) / (2s)$$

$$x = r \|\mathbf{b} - \mathbf{c}\| \cot(\varphi_{BC}/2) = (a + d + g)(a + d - g) / (2s)$$

$$y = r \|\mathbf{b} - \mathbf{d}\| \cot(\varphi_{BD}/2) = (a + c + f)(a + c - f) / (2s)$$

$$z = r \|\mathbf{c} - \mathbf{d}\| \cot(\varphi_{CD}/2) = (a + b + e)(a + b - e) / (2s)$$

We also **define** the corresponding "inverse" parameters as:

$$\tilde{u} = r \|\mathbf{a} - \mathbf{b}\| \tan(\varphi_{AB}/2) = (e + d - c)(e - d + c) / (2s)$$

$$\tilde{v} = r \|\mathbf{a} - \mathbf{c}\| \tan(\varphi_{AC}/2) = (f + d - b)(f - d + b) / (2s)$$

$$\tilde{w} = r \|\mathbf{a} - \mathbf{d}\| \tan(\varphi_{AD}/2) = (g + c - b)(g - c + b) / (2s)$$

$$\tilde{x} = r \|\mathbf{b} - \mathbf{c}\| \tan(\varphi_{BC}/2) = (g + d - a)(g - d + a) / (2s)$$

$$\tilde{y} = r \|\mathbf{b} - \mathbf{d}\| \tan(\varphi_{BD}/2) = (f + c - a)(f - c + a) / (2s)$$

$$\tilde{z} = r \|\mathbf{c} - \mathbf{d}\| \tan(\varphi_{CD}/2) = (e + b - a)(e - b + a) / (2s)$$



FORMULAE FOR THE NATURAL PARAMETERS, AND THE INVERSE PARAMETERS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

With $s := a + b + c + d$ as twice the exterior surface area, we have:

$$u = r \|\mathbf{a} - \mathbf{b}\| \cot(\varphi_{AB}/2) = (c + d + e)(c + d - e) / (2s)$$

$$v = r \|\mathbf{a} - \mathbf{c}\| \cot(\varphi_{AC}/2) = (b + d + f)(b + d - f) / (2s)$$

$$w = r \|\mathbf{a} - \mathbf{d}\| \cot(\varphi_{AD}/2) = (b + c + g)(b + c - g) / (2s)$$

$$x = r \|\mathbf{b} - \mathbf{c}\| \cot(\varphi_{BC}/2) = (a + d + g)(a + d - g) / (2s)$$

$$y = r \|\mathbf{b} - \mathbf{d}\| \cot(\varphi_{BD}/2) = (a + c + f)(a + c - f) / (2s)$$

$$z = r \|\mathbf{c} - \mathbf{d}\| \cot(\varphi_{CD}/2) = (a + b + e)(a + b - e) / (2s)$$

We also **define** the corresponding “inverse” parameters as:

$$\tilde{u} = r \|\mathbf{a} - \mathbf{b}\| \tan(\varphi_{AB}/2) = (e + d - c)(e - d + c) / (2s)$$

$$\tilde{v} = r \|\mathbf{a} - \mathbf{c}\| \tan(\varphi_{AC}/2) = (f + d - b)(f - d + b) / (2s)$$

$$\tilde{w} = r \|\mathbf{a} - \mathbf{d}\| \tan(\varphi_{AD}/2) = (g + c - b)(g - c + b) / (2s)$$

$$\tilde{x} = r \|\mathbf{b} - \mathbf{c}\| \tan(\varphi_{BC}/2) = (g + d - a)(g - d + a) / (2s)$$

$$\tilde{y} = r \|\mathbf{b} - \mathbf{d}\| \tan(\varphi_{BD}/2) = (f + c - a)(f - c + a) / (2s)$$

$$\tilde{z} = r \|\mathbf{c} - \mathbf{d}\| \tan(\varphi_{CD}/2) = (e + b - a)(e - b + a) / (2s)$$



HERON'S FORMULA FOR TETRAHEDRA

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

The inverse parameters are related to the areas of the triangles into which the exterior faces are divided by their “ex-touch” points. They may be expressed in terms of the natural parameters as:

$$\begin{aligned}\tilde{u} &= 2((v+x)(w+y) - uz)/s, & \tilde{z} &= 2((v+w)(x+y) - uz)/s \\ \tilde{v} &= 2((u+x)(w+z) - vy)/s, & \tilde{y} &= 2((u+w)(x+z) - vy)/s \\ \tilde{w} &= 2((u+y)(v+z) - wx)/s, & \tilde{x} &= 2((u+v)(y+z) - wx)/s\end{aligned}$$

Theorem:

The volume $t := 6|\overline{ABCD}|$ & inradius $r = t/s$ of a tetrahedron are given in terms of the natural parameters and $s = 2(u + \dots + z)$ by

$$t^4 = s^2 \Omega \quad \& \quad r^4 = \Omega/s^2, \quad \text{wherein}$$

$$\Omega = \Omega(u, v, w, x, y, z) := -\det \begin{bmatrix} 0 & u & v & w \\ u & 0 & x & y \\ v & x & 0 & z \\ w & y & z & 0 \end{bmatrix}.$$



HERON'S FORMULA FOR TETRAHEDRA

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

The inverse parameters are related to the areas of the triangles into which the exterior faces are divided by their "ex-touch" points.

They may be expressed in terms of the natural parameters as:

$$\tilde{u} = 2((v+x)(w+y) - uz)/s, \quad \tilde{z} = 2((v+w)(x+y) - uz)/s$$

$$\tilde{v} = 2((u+x)(w+z) - vy)/s, \quad \tilde{y} = 2((u+w)(x+z) - vy)/s$$

$$\tilde{w} = 2((u+y)(v+z) - wx)/s, \quad \tilde{x} = 2((u+v)(y+z) - wx)/s$$

Theorem:

The volume $t := 6|\overline{ABCD}|$ & inradius $r = t/s$ of a tetrahedron are given in terms of the natural parameters and $s = 2(u + \dots + z)$ by

$$t^4 = s^2 \Omega \quad \& \quad r^4 = \Omega/s^2, \quad \text{wherein}$$

$$\Omega = \Omega(u, v, w, x, y, z) := -\det \begin{bmatrix} 0 & u & v & w \\ u & 0 & x & y \\ v & x & 0 & z \\ w & y & z & 0 \end{bmatrix}.$$



PROOFS OF THE THEOREM, THE IN-TOUCH TETRAHEDRON, AND THE AREAL VECTORS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



The theorem can be proven by either:

- 1 substituting $D_{AB} \leftarrow u\tilde{u}/r^2$ etc. in the 4-point Cayley-Menger determinant $\Delta_D[A, B, C, D]$ and simplifying (a lot)
- 2 expressing the **in-touch tetrahedron** \overline{JKLM} volume in terms of the natural parameters & using the relation:

$$abcd \mathbf{n}_\infty \wedge \mathbf{j} \wedge \mathbf{k} \wedge \mathbf{l} \wedge \mathbf{m} = -\Omega(u, v, w, x, y, z) \mathbf{n}_\infty \wedge \mathbf{a} \wedge \mathbf{b} \wedge \mathbf{c} \wedge \mathbf{d}$$

(which follows as in 2D from the barycentric representations of $\mathbf{j}, \mathbf{k}, \mathbf{l}, \mathbf{m}$ together with the determinant multiplication theorem).

Yet another proof can be given based on the fact that the determinant of the Gram matrix of the **areal vectors** (facet normals weighted by the areas, as above) of any 3 facets also equals t^4 .



PROOFS OF THE THEOREM, THE IN-TOUCH TETRAHEDRON, AND THE AREAL VECTORS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



The theorem can be proven by either:

- 1 substituting $D_{AB} \leftarrow u\tilde{u}/r^2$ etc. in the 4-point Cayley-Menger determinant $\Delta_D[A, B, C, D]$ and simplifying (a lot)
- 2 expressing the **in-touch tetrahedron** \overline{JKLM} volume in terms of the natural parameters & using the relation:

$$abcd \mathbf{n}_\infty \wedge \mathbf{j} \wedge \mathbf{k} \wedge \mathbf{l} \wedge \mathbf{m} = -\Omega(u, v, w, x, y, z) \mathbf{n}_\infty \wedge \mathbf{a} \wedge \mathbf{b} \wedge \mathbf{c} \wedge \mathbf{d}$$

(which follows as in 2D from the barycentric representations of $\mathbf{j}, \mathbf{k}, \mathbf{l}, \mathbf{m}$ together with the determinant multiplication theorem).

Yet another proof can be given based on the fact that the determinant of the Gram matrix of the **areal vectors** (facet normals weighted by the areas, as above) of any 3 facets also equals t^4 .



PROOFS OF THE THEOREM, THE IN-TOUCH TETRAHEDRON, AND THE AREAL VECTORS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



The theorem can be proven by either:

- 1 substituting $D_{AB} \leftarrow u\tilde{u}/r^2$ etc. in the 4-point Cayley-Menger determinant $\Delta_D[A, B, C, D]$ and simplifying (a lot)
- 2 expressing the **in-touch tetrahedron** \overline{JKLM} volume in terms of the natural parameters & using the relation:

$$abcd \mathbf{n}_\infty \wedge \mathbf{j} \wedge \mathbf{k} \wedge \mathbf{l} \wedge \mathbf{m} = -\Omega(u, v, w, x, y, z) \mathbf{n}_\infty \wedge \mathbf{a} \wedge \mathbf{b} \wedge \mathbf{c} \wedge \mathbf{d}$$

(which follows as in 2D from the barycentric representations of $\mathbf{j}, \mathbf{k}, \mathbf{l}, \mathbf{m}$ together with the determinant multiplication theorem).

Yet another proof can be given based on the fact that the determinant of the Gram matrix of the **areal vectors** (facet normals weighted by the areas, as above) of any 3 facets also equals t^4 .



PROOFS OF THE THEOREM, THE IN-TOUCH TETRAHEDRON, AND THE AREAL VECTORS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



Yet another proof can be given based on the fact that the determinant of the Gram matrix of the **areal vectors** (facet normals weighted by the areas, as above) of any 3 facets also equals t^4 .



TABLE OF CONTENTS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- 1 Low-dimensional Euclidean geometry is not so elementary
- 2 Heron's formula and its relation to incenter geometry
- 3 Extending Heron's formula to tetrahedra (and beyond ...)
- 4 But its zeros are **really** weird in 3 (& higher?) dimensions
- 5 Orthocentric tetrahedra as extensions of triangles to 3D
- 6 Orthocentric systems in 2 & 3D, and in/excenter geometry
- 7 Open Questions and Future Work



THE PROJECTIVE NATURE OF THE ZEROS (AND GEOMETRIC INSIGHTS THAT FOLLOW)

So I will now address the question everyone is probably asking:

Why should I care about yet-another way of computing tetrahedron volumes (and a convoluted one at that)?

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



THE PROJECTIVE NATURE OF THE ZEROS (AND GEOMETRIC INSIGHTS THAT FOLLOW)

So I will now address the question everyone is probably asking:

Why should I care about yet-another way of computing tetrahedron volumes (and a convoluted one at that)?

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions




THE PROJECTIVE NATURE OF THE ZEROS (AND GEOMETRIC INSIGHTS THAT FOLLOW)

So I will now address the question everyone is probably asking:

Why should I care about yet-another way of computing tetrahedron volumes (and a convoluted one at that)?

In a non-degenerate tetrahedron the facet normals span three dimensions.



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions




THE PROJECTIVE NATURE OF THE ZEROS (AND GEOMETRIC INSIGHTS THAT FOLLOW)

So I will now address the question everyone is probably asking:

Why should I care about yet-another way of computing tetrahedron volumes (and a convoluted one at that)?

In a non-degenerate tetrahedron the facet normals span three dimensions.

When it's flattened (e.g. projected) into a plane, they become collinear.



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions






THE PROJECTIVE NATURE OF THE ZEROS (AND GEOMETRIC INSIGHTS THAT FOLLOW)

So I will now address the question everyone is probably asking:

Why should I care about yet-another way of computing tetrahedron volumes (and a convoluted one at that)?

In a non-degenerate tetrahedron the facet normals span three dimensions.

When it's flattened (e.g. projected) into a plane, they become collinear.



But if it's "squeezed" onto a line they get coplanar: →

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions




THE ZEROS AS LIMITS OF AFFINE SQUEEZE (AND STRETCH) TRANSFORMATIONS

Analytically, the zeros of Ω are the limits of a sequence of affine transformations of the form

$$\mathcal{A}_\sigma := \begin{bmatrix} \sigma^{-1} & 0 & 0 \\ 0 & \sigma^{-1} & 0 \\ 0 & 0 & \sigma \end{bmatrix}$$

applied to generic tetrahedra in space as $\sigma \rightarrow \infty$.

Areas & N.P.s vs. squeeze factor σ



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions




THE ZEROS AS LIMITS OF AFFINE SQUEEZE (AND STRETCH) TRANSFORMATIONS

Analytically, the zeros of Ω are the limits of a sequence of affine transformations of the form

$$\mathcal{A}_\sigma := \begin{bmatrix} \sigma^{-1} & 0 & 0 \\ 0 & \sigma^{-1} & 0 \\ 0 & 0 & \sigma \end{bmatrix}$$

applied to generic tetrahedra in space as $\sigma \rightarrow \infty$.

Areas & N.P.s vs. squeeze factor σ





THESE ZEROS CONSTITUTE A PROFOUND DIFFERENCE BETWEEN 2 & 3 DIMENSIONS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- Triangles with any given area and vertices at infinity also exist, but two of their Heron parameters become infinite.
- In contrast, tetrahedra with **finite** natural parameters and with $\Omega = 0$ do **not** correspond to quadruples of points in the Euclidean plane (in any obvious way), even though the set of planar quadruples also has $4 \cdot 2 - 3 = 5$ degrees of freedom.
- The **ratios** of their squared distances $u\tilde{u}/v\tilde{v}$, \dots , $y\tilde{y}/z\tilde{z}$ are generically finite, but the zeros of Ω are **not** quadruples on a line in the projective completion of Euclidean space, because such lines have only **one** point at infinity.
- These zeros are clearly non-physical, but they are perfectly well-defined mathematically and full of geometric structure.
- And it seems no one's **ever** before noticed that such a novel "completion" of the Euclidean symmetric product $\mathbb{E}^{3 \otimes 4}$ exists!



THESE ZEROS CONSTITUTE A PROFOUND DIFFERENCE BETWEEN 2 & 3 DIMENSIONS

- Triangles with any given area and vertices at infinity also exist, but two of their Heron parameters become infinite.
- In contrast, tetrahedra with **finite** natural parameters and with $\Omega = 0$ do **not** correspond to quadruples of points in the Euclidean plane (in any obvious way), even though the set of planar quadruples also has $4 \cdot 2 - 3 = 5$ degrees of freedom.
- The **ratios** of their squared distances $u\tilde{u}/v\tilde{v}, \dots, y\tilde{y}/z\tilde{z}$ are generically finite, but the zeros of Ω are **not** quadruples on a line in the projective completion of Euclidean space, because such lines have only **one** point at infinity.
- These zeros are clearly non-physical, but they are perfectly well-defined mathematically and full of geometric structure.
- And it seems no one's **ever** before noticed that such a novel "completion" of the Euclidean symmetric product $\mathbb{E}^{3 \otimes 4}$ exists!

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



THESE ZEROS CONSTITUTE A PROFOUND DIFFERENCE BETWEEN 2 & 3 DIMENSIONS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- Triangles with any given area and vertices at infinity also exist, but two of their Heron parameters become infinite.
- In contrast, tetrahedra with **finite** natural parameters and with $\Omega = 0$ do **not** correspond to quadruples of points in the Euclidean plane (in any obvious way), even though the set of planar quadruples also has $4 \cdot 2 - 3 = 5$ degrees of freedom.
- The **ratios** of their squared distances $u\tilde{u}/v\tilde{v}$, \dots , $y\tilde{y}/z\tilde{z}$ are generically finite, but the zeros of Ω are **not** quadruples on a line in the projective completion of Euclidean space, because such lines have only **one** point at infinity.
- These zeros are clearly non-physical, but they are perfectly well-defined mathematically and full of geometric structure.
- And it seems no one's **ever** before noticed that such a novel "completion" of the Euclidean symmetric product $\mathbb{E}^{3 \otimes 4}$ exists!



THESE ZEROS CONSTITUTE A PROFOUND DIFFERENCE BETWEEN 2 & 3 DIMENSIONS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- Triangles with any given area and vertices at infinity also exist, but two of their Heron parameters become infinite.
- In contrast, tetrahedra with **finite** natural parameters and with $\Omega = 0$ do **not** correspond to quadruples of points in the Euclidean plane (in any obvious way), even though the set of planar quadruples also has $4 \cdot 2 - 3 = 5$ degrees of freedom.
- The **ratios** of their squared distances $u\tilde{u}/v\tilde{v}$, \dots , $y\tilde{y}/z\tilde{z}$ are generically finite, but the zeros of Ω are **not** quadruples on a line in the projective completion of Euclidean space, because such lines have only **one** point at infinity.
- These zeros are clearly non-physical, but they are perfectly well-defined mathematically and full of geometric structure.
- And it seems no one's **ever** before noticed that such a novel "completion" of the Euclidean symmetric product $\mathbb{E}^{3 \otimes 4}$ exists!



THESE ZEROS CONSTITUTE A PROFOUND DIFFERENCE BETWEEN 2 & 3 DIMENSIONS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems


Open
questions

- Triangles with any given area and vertices at infinity also exist, but two of their Heron parameters become infinite.
- In contrast, tetrahedra with **finite** natural parameters and with $\Omega = 0$ do **not** correspond to quadruples of points in the Euclidean plane (in any obvious way), even though the set of planar quadruples also has $4 \cdot 2 - 3 = 5$ degrees of freedom.
- The **ratios** of their squared distances $u\tilde{u}/v\tilde{v}, \dots, y\tilde{y}/z\tilde{z}$ are generically finite, but the zeros of Ω are **not** quadruples on a line in the projective completion of Euclidean space, because such lines have only **one** point at infinity.
- These zeros are clearly non-physical, but they are perfectly well-defined mathematically and full of geometric structure.
- And it seems no one's **ever** before noticed that such a novel "completion" of the Euclidean symmetric product $\mathbb{E}^{3 \otimes 4}$ exists!



THE STRUCTURE OF THE 5D SET OF ZEROS AS A COMBINATORIAL LATTICE (PO-SET)

Hasse diagram of lattice structure induced by $\Omega = 0$ and all independent combinations of complementary products $u\tilde{u}, v\tilde{v}$ etc. vanishing (ranks on left are dimension of span of facet normals).



Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra


Orthocentric systems

Open questions



THE STRUCTURE OF THE 5D SET OF ZEROS AS A COMBINATORIAL LATTICE (PO-SET)

Hasse diagram of lattice structure induced by $\Omega = 0$ and all independent combinations of complementary products $u\tilde{u}, v\tilde{v}$ etc. vanishing (ranks on left are dimension of span of facet normals).



Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra


Orthocentric systems

Open questions



RANK 1 ZEROS CORRESPOND TO 4-POINT CONFIGURATIONS IN EQUI-AFFINE PLANE)

Plane divided into 16 regions by lines along sides of triangle \overline{ABC} and their parallels thru its vertices (labeled by pool ball icons ♫).



Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra


Orthocentric systems

Open questions



RANK 1 ZEROS CORRESPOND TO 4-POINT CONFIGURATIONS IN EQUI-AFFINE PLANE)

Plane divided into 16 regions by lines along sides of triangle \overline{ABC} and their parallels thru its vertices (labeled by pool ball icons ♫).



Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra


Orthocentric systems

Open questions



RANK 1 ZEROS CORRESPOND TO 4-POINT CONFIGURATIONS IN EQUI-AFFINE PLANE)

Plane divided into 16 regions by lines along sides of triangle \overline{ABC} and their parallels thru its vertices (labeled by pool ball icons ♫).



Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra


Orthocentric systems

Open questions



RANK 1 ZEROS CORRESPOND TO 4-POINT CONFIGURATIONS IN EQUI-AFFINE PLANE)

Plane divided into 16 regions by lines along sides of triangle \overline{ABC} and their parallels thru its vertices (labeled by pool ball icons ♫).




Dotted lines thru the vertices divide 3 of those regions into 4 subregions each, distinguished by which tetrahedron inequalities saturate, e.g. $\mathcal{T}_{AB}^{(2)} \equiv e + d = c$, $\mathcal{T}_{AB}^{(3)} \equiv e + c = d$, etc.

- Heron's Formula in Orthocentric Tetrahedra
- Havel & Sobczyk
- Not-so elementary
- Heron's formula for triangles
- Heron's formula for tetrahedra
- The zeros' projective nature
- Orthocentric tetrahedra
- Orthocentric systems
- Open questions



RANK 1 ZEROS CORRESPOND TO 4-POINT CONFIGURATIONS IN EQUI-AFFINE PLANE)

Plane divided into 16 regions by lines along sides of triangle \overline{ABC} and their parallels thru its vertices (labeled by pool ball icons ♫).



Dotted lines thru the vertices divide 3 of those regions into 4 subregions each, distinguished by which tetrahedron inequalities saturate, e.g. $\mathcal{T}_{AB}^{(2)} \equiv e + d = c$, $\mathcal{T}_{AB}^{(3)} \equiv e + c = d$, etc.

This classification of affine point configurations was studied intensively in the late 20th century by Goodman & Pollack, without knowledge of the tetrahedron inequalities.

Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra

Orthocentric systems

Open questions



TABLE OF CONTENTS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- 1 Low-dimensional Euclidean geometry is not so elementary
- 2 Heron's formula and its relation to incenter geometry
- 3 Extending Heron's formula to tetrahedra (and beyond ...)
- 4 But its zeros are **really** weird in 3 (& higher?) dimensions
- 5 Orthocentric tetrahedra as extensions of triangles to 3D**
- 6 Orthocentric systems in 2 & 3D, and in/excenter geometry
- 7 Open Questions and Future Work



WHAT EUCLID MISSED, GAUSS GOT RIGHT, AND THE TETRAHEDRON LEFT BEHIND

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



The concurrence of the altitudes of a triangle is curiously missing from Euclid's Elements.

Gauss proved it by noting that the orthocenter \bar{H} is the circumcenter \bar{O}_{DEF} of the "anti-medial" triangle \bar{DEF} .

In higher dimensions, however, the altitudes of simplices are **not** generally concurrent!



WHAT EUCLID MISSED, GAUSS GOT RIGHT, AND THE TETRAHEDRON LEFT BEHIND

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



The concurrence of the altitudes of a triangle is curiously missing from Euclid's Elements.

Gauss proved it by noting that the orthocenter \bar{H} is the circumcenter \bar{O}_{DEF} of the "anti-medial" triangle \bar{DEF} .

In higher dimensions, however, the altitudes of simplices are **not** generally concurrent!



WHAT EUCLID MISSED, GAUSS GOT RIGHT, AND THE TETRAHEDRON LEFT BEHIND

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



The concurrence of the altitudes of a triangle is curiously missing from Euclid's Elements.


Gauss proved it by noting that the orthocenter \bar{H} is the circumcenter \bar{O}_{DEF} of the "anti-medial" triangle \bar{DEF} .

In higher dimensions, however, the altitudes of simplices are **not** generally concurrent!



THE MONGE POINT AND THE MEDIAL OCTAHEDRON'S FACIAL ORTHOCENTERS

A tetrahedron's **Monge point** \bar{M} is where the perpendiculars thru the orthocenters \bar{H}_D etc. of its medial octahedron faces inside the tetrahedron meet.



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



THE MONGE POINT AND THE MEDIAL OCTAHEDRON'S FACIAL ORTHOCENTERS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature


Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

A tetrahedron's **Monge point** \bar{M} is where the perpendiculars thru the orthocenters \bar{H}_D etc. of its medial octahedron faces inside the tetrahedron meet.

When the altitudes likewise concur, they do so at \bar{M} , which is also the reflection of the circumcenter \bar{O} in the centroid \bar{G} , and they're all on the Euler line (green).





THE MONGE POINT AND THE MEDIAL OCTAHEDRON'S FACIAL ORTHOCENTERS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra


Orthocentric
systems

Open
questions

A tetrahedron's **Monge point** \bar{M} is where the perpendiculars thru the orthocenters \bar{H}_D etc. of its medial octahedron faces inside the tetrahedron meet.

When the altitudes likewise concur, they do so at \bar{M} , which is also the reflection of the circumcenter \bar{O} in the centroid \bar{G} , and they're all on the Euler line (green).

The perpendiculars thru the orthocenters of the octahedron's "surface" faces, or circumcenters \bar{O}_D etc. of \overline{ABCD} 's facets, meet at \bar{O} (blue lines).





ORTHOCENTRIC TETRAHEDRA: THE TRUE GENERALIZATION OF TRIANGLES TO 3D?

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

*Orthocentric Simplices as the True
Generalizations of Triangles*

Mowaffaq Hajja & Horst Martini

The Mathematical Intelligencer
2013



Simplices where the altitudes do concur are termed **orthocentric**, and behave more like triangles, e.g. they are equilateral iff the incenter and centroid coincide.

In particular, a tetrahedron is orthocentric if & only if either:

- $(b - a) \cdot (d - c) = (c - a) \cdot (d - b) = (c - b) \cdot (d - a) = 0$
- $\|b - a\|^2 + \|d - c\|^2 = \|c - a\|^2 + \|d - b\|^2 = \|c - b\|^2 + \|d - a\|^2$
- there exist $D_A, D_B, D_C, D_D \in \mathbb{R}$ such that their pairwise sums equal the squared inter-vertex distances, i.e.

$$\|b - a\|^2 = D_A + D_B, \dots, \|d - c\|^2 = D_C + D_D$$

Note that triangles always satisfy this last condition with

$$D_A := \frac{1}{2}(-a^2 + b^2 + c^2), D_B := \frac{1}{2}(a^2 - b^2 + c^2), D_C := \frac{1}{2}(a^2 + b^2 - c^2).$$



ORTHOCENTRIC TETRAHEDRA: THE TRUE GENERALIZATION OF TRIANGLES TO 3D?

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

*Orthocentric Simplices as the True
Generalizations of Triangles*

Mowaffaq Hajja & Horst Martini

The Mathematical Intelligencer
2013



Simplices where the altitudes do concur are termed **orthocentric**, and behave more like triangles, e.g. they are equilateral iff the incenter and centroid coincide.

In particular, a tetrahedron is orthocentric if & only if either:

- $(\mathbf{b} - \mathbf{a}) \cdot (\mathbf{d} - \mathbf{c}) = (\mathbf{c} - \mathbf{a}) \cdot (\mathbf{d} - \mathbf{b}) = (\mathbf{c} - \mathbf{b}) \cdot (\mathbf{d} - \mathbf{a}) = 0$
- $\|\mathbf{b} - \mathbf{a}\|^2 + \|\mathbf{d} - \mathbf{c}\|^2 = \|\mathbf{c} - \mathbf{a}\|^2 + \|\mathbf{d} - \mathbf{b}\|^2 = \|\mathbf{c} - \mathbf{b}\|^2 + \|\mathbf{d} - \mathbf{a}\|^2$
- there exist $D_A, D_B, D_C, D_D \in \mathbb{R}$ such that their pairwise sums equal the squared inter-vertex distances, i.e.

$$\|\mathbf{b} - \mathbf{a}\|^2 = D_A + D_B, \dots, \|\mathbf{d} - \mathbf{c}\|^2 = D_C + D_D$$

Note that triangles always satisfy this last condition with

$$D_A := \frac{1}{2}(-a^2 + b^2 + c^2), D_B := \frac{1}{2}(a^2 - b^2 + c^2), D_C := \frac{1}{2}(a^2 + b^2 - c^2).$$



ORTHOCENTRIC TETRAHEDRA: THE TRUE GENERALIZATION OF TRIANGLES TO 3D?

Orthocentric Simplices as the True Generalizations of Triangles

Mowaffaq Hajja & Horst Martini

The Mathematical Intelligencer
2013



Simplices where the altitudes do concur are termed **orthocentric**, and behave more like triangles, e.g. they are equilateral iff the incenter and centroid coincide.

In particular, a tetrahedron is orthocentric if & only if either:

- $(\mathbf{b} - \mathbf{a}) \cdot (\mathbf{d} - \mathbf{c}) = (\mathbf{c} - \mathbf{a}) \cdot (\mathbf{d} - \mathbf{b}) = (\mathbf{c} - \mathbf{b}) \cdot (\mathbf{d} - \mathbf{a}) = 0$
- $\|\mathbf{b} - \mathbf{a}\|^2 + \|\mathbf{d} - \mathbf{c}\|^2 = \|\mathbf{c} - \mathbf{a}\|^2 + \|\mathbf{d} - \mathbf{b}\|^2 = \|\mathbf{c} - \mathbf{b}\|^2 + \|\mathbf{d} - \mathbf{a}\|^2$
- there exist $D_A, D_B, D_C, D_D \in \mathbb{R}$ such that their pairwise sums equal the squared inter-vertex distances, i.e.

$$\|\mathbf{b} - \mathbf{a}\|^2 = D_A + D_B, \dots, \|\mathbf{d} - \mathbf{c}\|^2 = D_C + D_D$$

Note that triangles always satisfy this last condition with

$$D_A := \frac{1}{2}(-a^2 + b^2 + c^2), \quad D_B := \frac{1}{2}(a^2 - b^2 + c^2), \quad D_C := \frac{1}{2}(a^2 + b^2 - c^2).$$



THE AMAZINGLY SIMPLE DISTANCE GEOMETRY OF ORTHOCENTRIC TETRAHEDRA

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

The Cayley-Menger determinants and other **squared** distance-area-volume relations simplify amazingly in the orthocentric case:

- $a^2 = D_B D_C + D_B D_D + D_C D_D$ (and similarly b^2, c^2, d^2)
- $e^2 = (D_A + D_B)(D_C + D_D)$ (and similarly f^2, g^2)
- $(a^2 + b^2 - e^2)/2 = D_C D_D$ (and similarly for the rest)
- $t^2 = D_A D_B D_C + D_A D_B D_D + D_A D_C D_D + D_B D_C D_D$
- $R^2 = \left(a^2 D_A^2 + b^2 D_B^2 + c^2 D_C^2 + d^2 D_D^2 \right) / (4t^2)$ (circumradius)

Unfortunately we **cannot** get the squared inradius $r^2 = t^2/s^2$ (or the natural parameters) from the D 's without taking square roots; the exterior surface area $s = a + b + c + d$ involves four of those!

It **is** possible to go the other way, e.g. $D_A = (u\tilde{u} + v\tilde{v} - x\tilde{x})/(2r^2)$; moreover, a tetrahedron is orthocentric if & only if

$$u\tilde{u} + z\tilde{z} = v\tilde{v} + y\tilde{y} = w\tilde{w} + x\tilde{x} \quad (= (D_A + D_B + D_C + D_D)r^2).$$



THE AMAZINGLY SIMPLE DISTANCE GEOMETRY OF ORTHOCENTRIC TETRAHEDRA

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

The Cayley-Menger determinants and other **squared** distance-area-volume relations simplify amazingly in the orthocentric case:

- $a^2 = D_B D_C + D_B D_D + D_C D_D$ (and similarly b^2, c^2, d^2)
- $e^2 = (D_A + D_B)(D_C + D_D)$ (and similarly f^2, g^2)
- $(a^2 + b^2 - e^2)/2 = D_C D_D$ (and similarly for the rest)
- $t^2 = D_A D_B D_C + D_A D_B D_D + D_A D_C D_D + D_B D_C D_D$
- $R^2 = \left(a^2 D_A^2 + b^2 D_B^2 + c^2 D_C^2 + d^2 D_D^2 \right) / (4t^2)$ (circumradius)

Unfortunately we **cannot** get the squared inradius $r^2 = t^2/s^2$ (or the natural parameters) from the D 's without taking square roots; the exterior surface area $s = a + b + c + d$ involves four of those!

It **is** possible to go the other way, e.g. $D_A = (u\tilde{u} + v\tilde{v} - x\tilde{x})/(2r^2)$; moreover, a tetrahedron is orthocentric if & only if

$$u\tilde{u} + z\tilde{z} = v\tilde{v} + y\tilde{y} = w\tilde{w} + x\tilde{x} \quad (= (D_A + D_B + D_C + D_D)r^2).$$



THE AMAZINGLY SIMPLE DISTANCE GEOMETRY OF ORTHOCENTRIC TETRAHEDRA

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

The Cayley-Menger determinants and other **squared** distance-area-volume relations simplify amazingly in the orthocentric case:

- $a^2 = D_B D_C + D_B D_D + D_C D_D$ (and similarly b^2, c^2, d^2)
- $e^2 = (D_A + D_B)(D_C + D_D)$ (and similarly f^2, g^2)
- $(a^2 + b^2 - e^2)/2 = D_C D_D$ (and similarly for the rest)
- $t^2 = D_A D_B D_C + D_A D_B D_D + D_A D_C D_D + D_B D_C D_D$
- $R^2 = \left(a^2 D_A^2 + b^2 D_B^2 + c^2 D_C^2 + d^2 D_D^2 \right) / (4t^2)$ (circumradius)

Unfortunately we **cannot** get the squared inradius $r^2 = t^2/s^2$ (or the natural parameters) from the D 's without taking square roots; the exterior surface area $s = a + b + c + d$ involves four of those!

It **is** possible to go the other way, e.g. $D_A = (u\tilde{u} + v\tilde{v} - x\tilde{x})/(2r^2)$; moreover, a tetrahedron is orthocentric if & only if

$$u\tilde{u} + z\tilde{z} = v\tilde{v} + y\tilde{y} = w\tilde{w} + x\tilde{x} \quad (= (D_A + D_B + D_C + D_D)r^2).$$



THE AMAZINGLY SIMPLE DISTANCE GEOMETRY OF ORTHOCENTRIC TETRAHEDRA

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

The Cayley-Menger determinants and other **squared** distance-area-volume relations simplify amazingly in the orthocentric case:

- $a^2 = D_B D_C + D_B D_D + D_C D_D$ (and similarly b^2, c^2, d^2)
- $e^2 = (D_A + D_B)(D_C + D_D)$ (and similarly f^2, g^2)
- $(a^2 + b^2 - e^2)/2 = D_C D_D$ (and similarly for the rest)
- $t^2 = D_A D_B D_C + D_A D_B D_D + D_A D_C D_D + D_B D_C D_D$
- $R^2 = \left(a^2 D_A^2 + b^2 D_B^2 + c^2 D_C^2 + d^2 D_D^2 \right) / (4t^2)$ (circumradius)

Unfortunately we **cannot** get the squared inradius $r^2 = t^2/s^2$ (or the natural parameters) from the D 's without taking square roots; the exterior surface area $s = a + b + c + d$ involves four of those!

It **is** possible to go the other way, e.g. $D_A = (u\tilde{u} + v\tilde{v} - x\tilde{x})/(2r^2)$; moreover, a tetrahedron is orthocentric if & only if

$$u\tilde{u} + z\tilde{z} = v\tilde{v} + y\tilde{y} = w\tilde{w} + x\tilde{x} \quad \left(= (D_A + D_B + D_C + D_D)r^2 \right).$$



THE AMAZINGLY SIMPLE DISTANCE GEOMETRY OF ORTHOCENTRIC TETRAHEDRA

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

The Cayley-Menger determinants and other **squared** distance-area-volume relations simplify amazingly in the orthocentric case:

- $a^2 = D_B D_C + D_B D_D + D_C D_D$ (and similarly b^2, c^2, d^2)
- $e^2 = (D_A + D_B)(D_C + D_D)$ (and similarly f^2, g^2)
- $(a^2 + b^2 - e^2)/2 = D_C D_D$ (and similarly for the rest)
- $t^2 = D_A D_B D_C + D_A D_B D_D + D_A D_C D_D + D_B D_C D_D$
- $R^2 = \left(a^2 D_A^2 + b^2 D_B^2 + c^2 D_C^2 + d^2 D_D^2 \right) / (4t^2)$ (circumradius)

Unfortunately we **cannot** get the squared inradius $r^2 = t^2/s^2$ (or the natural parameters) from the D 's without taking square roots; the exterior surface area $s = a + b + c + d$ involves four of those!

It **is** possible to go the other way, e.g. $D_A = (u\tilde{u} + v\tilde{v} - x\tilde{x})/(2r^2)$; moreover, a tetrahedron is orthocentric if & only if

$$u\tilde{u} + z\tilde{z} = v\tilde{v} + y\tilde{y} = w\tilde{w} + x\tilde{x} \quad \left(= (D_A + D_B + D_C + D_D)r^2 \right).$$



THE AMAZINGLY SIMPLE DISTANCE GEOMETRY OF ORTHOCENTRIC TETRAHEDRA

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

The Cayley-Menger determinants and other **squared** distance-area-volume relations simplify amazingly in the orthocentric case:

- $a^2 = D_B D_C + D_B D_D + D_C D_D$ (and similarly b^2, c^2, d^2)
- $e^2 = (D_A + D_B)(D_C + D_D)$ (and similarly f^2, g^2)
- $(a^2 + b^2 - e^2)/2 = D_C D_D$ (and similarly for the rest)
- $t^2 = D_A D_B D_C + D_A D_B D_D + D_A D_C D_D + D_B D_C D_D$
- $R^2 = \left(a^2 D_A^2 + b^2 D_B^2 + c^2 D_C^2 + d^2 D_D^2 \right) / (4t^2)$ (circumradius)

Unfortunately we **cannot** get the squared inradius $r^2 = t^2/s^2$ (or the natural parameters) from the D 's without taking square roots; the exterior surface area $s = a + b + c + d$ involves four of those!

It **is** possible to go the other way, e.g. $D_A = (u\tilde{u} + v\tilde{v} - x\tilde{x})/(2r^2)$; moreover, a tetrahedron is orthocentric if & only if

$$u\tilde{u} + z\tilde{z} = v\tilde{v} + y\tilde{y} = w\tilde{w} + x\tilde{x} \quad (= (D_A + D_B + D_C + D_D)r^2).$$



THE AMAZINGLY SIMPLE DISTANCE GEOMETRY OF ORTHOCENTRIC TETRAHEDRA

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

The Cayley-Menger determinants and other **squared** distance-area-volume relations simplify amazingly in the orthocentric case:

- $a^2 = D_B D_C + D_B D_D + D_C D_D$ (and similarly b^2, c^2, d^2)
- $e^2 = (D_A + D_B)(D_C + D_D)$ (and similarly f^2, g^2)
- $(a^2 + b^2 - e^2)/2 = D_C D_D$ (and similarly for the rest)
- $t^2 = D_A D_B D_C + D_A D_B D_D + D_A D_C D_D + D_B D_C D_D$
- $R^2 = \left(a^2 D_A^2 + b^2 D_B^2 + c^2 D_C^2 + d^2 D_D^2 \right) / (4t^2)$ (circumradius)

Unfortunately we **cannot** get the squared inradius $r^2 = t^2/s^2$ (or the natural parameters) from the D 's without taking square roots; the exterior surface area $s = a + b + c + d$ involves four of those!

It **is** possible to go the other way, e.g. $D_A = (u\tilde{u} + v\tilde{v} - x\tilde{x})/(2r^2)$; moreover, a tetrahedron is orthocentric if & only if

$$u\tilde{u} + z\tilde{z} = v\tilde{v} + y\tilde{y} = w\tilde{w} + x\tilde{x} \quad (= (D_A + D_B + D_C + D_D)r^2).$$



THE AMAZINGLY SIMPLE DISTANCE GEOMETRY OF ORTHOCENTRIC TETRAHEDRA

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

The Cayley-Menger determinants and other **squared** distance-area-volume relations simplify amazingly in the orthocentric case:

- $a^2 = D_B D_C + D_B D_D + D_C D_D$ (and similarly b^2, c^2, d^2)
- $e^2 = (D_A + D_B)(D_C + D_D)$ (and similarly f^2, g^2)
- $(a^2 + b^2 - e^2)/2 = D_C D_D$ (and similarly for the rest)
- $t^2 = D_A D_B D_C + D_A D_B D_D + D_A D_C D_D + D_B D_C D_D$
- $R^2 = \left(a^2 D_A^2 + b^2 D_B^2 + c^2 D_C^2 + d^2 D_D^2 \right) / (4t^2)$ (circumradius)

Unfortunately we **cannot** get the squared inradius $r^2 = t^2/s^2$ (or the natural parameters) from the D 's without taking square roots; the exterior surface area $s = a + b + c + d$ involves four of those!

It **is** possible to go the other way, e.g. $D_A = (u\tilde{u} + v\tilde{v} - x\tilde{x})/(2r^2)$; moreover, a tetrahedron is orthocentric if & only if

$$u\tilde{u} + z\tilde{z} = v\tilde{v} + y\tilde{y} = w\tilde{w} + x\tilde{x} \quad (= (D_A + D_B + D_C + D_D)r^2).$$



CAN ORTHOCENTRIC TETRAHEDRA HAVE FACETAL VECTORS WHICH SPAN A PLANE?

These latter conditions don't involve r^2 , so they apply to degenerate tetrahedra, i.e. to the zeros of Ω , which leads to the question:

Can the squared distances in orthocentric tetrahedra become infinite while all the areas & natural parameters stay finite, or do they also behave like triangles in that way?

A detailed analysis of the equations shows that while most of the rank 2 zeros of Ω are eliminated by the orthocentricity constraints, one kind is **not**; in the previous lattice diagram, they are (in red):

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



CAN ORTHOCENTRIC TETRAHEDRA HAVE FACETAL VECTORS WHICH SPAN A PLANE?

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

These latter conditions don't involve r^2 , so they apply to degenerate tetrahedra, i.e. to the zeros of Ω , which leads to the question:

Can the squared distances in orthocentric tetrahedra become infinite while all the areas & natural parameters stay finite, or do they also behave like triangles in that way?

A detailed analysis of the equations shows that while most of the rank 2 zeros of Ω are eliminated by the orthocentricity constraints, one kind is **not**; in the previous lattice diagram, they are (in red):



CAN ORTHOCENTRIC TETRAHEDRA HAVE FACETAL VECTORS WHICH SPAN A PLANE?

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

These latter conditions don't involve r^2 , so they apply to degenerate tetrahedra, i.e. to the zeros of Ω , which leads to the question:

Can the squared distances in orthocentric tetrahedra become infinite while all the areas & natural parameters stay finite, or do they also behave like triangles in that way?

A detailed analysis of the equations shows that while most of the rank 2 zeros of Ω are eliminated by the orthocentricity constraints, one kind is **not**; in the previous lattice diagram, they are (in red):



CAN ORTHOCENTRIC TETRAHEDRA HAVE FACETAL VECTORS WHICH SPAN A PLANE?

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

These latter conditions don't involve r^2 , so they apply to degenerate tetrahedra, i.e. to the zeros of Ω , which leads to the question:

Can the squared distances in orthocentric tetrahedra become infinite while all the areas & natural parameters stay finite, or do they also behave like triangles in that way?

A detailed analysis of the equations shows that while most of the rank 2 zeros of Ω are eliminated by the orthocentricity constraints, one kind is **not**; in the previous lattice diagram, they are (in red):






TABLE OF CONTENTS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- 1 Low-dimensional Euclidean geometry is not so elementary
- 2 Heron's formula and its relation to incenter geometry
- 3 Extending Heron's formula to tetrahedra (and beyond ...)
- 4 But its zeros are **really** weird in 3 (& higher?) dimensions
- 5 Orthocentric tetrahedra as extensions of triangles to 3D
- 6 Orthocentric systems in 2 & 3D, and in/excenter geometry
- 7 Open Questions and Future Work



ORTHOCENTRIC SYSTEMS & IN/EXCENTER GEOMETRY IN THE PLANE

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



ORTHOCENTRIC SYSTEMS & IN/EXCENTER GEOMETRY IN THE PLANE



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

If \bar{H} is the orthocenter of a triangle \overline{ABC} , then \bar{A} is the orthocenter of \overline{ABH} , and likewise for \bar{B} , \bar{C} .

A, B, C : Excenters


G, G' : Centroids

H, H' : Orthocenters

I, J, K, L : Incenter/touch

O, O' : Circumcenters

N' : 9-point Center



ORTHOCENTRIC SYSTEMS & IN/EXCENTER GEOMETRY IN THE PLANE



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature


Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

If \bar{H} is the orthocenter of a triangle \bar{ABC} , then \bar{A} is the orthocenter of \bar{ABH} , and likewise for \bar{B} , \bar{C} .

Thus \bar{ABC} & \bar{H} is called an **orthocentric system**.



H, H' : Orthocenters

I, J, K, L : Incenter/touch

O, O' : Circumcenters

N' : 9-point Center

ORTHOCENTRIC SYSTEMS & IN/EXCENTER GEOMETRY IN THE PLANE



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



ORTHOCENTRIC SYSTEMS & IN/EXCENTER GEOMETRY IN THE PLANE



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



ORTHOCENTRIC SYSTEMS & IN/EXCENTER GEOMETRY IN THE PLANE



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions





ORTHOCENTRIC SYSTEMS & IN/EXCENTER GEOMETRY IN THE PLANE

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions





THE SIMILARITY OF THE IN-TOUCH AND EXCENTRAL TRIANGLES

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra


The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

It turns out that the in-touch & excentral triangles are **similar**, with a scale factor of $2R/r$, so that the circumradius of $A'B'C'$ is $2R$.





THE SIMILARITY OF THE IN-TOUCH AND EXCENTRAL TRIANGLES

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions





THE SIMILARITY OF THE IN-TOUCH AND EXCENTRAL TRIANGLES

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions





THE SIMILARITY OF THE IN-TOUCH AND EXCENTRAL TRIANGLES

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



Given my earlier result that the in-touch triangle has an area of $t_{it} = t \Omega / (abc)$, this also shows that the area of the excentral triangle is $tr^2s / (4tR) = 2Rs$.

The inverse homothety yields a scaled-down copy of the orthocentric system having the in-touch triangle's altitudes' feet as its orthic triangle.

THE EQUALITY OF ALL FOUR CIRCUM-RADII IN AN ORTHOCENTRIC SYSTEM



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra


The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

It follows that the radii of the circum-circles of the four triangles **all** equal $2R$, and in fact their circumcenters form an isometric copy of the orthocentric system that **orthogonally bisects** its edges.



THE EQUALITY OF ALL FOUR CIRCUM-RADII IN AN ORTHOCENTRIC SYSTEM



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature


Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

It follows that the radii of the circum-circles of the four triangles **all** equal $2R$, and in fact their circumcenters form an isometric copy of the orthocentric system that **orthogonally bisects** its edges.

Curiously, these circumcenters are also collinear with the **incenter** and the **in-touch points** (dotted **orange** lines).



THE EQUALITY OF ALL FOUR CIRCUM-RADII IN AN ORTHOCENTRIC SYSTEM



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



Curiously, these circumcenters are also collinear with the incenter and the in-touch points (dotted orange lines).

Clearly this mapping also goes both ways, and so constitutes a kind of duality relation I've not seen called out before.



INVERSION IN THE INCIRCLE, THE POLAR CIRCLES, AND CONFORMAL DUALITY

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra


The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

Inversion in the **incircle** exchanges the **vertices** with the **edge midpoints** of the in-touch triangle (\leftrightarrow).





INVERSION IN THE INCIRCLE, THE POLAR CIRCLES, AND CONFORMAL DUALITY

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature


Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

Inversion in the **incircle** exchanges the **vertices** with the **edge midpoints** of the in-touch triangle (\leftrightarrow).

Inversion in the **polar circles** (dotted) centered on the **excenters** exchanges the **vertices** with the **incenter** and the other 2 **excenters** (\leftrightarrow).





INVERSION IN THE INCIRCLE, THE POLAR CIRCLES, AND CONFORMAL DUALITY

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature


Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

Inversion in the **polar circles** (dotted) centered on the **excenters** exchanges the **vertices** with the **incenter** and the other 2 **excenters** (\leftrightarrow).

Note these circles are orthogonal so these inversions commute.





INVERSION IN THE INCIRCLE, THE POLAR CIRCLES, AND CONFORMAL DUALITY

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions





INVERSION IN THE INCIRCLE, THE POLAR CIRCLES, AND CONFORMAL DUALITY

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles


Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions




Inversion in the “imaginary” circle of radius $2\sqrt{rR}$, meaning inversion (\leftrightarrow) in the actual circle and reflection (\leftrightarrow) in its center \bar{I} (in either order), interchanges all 3 vertices with their respective excenters.

This operation, which Hertrich-Jeromin, King & O'Hare dubbed **conformal duality**, also interchanges the reciprocal bases of \overline{ABC} & $\overline{A'B'C'}$ in $\mathcal{G}_{3,1}$ (dashed circumcircles).



BUT ALMOST NONE OF THIS GENERALIZES TO THREE (OR HIGHER) DIMENSIONS!

If one constructs the in- & excenters of a tetrahedron (green) one finds, even if the tetrahedron is orthocentric, that its in-touch (red) and excentral (blue) tetrahedra:



- Are **not** similar to one another;
- Are not orthocentric, so their altitudes are **not** concurrent;
- And that the vertices of \overline{ABCD} are **not** the altitudes' feet on the facets of $\overline{A'B'C'D'}$.

But the green edges **do** intersect the blue edges in the vertices of an octahedron (**not** the medial)

that bisect the dihedral angles and span diagonals meeting at \bar{I} .

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



BUT ALMOST NONE OF THIS GENERALIZES TO THREE (OR HIGHER) DIMENSIONS!

Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra


The zeros' projective nature

Orthocentric tetrahedra

Orthocentric systems

Open questions

If one constructs the in- & excenters of a tetrahedron (green) one finds, even if the tetrahedron is orthocentric, that its in-touch (red) and excentral (blue) tetrahedra:




- Are **not** similar to one another;
- Are not orthocentric, so their altitudes are **not** concurrent;
- And that the vertices of \overline{ABCD} are **not** the altitudes' feet on the facets of $\overline{A'B'C'D'}$.

But the green edges **do** intersect the blue edges in the vertices of an octahedron (**not** the medial) that bisect the dihedral angles and span diagonals meeting at \bar{I} .



BUT ALMOST NONE OF THIS GENERALIZES TO THREE (OR HIGHER) DIMENSIONS!

If one constructs the in- & excenters of a tetrahedron (green) one finds, even if the tetrahedron is orthocentric, that its in-touch (red) and excentral (blue) tetrahedra:



- Are **not** similar to one another;
- Are not orthocentric, so their altitudes are **not** concurrent;
- And that the vertices of \overline{ABCD} are **not** the altitudes' feet on the facets of $\overline{A'B'C'D'}$.

But the green edges **do** intersect the blue edges in the vertices of an octahedron (**not** the medial)


that bisect the dihedral angles and span diagonals meeting at \bar{I} .

Heron's Formula in Orthocentric Tetrahedra
 Havel & Sobczyk
 Not-so elementary
 Heron's formula for triangles
 Heron's formula for tetrahedra
 The zeros' projective nature
 Orthocentric tetrahedra
 Orthocentric systems
 Open questions



BUT ALMOST NONE OF THIS GENERALIZES TO THREE (OR HIGHER) DIMENSIONS!

If one constructs the in- & excenters of a tetrahedron (green) one finds, even if the tetrahedron is orthocentric, that its in-touch (red) and excentral (blue) tetrahedra:



- Are **not** similar to one another;
- Are not orthocentric, so their altitudes are **not** concurrent;
- And that the vertices of \overline{ABCD} are **not** the altitudes' feet on the facets of $\overline{A'B'C'D'}$.

But the green edges **do** intersect the blue edges in the vertices of an octahedron (**not** the medial) that bisect the dihedral angles and span diagonals meeting at \bar{I} .

Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra

Orthocentric systems

Open questions



BUT ALMOST NONE OF THIS GENERALIZES TO THREE (OR HIGHER) DIMENSIONS!

Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra


The zeros' projective nature

Orthocentric tetrahedra

Orthocentric systems

Open questions

If one constructs the in- & excenters of a tetrahedron (green) one finds, even if the tetrahedron is orthocentric, that its in-touch (red) and excentral (blue) tetrahedra:



- Are **not** similar to one another;
- Are not orthocentric, so their altitudes are **not** concurrent;
- And that the vertices of \overline{ABCD} are **not** the altitudes' feet on the facets of $\overline{A'B'C'D'}$.


But the green edges **do** intersect the blue edges in the vertices of an octahedron (**not** the medial)

that bisect the dihedral angles **and** span diagonals meeting at \bar{I} .



THE TRICK IS TO MAKE THE EXCENTRAL TETRAHEDRON BE ORTHOCENTRIC!

- ▶ One way to construct an orthocentric tetrahedron is to take a planar orthocentric system and lift its orthocenter out of the plane.
- ▶ I have also found formulae that take an orthocentric tetrahedron's parameters $D_{A'}, D_{B'}, D_{C'}, D_{D'}$ and return the vertices $\bar{A}, \bar{B}, \bar{C}, \bar{D}$ of the tetrahedron with it as its excentral tetrahedron, but ...



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra


Orthocentric
systems

Open
questions



THE TRICK IS TO MAKE THE EXCENTRAL TETRAHEDRON BE ORTHOCENTRIC!

- One way to construct an orthocentric tetrahedron is to take a planar orthocentric system and lift its orthocenter out of the plane.
- I have also found formulae that take an orthocentric tetrahedron's parameters $D_{A'}, D_{B'}, D_{C'}, D_{D'}$ and return the vertices $\bar{A}, \bar{B}, \bar{C}, \bar{D}$ of the tetrahedron with it as its excentral tetrahedron, but ...



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



THE TRICK IS TO MAKE THE EXCENTRAL TETRAHEDRON BE ORTHOCENTRIC!

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra


The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- One way to construct an orthocentric tetrahedron is to take a planar orthocentric system and lift its orthocenter out of the plane.
- I have also found formulae that take an orthocentric tetrahedron's parameters $D_{A'}, D_{B'}, D_{C'}, D_{D'}$ and return the vertices $\bar{A}, \bar{B}, \bar{C}, \bar{D}$ of the tetrahedron with it as its excentral tetrahedron, but ...



- for small lifts the incenter \bar{I} coincides with the lifted vertex \bar{D}' , while the new tetrahedron's vertex \bar{D} is inside of $\bar{A}'\bar{B}'\bar{C}'\bar{D}'$, and that is obviously not right!



ANALOGIES WITH TRIANGLES WHEN THE EX-CENTRAL TETRAHEDRON IS ORTHOCENTRIC

► Happily, on lifting $\overline{D'}$ a bit more, $\overline{O'}$ passes inside, \overline{ABCD} changes **discontinuously**, and everything pops miraculously into place.

► **First** and foremost, \overline{I} now coincides with $\overline{H'}$, and $\overline{D'}$ is outside of \overline{ABCD} .

Second, the edges of \overline{ABCD} & $\overline{A'B'C'D'}$ become orthogonal.

Third, the diagonals of the excentral octahedron also become orthogonal to its edges (but not to those of \overline{ABCD}).

Fourth, the Euler lines of $\overline{A'B'C'D'}$ & \overline{JKLM} (not shown) coincide and are parallel that of \overline{ABCD} (dashed lines).

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



ANALOGIES WITH TRIANGLES WHEN THE EX-CENTRAL TETRAHEDRON IS ORTHOCENTRIC

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions


► Happily, on lifting $\overline{D'}$ a bit more, $\overline{O'}$ passes inside, \overline{ABCD} changes **discontinuously**, and everything pops miraculously into place.

► **First** and foremost, \overline{I} now coincides with $\overline{H'}$, and $\overline{D'}$ is outside of \overline{ABCD} .

Second, the edges of \overline{ABCD} & $\overline{A'B'C'D'}$ become orthogonal.

Third, the diagonals of the excentral octahedron also become orthogonal to its edges (but not to those of \overline{ABCD}).

Fourth, the Euler lines of $\overline{A'B'C'D'}$ & \overline{JKLM} (not shown) coincide and are parallel that of \overline{ABCD} (dashed lines).





ANALOGIES WITH TRIANGLES WHEN THE EX-CENTRAL TETRAHEDRON IS ORTHOCENTRIC


► Happily, on lifting $\overline{D'}$ a bit more, $\overline{O'}$ passes inside, \overline{ABCD} changes **discontinuously**, and everything pops miraculously into place.

► **First** and foremost, \overline{I} now coincides with $\overline{H'}$, and $\overline{D'}$ is outside of \overline{ABCD} .

Second, the edges of \overline{ABCD} & $\overline{A'B'C'D'}$ become orthogonal.

Third, the diagonals of the excentral octahedron also become orthogonal to its edges (but not to those of \overline{ABCD}).

Fourth, the Euler lines of $\overline{A'B'C'D'}$ & \overline{JKLM} (not shown) coincide and are parallel that of \overline{ABCD} (dashed lines).



Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra

Orthocentric systems

Open questions



ANALOGIES WITH TRIANGLES WHEN THE EX-CENTRAL TETRAHEDRON IS ORTHOCENTRIC


► Happily, on lifting $\overline{D'}$ a bit more, $\overline{O'}$ passes inside, \overline{ABCD} changes **discontinuously**, and everything pops miraculously into place.

► **First** and foremost, \overline{I} now coincides with $\overline{H'}$, and $\overline{D'}$ is outside of \overline{ABCD} .

Second, the edges of \overline{ABCD} & $\overline{A'B'C'D'}$ become orthogonal.

Third, the **diagonals** of the excentral octahedron also become orthogonal to its edges (but not to those of \overline{ABCD}).

Fourth, the Euler lines of $\overline{A'B'C'D'}$ & \overline{JKLM} (not shown) coincide and are parallel that of \overline{ABCD} (dashed lines).



Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra

Orthocentric systems

Open questions



ANALOGIES WITH TRIANGLES WHEN THE EX-CENTRAL TETRAHEDRON IS ORTHOCENTRIC


► Happily, on lifting $\overline{D'}$ a bit more, $\overline{O'}$ passes inside, \overline{ABCD} changes **discontinuously**, and everything pops miraculously into place.

► **First** and foremost, \overline{I} now coincides with $\overline{H'}$, and $\overline{D'}$ is outside of \overline{ABCD} .

Second, the edges of \overline{ABCD} & $\overline{A'B'C'D'}$ become orthogonal.

Third, the **diagonals** of the excentral octahedron also become orthogonal to its edges (but not to those of \overline{ABCD}).

Fourth, the Euler lines of $\overline{A'B'C'D'}$ & \overline{JKLM} (not shown) coincide and are parallel that of \overline{ABCD} (dashed lines).



Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions




ANALOGIES WITH TRIANGLES WHEN THE EX-CENTRAL TETRAHEDRON IS ORTHOCENTRIC

Last but not least: The in-touch \overline{JKLM} and excentral $\overline{A'B'C'D'}$ tetrahedra are similar as in triangles, but:

➤ The scale factor is not $2R/r$ as it is in triangles, nor any simple rational multiple thereof.

➤ The homothetic center (orange) of \overline{JKLM} & $\overline{A'B'C'D'}$ is on their Euler line, but I haven't otherwise characterized it.

➤ And alas, conformal duality still does not work the way it does in the plane.



Heron's Formula in Orthocentric Tetrahedra
 Havel & Sobczyk
 Not-so elementary
 Heron's formula for triangles
 Heron's formula for tetrahedra
 The zeros' projective nature
 Orthocentric tetrahedra
 Orthocentric systems
 Open questions




ANALOGIES WITH TRIANGLES WHEN THE EX-CENTRAL TETRAHEDRON IS ORTHOCENTRIC

Last but not least: The in-touch \overline{JKLM} and excentral $\overline{A'B'C'D'}$ tetrahedra are similar as in triangles, but:

➤ The scale factor is not $2R/r$ as it is in triangles, nor any simple rational multiple thereof.

➤ The homothetic center (orange) of \overline{JKLM} & $\overline{A'B'C'D'}$ is on their Euler line, but I haven't otherwise characterized it.

➤ And alas, conformal duality still does not work the way it does in the plane.



Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra

Orthocentric systems

Open questions




ANALOGIES WITH TRIANGLES WHEN THE EX-CENTRAL TETRAHEDRON IS ORTHOCENTRIC

Last but not least: The in-touch \overline{JKLM} and excentral $\overline{A'B'C'D'}$ tetrahedra are similar as in triangles, but:

- The scale factor is not $2R/r$ as it is in triangles, nor any simple rational multiple thereof.
- The homothetic center (orange) of \overline{JKLM} & $\overline{A'B'C'D'}$ is on their Euler line, but I haven't otherwise characterized it.

➤ And alas, conformal duality still does not work the way it does in the plane.



Heron's Formula in Orthocentric Tetrahedra

Havel & Sobczyk

Not-so elementary

Heron's formula for triangles

Heron's formula for tetrahedra

The zeros' projective nature

Orthocentric tetrahedra

Orthocentric systems

Open questions



ANALOGIES WITH TRIANGLES WHEN THE EX-CENTRAL TETRAHEDRON IS ORTHOCENTRIC

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

Last but not least: The in-touch \overline{JKLM} and excentral $\overline{A'B'C'D'}$ tetrahedra are similar as in triangles, but:

- The scale factor is not $2R/r$ as it is in triangles, nor any simple rational multiple thereof.
- The homothetic center (orange) of \overline{JKLM} & $\overline{A'B'C'D'}$ is on their Euler line, but I haven't otherwise characterized it.
- And alas, conformal duality still does not work the way it does in the plane.






TABLE OF CONTENTS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- 1 Low-dimensional Euclidean geometry is not so elementary
- 2 Heron's formula and its relation to incenter geometry
- 3 Extending Heron's formula to tetrahedra (and beyond ...)
- 4 But its zeros are **really** weird in 3 (& higher?) dimensions
- 5 Orthocentric tetrahedra as extensions of triangles to 3D
- 6 Orthocentric systems in 2 & 3D, and in/excenter geometry
- 7 **Open Questions and Future Work**



OPEN QUESTIONS ABOUT THESE ORTHOCENTRIC ZEROS AND IN GENERAL

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- Because the (orthocentric) excentral tetrahedron depends on four parameters and uniquely determines the base tetrahedron, the latter also constitutes a four-parameter manifold of some kind: **can it be independently characterized?**
- Although the excentral tetrahedron can't be "squashed" into a plane (or line) without its circumcenter getting outside it, you can squash its base tetrahedron into a plane by sending one or more excentral vertices to infinity: **can you reach all the zeros with an orthocentric excentral tetrahedron?**
- There is no (finite) quadruple of null vectors in $\mathcal{G}_{4,1}$ that corresponds to the rank 2 zeros of Ω : **what is the "simplest" geometric algebra containing $\mathcal{G}_{4,1}$ that can represent all the zeros explicitly?**

On that last question: I suspect it is $\mathcal{G}_{4,2}$, the geometric algebra of **Lie sphere** (or "contact") geometry.



OPEN QUESTIONS ABOUT THESE ORTHOCENTRIC ZEROS AND IN GENERAL

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- Because the (orthocentric) excentral tetrahedron depends on four parameters and uniquely determines the base tetrahedron, the latter also constitutes a four-parameter manifold of some kind: **can it be independently characterized?**
- Although the excentral tetrahedron can't be "squashed" into a plane (or line) without its circumcenter getting outside it, you can squash its base tetrahedron into a plane by sending one or more excentral vertices to infinity: **can you reach all the zeros with an orthocentric excentral tetrahedron?**
- There is no (finite) quadruple of null vectors in $\mathcal{G}_{4,1}$ that corresponds to the rank 2 zeros of Ω : **what is the "simplest" geometric algebra containing $\mathcal{G}_{4,1}$ that can represent all the zeros explicitly?**

On that last question: I suspect it is $\mathcal{G}_{4,2}$, the geometric algebra of **Lie sphere** (or "contact") geometry.



OPEN QUESTIONS ABOUT THESE ORTHOCENTRIC ZEROS AND IN GENERAL

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- Because the (orthocentric) excentral tetrahedron depends on four parameters and uniquely determines the base tetrahedron, the latter also constitutes a four-parameter manifold of some kind: **can it be independently characterized?**
- Although the excentral tetrahedron can't be "squashed" into a plane (or line) without its circumcenter getting outside it, you can squash its base tetrahedron into a plane by sending one or more excentral vertices to infinity: **can you reach all the zeros with an orthocentric excentral tetrahedron?**
- There is no (finite) quadruple of null vectors in $\mathcal{G}_{4,1}$ that corresponds to the rank 2 zeros of Ω : **what is the "simplest" geometric algebra containing $\mathcal{G}_{4,1}$ that can represent all the zeros explicitly?**

On that last question: I suspect it is $\mathcal{G}_{4,2}$, the geometric algebra of **Lie sphere** (or "contact") geometry.



OPEN QUESTIONS ABOUT THESE ORTHOCENTRIC ZEROS AND IN GENERAL

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions

- Because the (orthocentric) excentral tetrahedron depends on four parameters and uniquely determines the base tetrahedron, the latter also constitutes a four-parameter manifold of some kind: **can it be independently characterized?**
- Although the excentral tetrahedron can't be "squashed" into a plane (or line) without its circumcenter getting outside it, you can squash its base tetrahedron into a plane by sending one or more excentral vertices to infinity: **can you reach all the zeros with an orthocentric excentral tetrahedron?**
- There is no (finite) quadruple of null vectors in $\mathcal{G}_{4,1}$ that corresponds to the rank 2 zeros of Ω : **what is the "simplest" geometric algebra containing $\mathcal{G}_{4,1}$ that can represent all the zeros explicitly?**

On that last question: I suspect it is $\mathcal{G}_{4,2}$, the geometric algebra of **Lie sphere** (or "contact") geometry.



A COUPLE OF ACKNOWLEDGMENTS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



I would first like to acknowledge my co-author Garret Sobczyk (who was unable to attend for personal reasons), and in particular for the matrix representation of $\mathcal{G}_{4,1}$ that I used to validate all my calculations. As we all know, his contributions to the field of geometric algebra are second to none!



A COUPLE OF ACKNOWLEDGMENTS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



I would first like to acknowledge my co-author Garret Sobczyk (who was unable to attend for personal reasons), and in particular for the matrix representation of $\mathcal{G}_{4,1}$ that I used to validate all my calculations. As we all know, his contributions to the field of geometric algebra are second to none!

And since we're in The Netherlands, I'd also like to acknowledge Johan Jacob (Jaap) Seidel (1919-2001), whose remarks during my circa 1985 visit to him in Eindhoven eventually led me to see the connection between Hestenes' work on $\mathcal{G}_{n+1,1}$ and distance geometry.



A COUPLE OF ACKNOWLEDGMENTS

Heron's
Formula in
Orthocentric
Tetrahedra

Havel &
Sobczyk

Not-so
elementary

Heron's
formula for
triangles

Heron's
formula for
tetrahedra

The zeros'
projective
nature

Orthocentric
tetrahedra

Orthocentric
systems

Open
questions



I also thank the GeoGebra team for the software used to generate all the pretty pictures in this talk.

And thank YOU for your attention!

(handouts online at:
<http://dx.doi.org/10.13140/RG.2.2.17531.94240/2>)