

More on Fourier integrals

note by T. H. Koornwinder, thk@science.uva.nl

written for the course on Linear analysis, September–December 2005,
University of Amsterdam

last modified: January 4, 2006

This note gives a minor extension to Chap. 5 of the book *Fourier analysis, an introduction* by E. M. Stein and R. Shakarchi.

Dense subspaces of $L^p(\mathbb{R})$

By a *simple function* on \mathbb{R} we mean a finite linear combination of characteristic functions χ_E of measurable subsets E of \mathbb{R} . In particular, a simple function on \mathbb{R} is integrable iff it is a finite linear combination of characteristic functions χ_E with $\lambda(E) < \infty$.

The Lebesgue measure λ on \mathbb{R} is *regular*, i.e., for every measurable set $E \subset \mathbb{R}$ we have:

$$\begin{aligned}\lambda(E) &= \inf\{\lambda(V) : E \subset V \text{ and } V \text{ open}\}, \\ \lambda(E) &= \sup\{\lambda(K) : K \subset E \text{ and } K \text{ compact}\}.\end{aligned}$$

This follows from Theorem 2.18 in Rudin, *Real and complex analysis*.

Proposition Let $1 \leq p < \infty$. The following spaces are dense in $L^p(\mathbb{R})$:

1. The space of integrable simple functions on \mathbb{R} .
2. The linear span of the characteristic functions of bounded intervals in \mathbb{R} .
3. The space of continuous functions on \mathbb{R} of *compact support*, i.e., which vanish outside some bounded interval.

Proof We will prove these results for $p = 1$. The proof for other p is similar.

Proof of 1. Every $f \in L^1(\mathbb{R})$ can be written as $f = f_1 - f_2 + if_3 - if_4$ with f_1, f_2, f_3, f_4 nonnegative L^1 functions. So it is sufficient to prove that every nonnegative L^1 function f can be approximated in L^1 norm by integrable simple functions. There is an increasing sequence of nonnegative simple functions $t_n(x)$ which tend pointwise to f as $n \rightarrow \infty$. Then $\int t_n$ tends to $\int f$ as $n \rightarrow \infty$, so $\|f - t_n\|_1 \rightarrow 0$.

Proof of 2. By 1. it is sufficient to prove that, if $E \subset \mathbb{R}$ is measurable with $\lambda(E) < \infty$ then χ_E can be approximated in L^1 norm by finite linear combinations of characteristic functions of bounded intervals. Let $\varepsilon > 0$. By regularity of λ there is an open set $V \supset E$ such that $\lambda(V) < \lambda(E) + \frac{1}{2}\varepsilon < \infty$. Since V is a countable disjoint union of open intervals, there is a finite union $W \subset V$ of bounded open intervals such that $\lambda(W) > \lambda(V) - \frac{1}{2}\varepsilon$. Hence $\|\chi_E - \chi_W\| < \varepsilon$.

Proof of 3. Every characteristic function of a bounded interval can be approximated in L^1 norm by continuous functions of compact support. Now use 2. \square

We can use part 2. of this Proposition in order to prove the *Riemann-Lebesgue Lemma* for the Fourier transform:

If $f \in L^1(\mathbb{R})$ then $\widehat{f}(\xi) \rightarrow 0$ as $\xi \rightarrow \pm\infty$.

Just observe that the statement is true for $f = \chi_{[a,b]}$.

Exercises

Vrst 1. (For this exercise use results from both Fourier series and Fourier integrals.)
Below define $x^{-1} \sin x$ for $x = 0$ by continuity.

a) Let $t \in \mathbb{R}$. Show that for each $x \in (-\pi, \pi)$ we have

$$\sum_{n=-\infty}^{\infty} \frac{\sin(\pi(t-n))}{\pi(t-n)} e^{inx} = e^{ixt}$$

with pointwise convergence. What is the evaluation of the sum on the left-hand side for other real values of x ?

b) Show that, for all $n, m \in \mathbb{Z}$, we have

$$\int_{-\infty}^{\infty} \frac{\sin(\pi(t-n))}{\pi(t-n)} \frac{\sin(\pi(t-m))}{\pi(t-m)} dt = \delta_{n,m},$$

where the integral converges absolutely.

c) Does there exist $f \in L^2(\mathbb{R})$ with $f \neq 0$ such that

$$\int_{-\infty}^{\infty} f(t) \frac{\sin(\pi(t-n))}{\pi(t-n)} dt = 0 \quad \text{for all } n \in \mathbb{Z} ?$$

d) Let $f \in L^2([-\pi, \pi])$. Define \hat{f} as a function on \mathbb{R} by

$$\hat{f}(t) := \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-ixt} dx \quad (t \in \mathbb{R}). \quad (1)$$

(For $t \in \mathbb{Z}$ this defines the Fourier coefficients of f ; for general $t \in \mathbb{R}$ this defines the Fourier transform of a function on \mathbb{R} which vanishes outside $[-\pi, \pi]$.) Show that

$$\hat{f}(t) = \sum_{n=-\infty}^{\infty} \hat{f}(n) \frac{\sin(\pi(t-n))}{\pi(t-n)} \quad (t \in \mathbb{R}) \quad (2)$$

with absolutely convergent sum.

(This shows in particular the following. Let g be an L^2 function on \mathbb{R} which is the Fourier transform $g = \hat{f}$ of an L^2 function f on \mathbb{R} vanishing outside $[-\pi, \pi]$ (see (1)). So g is also continuous. Then g is completely determined by its restriction to \mathbb{Z} , with reconstruction formula given by (2).)

Hints to Problem 7 in Chapter 5 of Stein & Shakarchi

A result from (a) is:

$$h_k(x) = (-1)^k e^{x^2/2} \left(\frac{d}{dx}\right)^k e^{-x^2}.$$

Show, by once differentiating this formula, that

$$h_{k+1}(x) = \left(x - \frac{d}{dx}\right) h_k(x). \quad (3)$$

Use (3) in (c) in order to prove the result there by induction with respect to k .

Now show, by applying the operator $x + \frac{d}{dx}$ to both sides of (3) and by using induction with respect to k , that

$$\left(x + \frac{d}{dx}\right) h_{k+1}(x) = 2(k+1)h_k(x). \quad (4)$$

Now it has to be proved in (d) that

$$(Lh_k)(x) := \left(x^2 - \frac{d^2}{dx^2}\right) h_k(x) = (2k+1)h_k(x).$$

Show this by expressing the operator L in terms of $x - \frac{d}{dx}$ and $x + \frac{d}{dx}$ and by using (3) and (4).